MINISTRY OF EDUCATION AND SCIENCE
RUSSIAN FEDERATION
STATE EDUCATIONAL INSTITUTION
OF HIGHER PROFESSIONAL EDUCATION
BASHKIR STATE UNIVERSITY

arXiv:1110.4737v3 [math-ph] 10 May 2012

UDK 514.744.2

K.V. Andreev

On the spinor formalism for even n.

UFA-2011

I

English edition

Table of Contents

1 Introduction **5**

2 Clifford algebra **6**

3 Complex and real representations **9**

4 Real inclusion. Involution **14**

5 Elementary transformations of the orthogonal group **16**

6 Clifford equation. Double covering **24**

7 Quaternions **35**

8 Particular solutions of the Clifford equation **37**

9 On the structural constants of the non-division algebra. Sedenions **47**

10 Infinitesimal transformations **52**

11 Complex and real connections **53**

12 On the classification of tensors with the symmetries of the Riemann curvature tensor. Curvature spinors **61**

13 Twistor equation **65**

14 Spinor formalism for n=6 and n=8 **68**

14.1 General isomorphisms 68

14.2 On the classification of tensors with the symmetries of the Riemann curvature tensor for n=6. Curvature spinors 69

14.3 Geometric representation of a twistor in $\mathbb{R}^6_{(2,4)}$ 72

14.4 Structure constants of the octonion algebra. Spinor formalism for n = 8 77

14.5 Geometry of the inductive step from n=6 to n=8 79

14.5.1 Rosenfeld null-pair 79

14.5.2 Construction of the quadrics $\mathbb{C}Q_6$ and $\mathbb{C}\tilde{Q}_6$ 79

14.6 Correspondence $\mathbb{C}Q_6 \longmapsto \mathbb{C}\tilde{Q}_6$ 80
  14.6.1 Geometry of the transition to the connecting operators $\eta_\Lambda{}^{KL}$ for n = 8 83
14.7 Correspondence $\mathbb{C}\tilde{Q}_6 \longmapsto \mathbb{C}Q_6$ 87
  14.7.1 On the Majorana spinors. 89
  14.7.2 Theorem on two quadrics. 91
14.8 Explicit construction of the Lie spin-operators for n=4. 93

15 Acknowledgments 98

16 Appendix 99
16.1 Proof of the metric tensor and involution relations 99
16.2 Proof of the connecting operator relations 101
16.3 Proof of the infinitesimal transformation relations 103
16.4 Proof of the Lie bracket identity 104
16.5 Proof of the curvature tensor relations 105
16.6 Proof of the auxiliary identities for the theorem on the spin analogs of the curvature tensor 105
16.7 Proof of the twistor equation relations 107
16.8 Proof of the identities for n = 6 and n=8 109
  16.8.1 Proof of the fundamental identities for n = 6 109
  16.8.2 Proof of the 4-vector identities 110
  16.8.3 Proof of the 6-vector identities 110
  16.8.4 Proof of the Bianchi identity 112
  16.8.5 Proof of the identities related to the Weyl tensor 112
  16.8.6 Proof of the relations associated with the metric induced in the cross section of the cone K_6 113
  16.8.7 Proof of the first invariant relations 114
  16.8.8 Proof of the second invariant relations 116
  16.8.9 Proof of the octave alternative identity 117
  16.8.10 Proof of the central Moufang identity 118
  16.8.11 Proof of the A-operator identities for n = 8 119

List of Tables

---

1 The matrix form of the tensor S for the real inclusions of some 2-dimensional spaces in the complex space for $r^i = (T, Z)$. 45
2 The matrix form of the tensor S for the real inclusions of some 4-dimensional spaces in the complex space for $r^i = (T, X, Y, Z)$. 45
3 The matrix form of the tensor S for the real inclusions of some 6-dimensional spaces in the complex space for $r^i = (T, X, V, W, Y, Z)$. 46

List of Figures

1 Correspondence $\forall CP_2 \subset CP_3 \leftrightarrow R$ 82

2 Correspondence $CP_3 \supset CP_1 \leftrightarrow CP_1 \subset K_6$ 82

1 Introduction

In this article the answers to the following questions are considered for any even $n \geq 4$:

1. What is a Clifford algebra and how to construct it?
2. What is the real and complex representations?
3. What is an involution, and how it helps in a transition to real inclusions?
4. How to construct the complex and real representations of the Clifford generators (the connecting operators)?
5. How does the involution act on the Clifford generators (the connecting operators)?
6. How to lead a complex orthogonal matrix to a block diagonal form?
7. How to construct the basic isomorphisms (including the double coverings) and other relations in the explicit form with the help of the connecting operators?
8. How to construct a partial solution of the Clifford equation for the connecting operators for even $n \geq 4$?
9. How to construct the structure constants of the hypercomplex (sedenion) algebra (without division for n>8) by means of the connecting operators for $n \mod 8 = 0$?
10. How to enter and to coordinate the connection in the tangent and spinor bundles?
11. How to construct the Lie operator analogues and what are the conditions of their construction?
12. How to construct the curvature spinors?
13. What is the communication between the twistor equation, the derivational equation of the normalized Grassmannian and the conformal Killing equation?
14. What is the difference between the spinor formalism for $n \leq 8$ and the one for $n > 8$, and how to construct it for small dimensions?
15. How to construct the geometric representation of spinors (twistors) for $\mathbb{R}^6_{(2,4)}$?
16. How to construct a generalization of the Cartan triality principle to the Klein correspondence and what is the geometry of such the generalization for $n = 8$?
17. How to construct (one-to-one) the spin (spin-pair) analogues of the Lie operators for $n = 4$ in an explicit form?

The conclusion of all results of this article is made on the basis [33].[1]

---

[1] The Russian edition (on the pp. 124-261) contains the original variant of the article **with the corrected errors** and the original numbering of the pages (pp. 1-138)

2 Clifford algebra

In this section, it is told about how to construct the real 2n-dimensional Clifford algebra with any functional G according to [18, lecture 13, pp. 258-299]. In the case when the functional G in a suitable basis has the diagonal form with the identical quantity $\ll + \gg$ and $\ll - \gg$ on the main diagonal, the generators of such the Clifford algebra create the generators of the complex n-dimensional Clifford algebra which is isomorphic to the algebra of complex matrixes $\mathbb{C}(2^{\frac{n}{2}})$. The conclusion of all results of this section is made on the basis [18].

Let $\mathcal{V}$ be a vector space over $\mathbb{R}$. Then $\mathcal{V}$ will be a module over $\mathbb{R}$: 1). a (x+y) =ax+ay, 2). (a+b) x=ax+bx, 3). (ab) x=a (bx), 4). 1· x=x, where x, y $\in$ $\mathcal{V}$, a, b $\in$ $\mathbb{R}$. Let's consider the ideal

$$J(G) := \{T \otimes (x \otimes x - \frac{1}{2}G(x));\ T \in T_0(\mathcal{V}), x \in {T_0}^1(\mathcal{V})\} \tag{2.1}$$

in the algebra $T_0(\mathcal{V}) = {T_0}^0(\mathcal{V}) \oplus ... \oplus {T_0}^q(\mathcal{V}) \oplus ....$ Also we will define the Clifford algebra as $CL(G^{2n}) := T_0(\mathcal{V})/J(G)$. We will construct a representation of such the algebra into an polynomial algebra

$$\begin{array}{ll} \alpha : \mathcal{V} \longmapsto \mathcal{A}, & \tilde{\alpha} : T_0(\mathcal{V}) \longmapsto \mathcal{A}, \\ \alpha(\mathcal{V}) = B^{\tilde{\Lambda}} x_{\tilde{\Lambda}}, & \tilde{\alpha}(T_0(\mathcal{V})) = A + B^{\tilde{\Lambda}} x_{\tilde{\Lambda}} + C^{\tilde{\Lambda}\tilde{\Psi}} x_{\tilde{\Lambda}} x_{\tilde{\Psi}} + ... \ . \end{array} \tag{2.2}$$

Hereinafter $\tilde{\Lambda},\quad \tilde{\Psi},\quad ... = \overline{1,2n}$. Suppose that the relation $\hat{\alpha}(J(G)) = 0$ is executed for some mapping $\hat{\alpha} : CL(G) \longmapsto C\mathcal{A}$. It will mean that

$$x_{\tilde{\Lambda}} x_{\tilde{\Psi}} + x_{\tilde{\Psi}} x_{\tilde{\Lambda}} = G(x_{\tilde{\Lambda}}, x_{\tilde{\Psi}}). \tag{2.3}$$

Therefore, the form $G(x_{\tilde{\Lambda}}, x_{\tilde{\Psi}})$ in a suitable basis has the diagonal form. Suppose that for such the basis, the parities

$$x_{\tilde{\Lambda}} x_{\tilde{\Psi}} + x_{\tilde{\Psi}} x_{\tilde{\Lambda}} = 0,\ \ (\tilde{\Lambda} \neq \tilde{\Psi});\ \ {x_{\tilde{\Lambda}}}^2 = \pm 1 \tag{2.4}$$

are carried out that will define the mapping

$$\hat{\alpha}(Cl(G^{2n})) = A + B^{\tilde{\Lambda}} x_{\tilde{\Lambda}} + C^{[\tilde{\Lambda}\tilde{\Psi}]} x_{\tilde{\Lambda}} x_{\tilde{\Psi}} + ... \tag{2.5}$$

In turn, it means that the Clifford algebra is *finite-dimensional* and $dim\ CL(G^{2n}) = 2^{2n},\ \ dim\ \mathcal{V} = 2n$. Let's construct now the complex Clifford algebra $CL(g^n)$. For this purpose, we will consider the real Clifford algebra $CL(G^{2n}_{(n,n)})$. We will demand that $(x_{2\Lambda})^2 = 1,\ \ (x_{2\Lambda-1})^2 = -1$ and set

$$\sqrt{2} z_\Lambda = x_{2\Lambda} + i x_{2\Lambda-1},\ \ \sqrt{2}\bar{z}_\Lambda = x_{2\Lambda} - i x_{2\Lambda-1}. \tag{2.6}$$

Hereinafter $\Lambda,\quad \Psi,\quad ... = \overline{1,n}$. Then

$$\begin{array}{rcl} 2(z_\Lambda z_\Psi + z_\Psi z_\Lambda) & = & (x_{2\Lambda} + i x_{2\Lambda-1})(x_{2\Psi} + i x_{2\Psi-1}) + (x_{2\Psi} + i x_{2\Psi-1})(x_{2\Lambda} + i x_{2\Lambda-1}) = \\ & = & x_{2\Lambda} x_{2\Psi} - x_{2\Lambda-1} x_{2\Psi-1} + i(x_{2\Lambda-1} x_{2\Psi} + x_{2\Lambda} x_{2\Psi-1}) + \\ & + & x_{2\Psi} x_{2\Lambda} - x_{2\Psi-1} x_{2\Lambda-1} + i(x_{2\Psi-1} x_{2\Lambda} + x_{2\Psi} x_{2\Lambda-1}) = \\ & & = \begin{cases} 4, & \Lambda = \Psi, \\ 0, & \Lambda \neq \Psi, \end{cases} \\ 2(z_\Lambda \bar{z}_\Psi + \bar{z}_\Psi z_\Lambda) & = & (x_{2\Lambda} + i x_{2\Lambda-1})(x_{2\Psi} - i x_{2\Psi-1}) + (x_{2\Psi} - i x_{2\Psi-1})(x_{2\Lambda} + i x_{2\Lambda-1}) = \\ & = & x_{2\Lambda} x_{2\Psi} + x_{2\Lambda-1} x_{2\Psi-1} - i(-x_{2\Lambda-1} x_{2\Psi} + x_{2\Lambda} x_{2\Psi-1}) + \\ & + & x_{2\Psi} x_{2\Lambda} + x_{2\Psi-1} x_{2\Lambda-1} - i(x_{2\Psi-1} x_{2\Lambda} - x_{2\Psi} x_{2\Lambda-1}) = 0, \end{array} \tag{2.7}$$

$$z_\Lambda z_\Psi + z_\Psi z_\Lambda = g(z_\Lambda, z_\Psi). \tag{2.8}$$

Therefore, the factorization of the complex algebra $T_0(\mathcal{V}^{\mathbb{C}}) = T_0(\mathcal{V}^{\mathbb{C}})(z)$ by the ideal $J(g) = \{T \otimes (z \otimes z - \frac{1}{2}g(z));\quad T \in T_0(\mathcal{V}^{\mathbb{C}}),\quad z \in {T_0}^1(\mathcal{V}^{\mathbb{C}})\}$ will define the complex algebra $CL(g^n) \cong T_0(\mathcal{V}^{\mathbb{C}})/J(g)$.

Theorem 2.1. *For the Clifford algebra, there is the decomposition*

$$\begin{gathered}
CL(G^{(2n+2)}_{(n+1,n+1)}) = CL(G^{2n}_{(n,n)}) \otimes CL(G^2_{(1,1)}),\\
CL(G^{(2n+2)}_{(2n+2,0)}) = CL(G^{2n}_{(2n,0)}) \otimes CL(G^2_{(2,0)}),\quad CL(g^n) = CL(g^{n-2}) \otimes CL(g^2)
\end{gathered} \tag{2.9}$$

with the multiplication $a \cdot b = (\tilde{a},\ \tilde{\tilde{a}}) \cdot (\tilde{b},\ \tilde{\tilde{b}}) = (\tilde{a}\tilde{b},\ \tilde{\tilde{a}}\tilde{\tilde{b}})$.

Proof. Let $x_{\tilde{\Lambda}}$ be a basis in $\mathcal{V}^{2n}$, and let x_{2n+1}, x_{2n+2} be a basis in $\mathcal{V}^2$

$$\begin{gathered}
x_{\tilde{\Lambda}} x_{\tilde{\Psi}} + x_{\tilde{\Psi}} x_{\tilde{\Lambda}} = 0,\ \ (\tilde{\Lambda} \neq \tilde{\Psi});\quad (x_{2\Lambda})^2 = 1;\ \ (x_{2\Lambda-1})^2 = -1,\\
x_{2n+1}x_{2n+2} + x_{2n+2}x_{2n+1} = 0;\quad (x_{2n+2})^2 = -1;\ \ (x_{2n+1})^2 = 1.
\end{gathered} \tag{2.10}$$

It will give a chance to define the basis in $\mathcal{V}^{2n+2}$ as

$$X_{\tilde{\Lambda}} = x_{\tilde{\Lambda}} x_{2n+1} x_{2n+2},\quad X_{2n+1} = x_{2n+2},\quad X_{2n+2} = x_{2n+1} \tag{2.11}$$

with the conditions

$$\begin{gathered}
X_{\tilde{\Lambda}} X_{\tilde{\Psi}} + X_{\tilde{\Psi}} X_{\tilde{\Lambda}} = 0,\ \ (\tilde{\Lambda} \neq \tilde{\Psi}),\ \ (X_{2\Lambda})^2 = 1,\ \ (X_{2\Lambda-1})^2 = -1,\\
\text{in this equation } \Lambda = \overline{1, n+1},\quad \tilde{\Lambda} = \overline{1, 2n+2}.
\end{gathered} \tag{2.12}$$

Then for any $a \in CL(G^{2n+2}_{(n+1,n+1)})$, the decomposition

$$\begin{gathered}
a = a_1 X_{2n+2} X_{2n+1} + a_2 X_{2n+2} + a_3 X_{2n+1} + a_4 =\\
= a_1 \otimes x_{2n+1}x_{2n+2} + a_2 \otimes x_{2n+1} + a_3 \otimes x_{2n+2} + a_4 \otimes 1
\end{gathered} \tag{2.13}$$

is executed. Let's spend the replacement of the basis

$$\begin{gathered}
y_{2n+1} = (x_{2n+1} + x_{2n+2})/2,\quad y_{2n+2} = (x_{2n+1} - x_{2n+2})/2,\\
(y_{2n+1})^2 = 0,\quad (y_{2n+2})^2 = 0,\quad 2y_{2n+1}y_{2n+2} = 1 - x_{2n+1}x_{2n+2}.
\end{gathered} \tag{2.14}$$

Then, the decomposition (2.13) will have the form

$$a = (-a_1 + a_4)y_{2n+1}y_{2n+2} + (a_2 + a_3)y_{2n+1} + (a_2 - a_3)y_{2n+2} + (a_1 + a_4)y_{2n+2}y_{2n+1}. \tag{2.15}$$

Hence, any element $a \in CL(G^2_{(1,1)})$ is decomposed as

$$\begin{gathered}
a = a_1 x_{2n+1}x_{2n+2} + a_2 x_{2n+1} + a_3 x_{2n+2} + a_4 =\\
= \begin{pmatrix} y_{2n+1}y_{2n+2} & y_{2n+2} \end{pmatrix} \begin{pmatrix} -a_1 + a_4 & a_2 + a_3 \\ a_2 - a_3 & a_1 + a_4 \end{pmatrix} \begin{pmatrix} y_{2n+1}y_{2n+2} \\ y_{2n+1} \end{pmatrix},
\end{gathered} \tag{2.16}$$

and for any two elements $a, b \in CL(G^{2n+2}_{(n+1,n+1)})$, the relation

$$\begin{aligned}
c := a \cdot b \quad = \quad & (a_1 \otimes x_{2n+1}x_{2n+2} \ + \ a_2 \otimes x_{2n+1} \ + \ a_3 \otimes x_{2n+2} \ + \ a_4 \otimes 1)\cdot\\
& \cdot(b_1 \otimes x_{2n+1}x_{2n+2} \ + \ b_2 \otimes x_{2n+1} \ + \ b_3 \otimes x_{2n+2} \ + \ b_4 \otimes 1) =
\end{aligned}$$

$$= \underbrace{(a_2b_3 - a_3b_2 + a_1b_4 + a_4b_1)}_{:=c_1} \otimes x_{2n+1}x_{2n+2} \quad + \quad \underbrace{(a_2b_4 + a_4b_2 - a_1b_3 + a_3b_1)}_{:=c_2} \otimes x_{2n+1} +$$
$$+ \underbrace{(a_3b_4 + a_4b_3 - a_1b_2 + a_2b_1)}_{:=c_3} \otimes x_{2n+2} \quad + \quad \underbrace{(a_1b_1 + a_4b_4 + a_2b_2 - a_3b_3)}_{:=c_4} \tag{2.17}$$

is executed. At the same time, for any two elements $a, b \in CL(G^2_{(1,1)})$, the identity

$$c := a \cdot b = \begin{pmatrix} -a_1 + a_4 & a_2 + a_3 \\ a_2 - a_3 & a_1 + a_4 \end{pmatrix} \cdot \begin{pmatrix} -b_1 + b_4 & b_2 + b_3 \\ b_2 - b_3 & b_1 + b_4 \end{pmatrix} =$$
$$= \begin{pmatrix} a_1b_1 - a_1b_4 - a_4b_1 + a_4b_4 + a_2b_2 + a_3b_2 - a_2b_3 - a_3b_3 \\ -a_2b_1 + a_3b_1 + a_2b_4 - a_3b_4 + a_1b_2 + a_4b_2 - a_1b_3 - a_4b_3 \end{pmatrix.$$
$$\begin{matrix} -a_1b_2 + a_4b_2 - a_1b_3 + a_4b_3 + a_2b_1 + a_3b_1 + a_2b_4 + a_3b_4 \\ a_2b_2 - a_3b_2 + a_2b_3 - a_3b_3 + a_1b_1 + a_4b_1 + a_1b_4 + a_4b_4 \end{pmatrix} =$$
$$= \begin{pmatrix} -c_1 + c_4 & c_2 + c_3 \\ c_2 - c_3 & c_1 + c_4 \end{pmatrix}. \tag{2.18}$$

is true. That will allow to define the correspondence between the generators of the Clifford algebra $CL(G^2_{(1,1)})$ and their matrix representations

$$x_{2n+1} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad x_{2n+2} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \tag{2.19}$$

This means that for any two elements $a, b \in CL(G^{2n+2}_{(n+1,n+1)})$, for the similar matrix representation, $c_1, c_2, c_3, c_4 \in CL(G^{2n}_{(n,n)})$. Other two representations are similarly proved also. ☐

Theorem 2.2. *The complex Clifford algebra for even n is isomorphic to the algebra of complex matrixes of kind*

$$CL(g) \cong \mathbb{C}(2^{n/2}). \tag{2.20}$$

Proof. Let's determine according to the proof of Theorem 2.1 and replacing X on x

$$\begin{gathered} \sqrt{2}z_{n+1} = x_{2n+2} + ix_{2n+1}, \quad \sqrt{2}z_{n+2} = x_{2n+4} + ix_{2n+3}, \\ z_{n+1}z_{n+1} = 1, \quad z_{n+1}\bar{z}_{n+1} = ix_{2n+1}x_{2n+2} = -\bar{z}_{n+1}z_{n+1}, \\ z_{n+2}z_{n+2} = 1, \quad z_{n+2}\bar{z}_{n+2} = ix_{2n+3}x_{2n+4} = -\bar{z}_{n+2}z_{n+2}, \\ z_{n+1}z_{n+2} + z_{n+2}z_{n+1} = 0. \end{gathered} \tag{2.21}$$

This will allow to define the correspondence between the generators of the Clifford algebra $CL(G^4_{(2,2)})$ and their matrix representations

$$\begin{gathered} x_{2n+2} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad x_{2n+1} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \\ x_{2n+4} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \quad x_{2n+3} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}. \end{gathered} \tag{2.22}$$

Then for the generators $CL(g^2)$ in the special basis similar to the one of Example 3.2[2], we will obtain

$$\sqrt{2}z_{n+1} = \begin{pmatrix} 0 & 1+i \\ 1-i & 0 \end{pmatrix}, \quad \sqrt{2}z_{n+2} = \begin{pmatrix} 0 & 1-i \\ 1+i & 0 \end{pmatrix}, \tag{2.23}$$

as will prove our theorem. Strictly to construct the specified conformity in the common view, the knowledge of some additional facts, which are worthy to devote them the following section, is necessary. □

Thus, the real Clifford algebra $CL(G^{2n}_{(n,n)})$ is isomorphic to the real algebra of n-dimension matrixes

$$CL(G^{2n}_{(n,n)}) \cong \mathbb{R}(2^n). \tag{2.24}$$

Therefore, there is a mapping

$$\gamma: \quad \mathbb{R}(2^n) \longmapsto (\mathcal{V}^{\mathbb{R}})^{2n}. \tag{2.25}$$

On the basis (2.3), the operators $\gamma_{\tilde{\Lambda}}$ should satisfy the equation

$$\gamma_{\tilde{\Lambda}}\gamma_{\tilde{\Psi}} + \gamma_{\tilde{\Psi}}\gamma_{\tilde{\Lambda}} = G_{\tilde{\Lambda}\tilde{\Psi}}, \tag{2.26}$$

which is called *Clifford equation.* At the same time, the complex Clifford algebra is represented by the matrix algebra $\mathbb{C}(2^{n/2})$. The multiplication in the complex Clifford algebra delegates the matrix multiplication in $\mathbb{C}(2^{n/2})$. Therefore, one can construct mapping

$$\gamma: \quad \mathbb{C}(2^{n/2}) \longmapsto (\mathcal{V}^{\mathbb{C}})^{n}. \tag{2.27}$$

From (2.8), it follows that the operators γ_Λ should satisfy the complex Clifford equation

$$\gamma_\Lambda\gamma_\Psi + \gamma_\Psi\gamma_\Lambda = g_{\Lambda\Psi}. \tag{2.28}$$

3 Complex and real representations

In this section, it is told about how locally to construct the n-dimensional complex representation on the real one of the 2n-dimensional pseudo-Riemannian manifold with tangent bundle fibers isomorphic to $\mathbb{R}^{2n}_{(n,n)}$. For this purpose, we construct the corresponding complex reparametrization of an atlas in a neighborhood of some point that induces the Neifeld operators [15] connected to the Norden affinor [17] in the tangent bundle. Accordingly, by means of the connecting operators $\gamma_{\tilde{\Lambda}}$, the Neifeld operator analogues are induced to within a sign in the spinor bundle. It will give a chance to pass to the complex representation of the connecting operators γ_Λ. The conclusion of all results of this section is made on the basis [15], [17].

Under *complex analytical Riemannian space* $\mathbb{C}V_n$, we will further understand an analytical complex bundle supplied with *analytical quadratic metric*, i.e., a metric defined by means of a symmetric nonsingular tensor $g_{\Lambda\Psi}$ (here $\Lambda, \Psi, ... = \overline{1,n}$; $\tilde{\Lambda}, \tilde{\Psi}, ... = \overline{1,2n}$), which the coordinates are analytical functions of the point coordinates. To this tensor, there corresponds *complex Riemannian torsion-free connection*, the coefficients of which are defined by the Christoffel symbols, and hence these coefficients are analytical functions. The tangent bundle to this manifold $\tau^{\mathbb{C}}(\mathbb{C}V_n)$ has fibers $\tau_x^{\mathbb{C}} \cong \mathbb{C}\mathbb{R}^n$ that is fibers isomorphic to the n-dimensional complex Euclidian space, the metric of which is defined by the value of the metric tensor at the given point. The real representation V_{2n} of $\mathbb{C}V_n$ has the tangent bundle $\tau^{\mathbb{R}}(V_{2n})$ with fibers isomorphic to $\mathbb{R}^{2n}_{(n,n)}$. Let on V_{2n}, an atlas $(U; x^{\tilde{\Lambda}})$ be set. We will consider *reparametrization* of this atlas $(U; w^{\Lambda})$ such that $w^{\Lambda} = \frac{1}{\sqrt{2}}(u^{\Lambda}(x^{\tilde{\Lambda}}) + iv^{\Lambda}(x^{\tilde{\Lambda}}))$, which is locally solvable as $x^{\tilde{\Lambda}} = x^{\tilde{\Lambda}}(u^{\Lambda}, v^{\Lambda})$. Set

---

[2] In the example, $x_{\tilde{\Lambda}} \longmapsto \frac{1}{\sqrt{2}}x_{\tilde{\Lambda}}$, $z_\Lambda \longmapsto \frac{1}{\sqrt{2}}z_\Lambda$.

$$m^{\Lambda}{}_{\tilde{\Lambda}} := \frac{1}{\sqrt{2}}(\frac{\partial u^{\Lambda}}{\partial x^{\tilde{\Lambda}}} + i\frac{\partial v^{\Lambda}}{\partial x^{\tilde{\Lambda}}}) =: \frac{\partial w^{\Lambda}}{\partial x^{\tilde{\Lambda}}}, \quad m_{\Lambda}{}^{\tilde{\Lambda}} := \frac{1}{\sqrt{2}}(\frac{\partial x^{\tilde{\Lambda}}}{\partial u^{\Lambda}} - i\frac{\partial x^{\tilde{\Lambda}}}{\partial v^{\Lambda}}) =: \frac{\partial x^{\tilde{\Lambda}}}{\partial w^{\Lambda}}. \tag{3.1}$$

Then

$$\Delta_{\tilde{\Lambda}}{}^{\tilde{\Psi}} := m^{\Lambda}{}_{\tilde{\Lambda}} m_{\Lambda}{}^{\tilde{\Psi}} = \frac{1}{2}((\underbrace{\frac{\partial u^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ \frac{\partial x^{\tilde{\Psi}}}{\partial u^{\Lambda}} + \frac{\partial v^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ \frac{\partial x^{\tilde{\Psi}}}{\partial v^{\Lambda}}}_{=\delta_{\tilde{\Lambda}}{}^{\tilde{\Psi}}}) + i(\underbrace{\frac{\partial v^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ \frac{\partial x^{\tilde{\Psi}}}{\partial u^{\Lambda}} - \frac{\partial u^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ \frac{\partial x^{\tilde{\Psi}}}{\partial v^{\Lambda}}}_{=:f_{\tilde{\Lambda}}{}^{\tilde{\Psi}}})), \tag{3.2}$$

that will define the complex structure f on V_{2n}, where $\Delta_{\Psi}{}^{\Lambda}$ is *Norden affinor*. We will define *Neifeld operators* $\bar{m}^{\Psi'}{}_{\tilde{\Psi}} := \overline{m^{\Psi}{}_{\tilde{\Psi}}}$ as

$$0 =: m^{\Lambda}{}_{\tilde{\Lambda}} \bar{m}_{\Psi'}{}^{\tilde{\Lambda}} = \tfrac{1}{2}((\underbrace{\frac{\partial u^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ \frac{\partial x^{\tilde{\Lambda}}}{\partial u^{\Psi'}} - \frac{\partial v^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ \frac{\partial x^{\tilde{\Lambda}}}{\partial v^{\Psi'}}}_{\equiv 0}) + i\ (\tfrac{\partial u^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ \tfrac{\partial x^{\tilde{\Lambda}}}{\partial v^{\Psi'}} + \tfrac{\partial v^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ \tfrac{\partial x^{\tilde{\Lambda}}}{\partial u^{\Psi'}})), \tag{3.3}$$

$$\delta_{\Psi}{}^{\Lambda} =: m^{\Lambda}{}_{\tilde{\Lambda}} m_{\Psi}{}^{\tilde{\Lambda}} = \tfrac{1}{2}((\underbrace{\frac{\partial u^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ \frac{\partial x^{\tilde{\Lambda}}}{\partial u^{\Psi}} + \frac{\partial v^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ \frac{\partial x^{\tilde{\Lambda}}}{\partial v^{\Psi}}}_{=2\delta_{\Psi}{}^{\Lambda}}) - i\ (\tfrac{\partial u^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ \tfrac{\partial x^{\tilde{\Lambda}}}{\partial v^{\Psi}} - \tfrac{\partial v^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ \tfrac{\partial x^{\tilde{\Lambda}}}{\partial u^{\Psi}})), \tag{3.4}$$

Then, $\frac{\partial u^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ \frac{\partial x^{\tilde{\Lambda}}}{\partial v^{\Psi}} \equiv 0$ (where $v^{\Psi'} = \delta_{\Psi}{}^{\Psi'} v^{\Psi}$ and $u^{\Psi'} = \delta_{\Psi}{}^{\Psi'} u^{\Psi}$, and the tensor $\delta_{\Psi}{}^{\Psi'}$ has the identity matrix). Hence,

$$\frac{\partial u^{\Lambda}}{\partial x^{\tilde{\Lambda}}} f_{\tilde{\Psi}}{}^{\tilde{\Lambda}} = \frac{\partial u^{\Lambda}}{\partial x^{\tilde{\Lambda}}}\ (\frac{\partial v^{\Psi}}{\partial x^{\tilde{\Psi}}}\ \frac{\partial x^{\tilde{\Lambda}}}{\partial u^{\Psi}} - \frac{\partial u^{\Psi}}{\partial x^{\tilde{\Psi}}}\ \frac{\partial x^{\tilde{\Lambda}}}{\partial v^{\Psi}}) = \frac{\partial v^{\Lambda}}{\partial x^{\tilde{\Psi}}} \tag{3.5}$$

are *Cauchy-Riemann conditions* if the complex structure has the canonical form in the given basis

$$f_{\tilde{\Lambda}}{}^{\tilde{\Psi}} = \begin{pmatrix} 0 & E \\ -E & 0 \end{pmatrix}, \tag{3.6}$$

where E is the identity matrix of dimension $n \times n$. It is easy to check up that $f^2 = -E$. If now to demand that maps of the atlas on $\mathbb{C}V_n$ have been coordinated by means of holomorphic transformations then by such a way, one can identify the two manifolds: $\mathbb{C}V_n$ and V_{2n}. Thus, it is possible to construct the isomorphism between fibers of the tangent bundles: $\tau^{\mathbb{C}}(\mathbb{C}V_n)$ and $\tau^{\mathbb{C}}(V_{2n})$ as well as between fibers of their real representations: $\tau^{\mathbb{R}}(\mathbb{C}V_n)$ and $\tau^{\mathbb{R}}(V_{2n})$.

Let in the same basis in which the complex structure has the canonical form, our metric be

$$G_{\tilde{\Lambda}\tilde{\Psi}} = \begin{pmatrix} E & 0 \\ 0 & -E \end{pmatrix}, \tag{3.7}$$

where E is the identity matrix of the dimension $n \times n$. The special operators $m_{\Lambda}{}^{\tilde{\Lambda}}$, $m^{\Lambda}{}_{\tilde{\Lambda}}$ which by definition, should satisfy the following system of the equations

$$\begin{cases} m_{\Lambda}{}^{\tilde{\Lambda}} m^{\Psi}{}_{\tilde{\Lambda}} = \delta_{\Lambda}{}^{\Psi}, \\ \\ m_{\Lambda}{}^{\tilde{\Lambda}} \bar{m}^{\Psi'}{}_{\tilde{\Lambda}} = 0, \\ \\ m_{\Lambda}{}^{\tilde{\Lambda}} m^{\Lambda}{}_{\tilde{\Psi}} = \tfrac{1}{2}(\delta_{\tilde{\Psi}}{}^{\tilde{\Lambda}} + i f_{\tilde{\Psi}}{}^{\tilde{\Lambda}}) := \triangle_{\tilde{\Psi}}{}^{\tilde{\Lambda}}, \end{cases} \tag{3.8}$$

answer for the conformity $V_{2n} \longleftrightarrow \mathbb{C}V_n$. Then from the average equation, the parities

$$m^{\Psi}{}_{\tilde{\Omega}} = m^{\Lambda}{}_{\tilde{\Omega}}(m_{\Lambda}{}^{\tilde{\Lambda}} m^{\Psi}{}_{\tilde{\Lambda}}) = (m^{\Lambda}{}_{\tilde{\Omega}} m_{\Lambda}{}^{\tilde{\Lambda}}) m^{\Psi}{}_{\tilde{\Lambda}} = \frac{1}{2} m^{\Psi}{}_{\tilde{\Omega}} + \frac{i}{2} f_{\tilde{\Omega}}{}^{\tilde{\Lambda}} m^{\Psi}{}_{\tilde{\Lambda}}. \tag{3.9}$$

will follow. Whence, the relations

$$m^{\Psi}{}_{\tilde{\Omega}} = i\ f_{\tilde{\Omega}}{}^{\tilde{\Lambda}} m^{\Psi}{}_{\tilde{\Lambda}} \tag{3.10}$$

are the Cauchy-Riemann conditions again. Therefore, in the presence of the metric $G_{\tilde{\Lambda}\tilde{\Psi}}$ for $\mathbb{R}^{2n}_{(n,n)}$, the metric

$$\begin{gathered} g_{\Lambda\Psi} := G_{\tilde{\Lambda}\tilde{\Psi}} m_{\Lambda}{}^{\tilde{\Lambda}} m_{\Psi}{}^{\tilde{\Psi}}, \quad \bar{g}_{\Lambda'\Psi'} := G_{\tilde{\Lambda}\tilde{\Psi}} \bar{m}_{\Lambda'}{}^{\tilde{\Lambda}} \bar{m}_{\Psi'}{}^{\tilde{\Psi}}, \\ g_{\Lambda'\Psi} := G_{\tilde{\Lambda}\tilde{\Psi}} \bar{m}_{\Lambda'}{}^{\tilde{\Lambda}} m_{\Psi}{}^{\tilde{\Psi}}, \quad \bar{g}_{\Lambda\Psi'} := G_{\tilde{\Lambda}\tilde{\Psi}} \bar{m}_{\Lambda}{}^{\tilde{\Lambda}} \bar{m}_{\Psi'}{}^{\tilde{\Psi}} \end{gathered} \tag{3.11}$$

is induced in the complex representation. In the standard complex representation (where instead of the space $\mathbb{R}^{2n}_{(n,n)}$, the space $\mathbb{R}^{2n}$ undertakes), this means that $g_{\Lambda'\Psi'} = \bar{g}_{\Lambda'\Psi'} = 0$ and leads to the Hermitian metric $g_{\Lambda'\Psi}$. However, in our case, it is necessary to take advantage of the conditions

$$g_{\Lambda'\Psi} = \bar{g}_{\Lambda\Psi'} = 0. \tag{3.12}$$

It will lead to some restrictions on the metric tensor

$$\begin{gathered} 0 = g_{\Lambda'\Psi} \bar{m}^{\Lambda'}{}_{\tilde{\Omega}} m^{\Psi}{}_{\tilde{\Theta}} = (G_{\tilde{\Lambda}\tilde{\Psi}} \bar{m}_{\Lambda'}{}^{\tilde{\Lambda}} m_{\Psi}{}^{\tilde{\Psi}}) \bar{m}^{\Lambda'}{}_{\tilde{\Omega}} m^{\Psi}{}_{\tilde{\Theta}} = G_{\tilde{\Lambda}\tilde{\Psi}} (\bar{m}_{\Lambda'}{}^{\tilde{\Lambda}} \bar{m}^{\Lambda'}{}_{\tilde{\Omega}})(m_{\Psi}{}^{\tilde{\Psi}} m^{\Psi}{}_{\tilde{\Theta}}) = \\ = G_{\tilde{\Lambda}\tilde{\Psi}} \cdot \tfrac{1}{2}(\delta_{\tilde{\Omega}}{}^{\tilde{\Lambda}} - i f_{\tilde{\Omega}}{}^{\tilde{\Lambda}}) \cdot \tfrac{1}{2}(\delta_{\tilde{\Theta}}{}^{\tilde{\Psi}} + i f_{\tilde{\Theta}}{}^{\tilde{\Psi}}) = \\ = \tfrac{1}{4}(\underbrace{G_{\tilde{\Omega}\tilde{\Theta}} + (f_{\tilde{\Omega}}{}^{\tilde{\Lambda}} f_{\tilde{\Theta}}{}^{\tilde{\Psi}} G_{\tilde{\Lambda}\tilde{\Psi}})}_{=0} + i(\underbrace{(f_{\tilde{\Omega}}{}^{\tilde{\Lambda}} G_{\tilde{\Lambda}\tilde{\Theta}}) - (f_{\tilde{\Theta}}{}^{\tilde{\Psi}} G_{\tilde{\Omega}\tilde{\Psi}})}_{=0})) = 0. \end{gathered} \tag{3.13}$$

Therefore, our pseudo-Euclidian metric with the canonical complex structure satisfies the equation

$$f_{\tilde{\Omega}}{}^{\tilde{\Lambda}} G_{\tilde{\Lambda}\tilde{\Psi}} f_{\tilde{\Theta}}{}^{\tilde{\Psi}} = -G_{\tilde{\Omega}\tilde{\Theta}}, \tag{3.14}$$

$$\begin{pmatrix} 0 & E \\ -E & 0 \end{pmatrix} \begin{pmatrix} E & 0 \\ 0 & -E \end{pmatrix} \begin{pmatrix} 0 & -E \\ E & 0 \end{pmatrix} = - \begin{pmatrix} E & 0 \\ 0 & -E \end{pmatrix}.$$

Besides, it will be executed

$$G_{\tilde{\Lambda}\tilde{\Psi}} = m^{\Lambda}{}_{\tilde{\Lambda}} m^{\Psi}{}_{\tilde{\Psi}} g_{\Lambda\Psi} + \bar{m}^{\Lambda'}{}_{\tilde{\Lambda}} \bar{m}^{\Psi'}{}_{\tilde{\Psi}} \bar{g}_{\Lambda'\Psi'}. \tag{3.15}$$

Example 3.1. *At n=1 for* $\mathbb{R}^2_{(1,1)}$

$$\begin{gathered} m_\Lambda{}^{\tilde{\Lambda}} = \tfrac{1}{\sqrt{2}}(1, i), \quad (m^T)_{\tilde{\Lambda}}{}^{\Lambda} = \tfrac{1}{\sqrt{2}} \left(\begin{array}{c} 1 \\ -i \end{array} \right), \quad g_{\Lambda\Psi} = 1, \\ G_{\tilde{\Lambda}\tilde{\Psi}} = \left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array} \right), \quad f_{\tilde{\Lambda}}{}^{\tilde{\Psi}} = \left(\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array} \right). \end{gathered} \tag{3.16}$$

Now the operators (3.8) can be applied to (2.25). For this purpose, we will consider the decomposition for n=2 (N=1)

$$f_{\tilde{\Psi}}{}^{\tilde{\Lambda}}\gamma_{\tilde{\Lambda}} = \tilde{F}\gamma_{\tilde{\Psi}}\tilde{\tilde{F}}, \quad \tilde{F}^4 = \tilde{\tilde{F}}^4 = E. \tag{3.17}$$

The proof of this formula will be given in the following section (6.15). From (3.17), it will follow

$$\begin{gathered} \triangle_{\tilde{\Psi}}{}^{\tilde{\Lambda}}\gamma_{\tilde{\Lambda}} = \tfrac{1}{2}(\delta_{\tilde{\Psi}}{}^{\tilde{\Lambda}} + i f_{\tilde{\Psi}}{}^{\tilde{\Lambda}})\gamma_{\tilde{\Lambda}} = \\ = \underbrace{\frac{1}{2}(\delta + \frac{(1+i)}{2}\tilde{F} + \frac{(1-i)}{2}\tilde{F}^3)}_{:=\tilde{\triangle}_+} \gamma_{\tilde{\Psi}} \underbrace{\frac{1}{2}(\delta + \frac{(1+i)}{2}\tilde{\tilde{F}} + \frac{(1-i)}{2}\tilde{\tilde{F}}^3)}_{:=\tilde{\tilde{\triangle}}_+} + \\ + \underbrace{\frac{1}{2}(\delta - \frac{(1+i)}{2}\tilde{F} - \frac{(1-i)}{2}\tilde{F}^3)}_{:=\tilde{\triangle}_-} \gamma_{\tilde{\Psi}} \underbrace{\frac{1}{2}(\delta - \frac{(1+i)}{2}\tilde{\tilde{F}} - \frac{(1-i)}{2}\tilde{\tilde{F}}^3)}_{:=\tilde{\tilde{\triangle}}_-} = \\ = \tilde{\triangle}_+\gamma_{\tilde{\Psi}}\tilde{\tilde{\triangle}}_+ + \tilde{\triangle}_-\gamma_{\tilde{\Psi}}\tilde{\tilde{\triangle}}_-. \end{gathered} \tag{3.18}$$

This means that one can set for n=2 (N=1)

$$\left\{ \begin{array}{l} \tilde{m}_{\pm\hat{A}}{}^{\tilde{B}}\tilde{m}_{\pm}{}^{\hat{C}}{}_{\tilde{B}} = \delta_{\hat{A}}{}^{\hat{C}}, \\ \\ \tilde{m}_{\pm\hat{A}}{}^{\tilde{B}}\tilde{m}_{\mp}{}^{\hat{C}}{}_{\tilde{B}} = 0, \\ \\ \tilde{m}_{\pm\hat{A}}{}^{\tilde{B}}\tilde{m}_{\pm}{}^{\hat{A}}{}_{\tilde{C}} = \\ = \underbrace{\tfrac{1}{2}(\delta_{\tilde{C}}{}^{\tilde{B}} \pm \tfrac{(1+i)}{2}\tilde{F}_{\tilde{C}}{}^{\tilde{B}} \pm \tfrac{(1-i)}{2}(\tilde{F}^3)_{\tilde{C}}{}^{\tilde{B}})}_{:=\tilde{\triangle}_{\pm\tilde{C}}{}^{\tilde{B}}}, \end{array} \right. \quad \left\{ \begin{array}{l} \tilde{\tilde{m}}_{\pm\hat{A}}{}^{\tilde{B}}\tilde{\tilde{m}}_{\pm}{}^{\hat{C}}{}_{\tilde{B}} = \delta_{\hat{A}}{}^{\hat{C}}, \\ \\ \tilde{\tilde{m}}_{\pm\hat{A}}{}^{\tilde{B}}\tilde{\tilde{m}}_{\mp}{}^{\hat{C}}{}_{\tilde{B}} = 0, \\ \\ \tilde{\tilde{m}}_{\pm\hat{A}}{}^{\tilde{B}}\tilde{\tilde{m}}_{\pm}{}^{\hat{A}}{}_{\tilde{C}} = \\ = \underbrace{\tfrac{1}{2}(\delta_{\tilde{C}}{}^{\tilde{B}} \pm \tfrac{(1+i)}{2}\tilde{\tilde{F}}_{\tilde{C}}{}^{\tilde{B}} \pm \tfrac{(1-i)}{2}(\tilde{\tilde{F}}^3)_{\tilde{C}}{}^{\tilde{B}})}_{:=\tilde{\tilde{\triangle}}_{\pm\tilde{C}}{}^{\tilde{B}}}. \end{array} \right. \tag{3.19}$$

Here $\tilde{A}, \tilde{B}, \tilde{C}, \tilde{D}, ... = \overline{1, (2N)^2}$, $\hat{A}, \hat{B}, \hat{C}, \hat{D}, ... = \overline{1, 2N}$. Note that we can introduce the operation of *conjugation* (not complex!) such that the operators $\tilde{m}_\pm$ ($\tilde{\tilde{m}}_\pm$) and $\tilde{m}_\mp$ ($\tilde{\tilde{m}}_\mp$) are conjugate to each other for such the operation. Hence, as the operators γ_Λ, one can take either $\gamma_{+\Lambda}$ or $\gamma_{-\Lambda}$

$$\gamma_{\pm\Lambda\hat{A}}{}^{\hat{B}} := m_\Lambda{}^{\tilde{\Lambda}}\gamma_{\tilde{\Lambda}\tilde{C}}{}^{\tilde{D}}\tilde{m}_{\pm\hat{A}}{}^{\tilde{C}}\tilde{\tilde{m}}_{\pm}{}^{\hat{B}}{}_{\tilde{D}}. \tag{3.20}$$

Example 3.2. *At n=2 for* $\mathbb{R}^4_{(2,2)}$

$$m_{\Lambda}{}^{\tilde{\Lambda}} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & i & 0 & 0 \\ 0 & 0 & 1 & i \end{pmatrix}, \quad (m^T)_{\tilde{\Lambda}}{}^{\Lambda} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ -i & 0 \\ 0 & 1 \\ 0 & -i \end{pmatrix}, \quad g_{\Lambda\Psi} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$
$$G_{\tilde{\Lambda}\tilde{\Psi}} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad f_{\tilde{\Lambda}}{}^{\tilde{\Psi}} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}. \tag{3.21}$$

$\gamma_{\tilde{\Lambda}}$ *are defined according to (2.26),* γ_{Λ} *are defined according to (2.28).*

$$\begin{gathered} m_{\hat{A}}{}^{\tilde{B}} := \tilde{m}_{+\hat{A}}{}^{\tilde{B}} = \tilde{\tilde{m}}_{+\hat{A}}{}^{\tilde{B}} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & i & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}, \\ (m^T)_{\tilde{B}}{}^{\hat{A}} = (\tilde{m}^T_{+})_{\tilde{B}}{}^{\hat{A}} = (\tilde{\tilde{m}}^T_{+})_{\tilde{B}}{}^{\hat{A}} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ -i & 0 \\ 0 & 1 \\ 0 & 1 \end{pmatrix}, \\ \tilde{m}_{-\hat{A}}{}^{\tilde{B}} = \tilde{\tilde{m}}_{-\hat{A}}{}^{\tilde{B}} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & -i & 0 & 0 \\ 0 & 0 & 1 & -1 \end{pmatrix}, \\ (\tilde{m}^T_{-})_{\tilde{B}}{}^{\hat{A}} = (\tilde{\tilde{m}}^T_{-})_{\tilde{B}}{}^{\hat{A}} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ i & 0 \\ 0 & 1 \\ 0 & -1 \end{pmatrix}, \\ F^{\tilde{A}}{}_{\tilde{B}} := \tilde{F}^{\tilde{A}}{}_{\tilde{B}} = \tilde{\tilde{F}}^{\tilde{A}}{}_{\tilde{B}} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}. \end{gathered} \tag{3.22}$$

By the way, it once again says that the operator F *is a matrix fourth root from the matrix identity. The square* F^2 *corresponds to the trivial transformation multiplied by -1 in the space* $\mathbb{R}^4_{(2,2)}$.

In the case of arbitrary n, the algorithm for the reduction of the spinor dimension is as follows.

Algorithm 3.1. *The complex structure can be represented as a product of elementary transformations of the dimension* 4×4. *Each I-th elementary transformation generates the operator* $(m_I)_{\Lambda}{}^{\tilde{\Lambda}}$ *which is responsible for the transition from the real representation of the subspace* $(R_I)^4_{(2,2)} \subset \mathbb{R}^{2n}_{(2n,2n)}$ *to the complex representation* $(C_I)^2 \subset R^{2n-4}_{(2n-2,2n-2)} \oplus (C_I)^2$, *and hence this lowers the dimension of the space on 2. Accordingly, in the spinor space, the operators* $\tilde{\tilde{m}}_{I_{\pm}}$, $\tilde{m}_{I_{\pm}}$ *which can reduce the dimension of the spinor space by 2 times are generated. Obviously, the n/2 steps are required* ($I = \overline{1, \frac{n}{2}}$). *This would reduce the spinor dimension by* $2^{\frac{n}{2}} = 2N$ *times and would lead to the complex matrix representation of the dimension* $2N \times 2N$. *The*

operators $\tilde{\tilde{m}}_{I_\pm}$*,* $\tilde{m}_{I_\pm}$ *will satisfy the following relations*

$$\begin{cases} \tilde{m}_{I_\pm \hat{A}}{}^{\tilde{B}} \tilde{m}_{I_\pm}{}^{\hat{C}}{}_{\tilde{B}} = \delta_{\hat{A}}{}^{\hat{C}}, \\ \\ \tilde{m}_{I_\pm \hat{A}}{}^{\tilde{B}} \tilde{m}_{I_\mp}{}^{\hat{C}}{}_{\tilde{B}} = 0, \\ \\ \tilde{m}_{I_\pm \hat{A}}{}^{\tilde{B}} \tilde{m}_{I_\pm}{}^{\hat{A}}{}_{\tilde{C}} = \tilde{\triangle}_{I_\pm \tilde{C}}{}^{\tilde{B}}, \end{cases} \qquad \begin{cases} \tilde{\tilde{m}}_{I_\pm \hat{A}}{}^{\tilde{B}} \tilde{\tilde{m}}_{I_\pm}{}^{\hat{C}}{}_{\tilde{B}} = \delta_{\hat{A}}{}^{\hat{C}}, \\ \\ \tilde{\tilde{m}}_{I_\pm \hat{A}}{}^{\tilde{B}} \tilde{\tilde{m}}_{I_\mp}{}^{\hat{C}}{}_{\tilde{B}} = 0, \\ \\ \tilde{\tilde{m}}_{I_\pm \hat{A}}{}^{\tilde{B}} \tilde{\tilde{m}}_{I_\pm}{}^{\hat{A}}{}_{\tilde{C}} = \tilde{\tilde{\triangle}}_{I_\pm \tilde{C}}{}^{\tilde{B}}. \end{cases} \tag{3.23}$$

Here for I step $\tilde{A}, \quad \tilde{B}, \quad \tilde{C}, \quad \tilde{D}, ... = \overline{1, \frac{(2N)^2}{2^{I-1}}}$, $\hat{A}, \hat{B}, \hat{C}, \hat{D}, ... = \overline{1, \frac{(2N)^2}{2^I}}$. *Then we can construct the operators* ($\tilde{m}_{J_{z_J}} := \tilde{m}_{J_\pm \hat{A}}{}^{\tilde{B}}$, $\tilde{\tilde{m}}^*_{J_{z_J}} := \tilde{\tilde{m}}_{J_\pm}{}^{\hat{A}}{}_{\tilde{B}}$)

$$\tilde{M}_K := \tilde{m}_{\frac{n}{2}_{z_{\frac{n}{2}}}} \tilde{m}_{\frac{n}{2}-1_{z_{\frac{n}{2}-1}}} \cdot ... \cdot \tilde{m}_{2_{z_2}} \tilde{m}_{1_{z_1}}, \quad \tilde{\tilde{M}}_K := \tilde{\tilde{m}}^*_{\frac{n}{2}_{z_{\frac{n}{2}}}} \tilde{\tilde{m}}^*_{\frac{n}{2}-1_{z_{\frac{n}{2}-1}}} \cdot ... \cdot \tilde{\tilde{m}}^*_{2_{z_2}} \tilde{\tilde{m}}^*_{1_{z_1}}, \tag{3.24}$$

where z_J $(J = \overline{1, \frac{n}{2}})$ *is equal to 0 for the sign* $\ll - \gg$ *or 1 for the sign* $\ll + \gg$ *then* $K = \sum\limits_{J=1}^{\frac{n}{2}} z_J \cdot 2^{J-1} + 1$. *If it does not matter which of the K-th operators will be selected then the number K will be omitted. Using these operators, one can define*

$$(\gamma_K)_\Lambda := \tilde{M}_K (m_\Lambda{}^{\tilde{\Lambda}} \gamma_{\tilde{\Lambda}}) (\tilde{\tilde{M}}_K)^T. \tag{3.25}$$

The corresponding example will be analyzed below (Algorithm 8.1).

4 Real inclusion. Involution

In this section it is told about how locally to enclose a real n-dimensional (pseudo-) Riemannian space into the complex representation $\mathbb{C}V_n$ of the 2n-dimensional pseudo-Riemannian space. For this purpose, a real surface in $\mathbb{C}V_n$ with a real parametrization is considered. This enclosure is induced in the tangent bundle by means of an inclusion operator $H_i{}^\Lambda$ [15] with which the help one can obtain the fiber of a real tangent bundle equipped with the (pseudo-)Euclidian metric. The index of such the metric will be significantly depend on the kind of the inclusion operator. The conclusion of all results of this section is made on the basis [15].

We will consider the real (pseudo-) Riemannian space V_n as a surface of the real dimension n in the space $\mathbb{C}V_n$, i.e., a surface defined by means of the parametrical equation

$$w^\Lambda = w^\Lambda(u^i), \quad (\Lambda, \Psi, ..., i, j, g, h = \overline{1, n}), \tag{4.1}$$

where w^Λ are the complex coordinates of a point x of the base, and u^i are parameters: the local coordinates of a point of the space V_n. The partial derivatives ($\partial_i w^\Lambda =: H_i{}^\Lambda$) locally define *inclusion* of the real tangent spaces (4.1) in the complex tangent space $\tau_x^{\mathbb{C}}$ as follows

$$H : \tau_x^{\mathbb{R}} \longmapsto \tau_x^{\mathbb{C}}, \tag{4.2}$$

$$\begin{gathered} w^\Lambda = w^\Lambda(u^i(t)), \quad r^\Lambda := \tfrac{dw^\Lambda}{dt} = H_i{}^\Lambda \tfrac{du^i}{dt} =: H_i{}^\Lambda r^i, \\ \tfrac{du^i}{dt} \in \tau_x^{\mathbb{R}} \longmapsto \tfrac{dw^\Lambda}{dt} \in \tau_x^{\mathbb{C}}, \end{gathered} \tag{4.3}$$

where the differentiation is conducted along a real curve $\gamma(t)$ of the surface (4.1). Since $\| H_i{}^\Lambda \|$ is a nonsingular Jacobian matrix then there is the operator $H^i{}_\Lambda$ such that

$$\left\{ \begin{array}{c} H^i{}_\Lambda H_i{}^\Psi = \delta_\Lambda{}^\Psi, \\ H^i{}_\Lambda H_j{}^\Lambda = \delta_j{}^i. \end{array} \right. \tag{4.4}$$

From here, it follows that the operator $H_i{}^\Lambda$ defines *involution*

$$S_\Lambda{}^{\Psi'} = H^i{}_\Lambda \bar{H}_i{}^{\Psi'} \tag{4.5}$$

in the complex space, where the coordinates $\bar{H}_i{}^{\Psi'}$ are in *complex conjugate* to the coordinates $H_i{}^\Psi$. Therefore,

$$r^i = H^i{}_\Lambda r^\Lambda = \overline{H^i{}_\Lambda r^\Lambda} \quad \Rightarrow \quad S_\Lambda{}^{\Psi'} r^\Lambda = \bar{r}^{\Psi'}. \tag{4.6}$$

It is a necessary and sufficient condition for the vector $r^\Lambda \in \tau_x^{\mathbb{C}}$ to be real. Thus,

$$S_\Lambda{}^{\Psi'} \bar{S}_{\Psi'}{}^\Phi = \delta_\Lambda{}^\Phi. \tag{4.7}$$

We will define the metric of V_n (the real (pseudo-) Riemannian space) by the condition

$$g_{\Lambda\Psi} r^\Lambda r^\Psi = \overline{g_{\Lambda\Psi} r^\Lambda r^\Psi}, \quad \forall \bar{r}^{\Psi'} = S_\Lambda{}^{\Psi'} r^\Lambda. \tag{4.8}$$

This means that a real tensor of the space V_n is defined as the tensor self-conjugated under the specified Hermitian involution

$$g_{\Lambda\Psi} = S_\Lambda{}^{\Phi'} S_\Psi{}^{\Theta'} \bar{g}_{\Phi'\Theta'}. \tag{4.9}$$

Therefore, the tensor

$$g_{ij} := H_i{}^\Lambda H_j{}^\Psi g_{\Lambda\Psi} = \overline{H_i{}^\Lambda H_j{}^\Psi g_{\Lambda\Psi}} \tag{4.10}$$

will be the metric tensor of $V_n \subset \mathbb{C}V_n$. The kind of the metric g_{ij} significantly depends on the structure of the operator $H_i{}^\Lambda$ and hence the involution $S_\Lambda{}^{\Psi'}$.

Example 4.1. *Let there be the complex Euclidian space $\mathbb{CR}^4$. An inclusion of a real space in the complex can be obtain by means of one of the three various ways*

$$H_i{}^\Lambda = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad H_i{}^\Lambda = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \end{pmatrix}, \quad H_i{}^\Lambda = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \end{pmatrix}.$$

$$G_{\Lambda\Psi} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

$$g_{ij} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad g_{ij} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad g_{ij} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}. \tag{4.11}$$

5 Elementary transformations of the orthogonal group

In this section, it is told about how to lead a pseudo-orthogonal transformation to a block-diagonal form, and then to pass to the complex representation of such the transformation. The conclusion of all results of this section is made on the basis [10], [20].

Let's consider an orthogonal transformation in the space $\mathbb{R}^{2n}_{(n,n)}$ which is set by the formula

$$G_{\tilde{\Lambda}\tilde{\Psi}} S_{\tilde{\Omega}}{}^{\tilde{\Lambda}} S_{\tilde{\Gamma}}{}^{\tilde{\Psi}} = G_{\tilde{\Omega}\tilde{\Gamma}}. \tag{5.1}$$

Thus, a basis is chosen so that $\| G_{\tilde{\Lambda}\tilde{\Psi}} \|$ will have the diagonal form.

Theorem 5.1. *Conformal transformations of the space $\mathbb{R}^{2n-2}_{(n-1,n-1)}$ form the group $O(n,n)$ consisting of*
1. ***rotations*** *from $O(n-1,n-1)$,* *3.* ***translations,*** *5.* ***superpositions 1-4.***
2. ***dilations,*** *4.* ***inversions,***
Then any transformation of $O(n,n)$ can be represented as a product of ***elementary transformations***

$$S_{\tilde{\Lambda}_1}{}^{\tilde{\Lambda}_{J+1}} = \pm \prod_{I=1}^{J} (\pm (r_I)_{\tilde{\Lambda}_I} (r_I)^{\tilde{\Lambda}_{I+1}} - \delta_{\tilde{\Lambda}_I}{}^{\tilde{\Lambda}_{I+1}}), \quad (r_I)_{\tilde{\Lambda}_I} (r_I)^{\tilde{\Lambda}_I} = \pm 2. \tag{5.2}$$

Proof. The elementary transformations (5.2) are really orthogonal

$$(\pm r_{\tilde{\Lambda}} r^{\tilde{\Psi}} - \delta_{\tilde{\Lambda}}{}^{\tilde{\Psi}})(\pm r_{\tilde{\Psi}} r^{\tilde{\Theta}} - \delta_{\tilde{\Psi}}{}^{\tilde{\Theta}}) = (\pm 2) r_{\tilde{\Lambda}} r^{\tilde{\Theta}} \mp r_{\tilde{\Lambda}} r^{\tilde{\Theta}} \mp r_{\tilde{\Lambda}} r^{\tilde{\Theta}} + \delta_{\tilde{\Lambda}}{}^{\tilde{\Theta}} = \delta_{\tilde{\Lambda}}{}^{\tilde{\Theta}}. \tag{5.3}$$

Usual rotations from the group $O(2)$ can be represented as (5.2). At the same time, $r^{\tilde{\Lambda}} = \sqrt{2}(\cos\frac{\alpha}{2},\ \ \sin\frac{\alpha}{2})$, and there are the two disconnected classes of rotations

$$a).\begin{pmatrix} \cos\alpha & \sin\alpha \\ \sin\alpha & -\cos\alpha \end{pmatrix}, \quad b).\begin{pmatrix} \cos\alpha & \sin\alpha \\ -\sin\alpha & \cos\alpha \end{pmatrix}. \tag{5.4}$$

However, the transformation from b). is obtained by a superposition of the two transformations from a).

1. A transformation from the group $O(1,1)$ is *boost* of one of the 4 forms

$$\begin{array}{llll} a). & \begin{pmatrix} \operatorname{ch}\theta & \operatorname{sh}\theta \\ \operatorname{sh}\theta & \operatorname{ch}\theta \end{pmatrix}, & b). & \begin{pmatrix} \operatorname{ch}\theta & -\operatorname{sh}\theta \\ \operatorname{sh}\theta & -\operatorname{ch}\theta \end{pmatrix}, \\ c). & \begin{pmatrix} -\operatorname{ch}\theta & \operatorname{sh}\theta \\ -\operatorname{sh}\theta & \operatorname{ch}\theta \end{pmatrix}, & d). & \begin{pmatrix} -\operatorname{ch}\theta & -\operatorname{sh}\theta \\ -\operatorname{sh}\theta & -\operatorname{ch}\theta \end{pmatrix}. \end{array} \tag{5.5}$$

   The transformation from b). is represented as in the equation (5.2), where $r^{\tilde{\Lambda}} = \sqrt{2}(\operatorname{ch}\frac{\theta}{2},\ \ \operatorname{sh}\frac{\theta}{2})$. The transformation from a). is a superposition of the two transformations from b). The transformations from c). and d). differ only in the sign from the transformations from b). and a). respectively. This describes the four disconnected components of $O(1,1)$.

2. Let *one-dimensional dilation* in $\mathbb{R}^1$ has the form $\tilde{x} = \lambda x$. Consider the light cone in $\mathbb{R}^3_{(1,2)}$ defined as $T^2 - Z^2 - X^2 = 0$. We cut it by the plane $T + Z = 1$ and perform the stereographic projection of the cross section onto the line $T = 1$, $Z = 0$ that induces the one-dimensional space $\mathbb{R}^1$ with the single coordinate $x = \frac{X}{T-Z}$. Then $\tilde{T} + \tilde{Z} = \lambda(T+Z)$, $\tilde{T} - \tilde{Z} = \lambda^{-1}(T-Z)$, $\tilde{X} = X$. If $\lambda > 0$ then this transformation will be the boost from a).; if $\lambda < 0$, this transformation will the boost from d).

3. Consider *translation* $\tilde{x} = x + a$ in the space $\mathbb{R}^1$. This will lead to the relations: $\tilde{T} - \tilde{Z} = T - Z$, $\tilde{T} + \tilde{Z} = T + Z + 2aX + a^2(T-Z)$, $\tilde{X} = X + a(T-Z)$. The fixed vector of this transformation will be an isotropic vector of the form $(b, 0, b)$. It is impossible to translate an isotropic vector to a non-isotropic vector by any pseudo-orthogonal transformation since such the transformations keep the constant value $T^2 - X^2 - Z^2$. This means that we can not apply such the transformations to the basis to direct an isotropic vector along a non-isotropic axis. However, this does not prevent for the dilation to be decomposed into elementary transformations: *rotation*, *boost*, and one more such *rotation*. To do this, as the rotation we take the elementary transformation

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\alpha & -\sin\alpha \\ 0 & \sin\alpha & \cos\alpha \end{pmatrix} \tag{5.6}$$

with $\operatorname{tg}\alpha = \frac{a}{2}$, and as the boost from a)., we take the elementary transformation

$$\begin{pmatrix} \operatorname{ch}\theta & \operatorname{sh}\theta & 0 \\ \operatorname{sh}\theta & \operatorname{ch}\theta & 0 \\ 0 & 0 & 1 \end{pmatrix} \tag{5.7}$$

with $\operatorname{ch}\theta = \frac{a^2}{2} + 1$. Then, this composition will have the form

$$\begin{gathered} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{1+\frac{a^2}{4}}} & -\frac{\frac{a}{2}}{\sqrt{1+\frac{a^2}{4}}} \\ 0 & \frac{\frac{a}{2}}{\sqrt{1+\frac{a^2}{4}}} & \frac{1}{\sqrt{1+\frac{a^2}{4}}} \end{pmatrix} \begin{pmatrix} 1+\frac{a^2}{2} & a\sqrt{1+\frac{a^2}{4}} & 0 \\ a\sqrt{1+\frac{a^2}{4}} & 1+\frac{a^2}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot \\ \cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{1+\frac{a^2}{4}}} & -\frac{\frac{a}{2}}{\sqrt{1+\frac{a^2}{4}}} \\ 0 & \frac{\frac{a}{2}}{\sqrt{1+\frac{a^2}{4}}} & \frac{1}{\sqrt{1+\frac{a^2}{4}}} \end{pmatrix} = \begin{pmatrix} 1+\frac{a^2}{2} & a & -\frac{a^2}{2} \\ a & 1 & -a \\ \frac{a^2}{2} & a & 1-\frac{a^2}{2} \end{pmatrix}. \end{gathered} \tag{5.8}$$

4. *Inversion* $\tilde{x} = \frac{1}{x}$ in $\mathbb{R}^1$ induces the rotation $\tilde{T} = T$, $\tilde{Z} = -Z$, $\tilde{X} = X$.

5. *Superpositions* of all one-dimensional transformations of the form 1-4 and *rotations* from $O(n-1, n-1)$ represent the conformal transformation group of the space $\mathbb{R}^{2n-2}_{(n-1,n-1)}$ which will be isomorphic to the group $O(n, n)$.

□

Example 5.1. *Let's consider the group of pseudo-orthogonal transformations* $O(1,2)$ *of the space* $\mathbb{R}^3_{(1,2)}$. *As is known, eigenvalues of any such a transformation are roots of a polynomial of third degree. One of such the values should be obligatory a real number. Since pseudo-orthogonal transformations satisfy the parity (5.1) then the square of such the real number is equal to 1. Therefore, any transformation from the group* $O(1,2)$ *possesses a fixed axis. However, not always this axis can be combined with a coordinate axis using pseudo-orthogonal transformations of the basis only.*

1. *Let's consider the composition of a rotation and a boost for the space* $\mathbb{R}^3_{(1,2)}$

$$\begin{pmatrix} \operatorname{ch}\theta & -\operatorname{sh}\theta & 0 \\ \operatorname{sh}\theta & -\operatorname{ch}\theta & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\alpha & \sin\alpha \\ 0 & \sin\alpha & -\cos\alpha \end{pmatrix} = \begin{pmatrix} \operatorname{ch}\theta & -\cos\alpha\operatorname{sh}\theta & -\sin\alpha\operatorname{sh}\theta \\ \operatorname{sh}\theta & -\cos\alpha\operatorname{ch}\theta & -\sin\alpha\operatorname{ch}\theta \\ 0 & \sin\alpha & -\cos\alpha \end{pmatrix}. \tag{5.9}$$

Let this composition will leave fixed the vector (a,b,c)

$$\left\{ \begin{array}{rcll} a\operatorname{ch}\theta - b\cos\alpha\operatorname{sh}\theta - c\sin\alpha\operatorname{sh}\theta & = & a, & \frac{c}{b} = \operatorname{tg}\frac{\alpha}{2}, \quad \frac{a}{b} = \operatorname{cth}\frac{\theta}{2}, \\ a\operatorname{sh}\theta - b\cos\alpha\operatorname{ch}\theta - c\sin\alpha\operatorname{ch}\theta & = & b, & \\ b\sin\alpha - c\cos\alpha & = & c, & r = (b\operatorname{cth}\frac{\theta}{2}, b, b\operatorname{tg}\frac{\alpha}{2}). \end{array} \right. \tag{5.10}$$

Let's spend the basis transformation

$$\begin{pmatrix} \operatorname{ch}\frac{\theta}{2} & -\operatorname{sh}\frac{\theta}{2} & 0 \\ -\operatorname{sh}\frac{\theta}{2} & \operatorname{ch}\frac{\theta}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} b\operatorname{cth}\frac{\theta}{2} \\ b \\ b\operatorname{tg}\frac{\alpha}{2} \end{pmatrix} = \begin{pmatrix} b/\operatorname{sh}\frac{\theta}{2} \\ 0 \\ b\operatorname{tg}\frac{\alpha}{2} \end{pmatrix}. \tag{5.11}$$

As now $|r|^2 = b^2(1/\operatorname{sh}^2\frac{\theta}{2} - \operatorname{tg}^2\frac{\alpha}{2})$, *we will set* $b := \pm\frac{1}{\sqrt{|1/\operatorname{sh}^2\frac{\theta}{2} - \operatorname{tg}^2\frac{\alpha}{2}|}}$. *Then there is the basis transformation*

$$\begin{pmatrix} \operatorname{ch}\phi & 0 & \operatorname{sh}\phi \\ 0 & 1 & 0 \\ \operatorname{sh}\phi & 0 & \operatorname{ch}\phi \end{pmatrix} \begin{pmatrix} b/\operatorname{sh}\frac{\theta}{2} \\ 0 \\ b\operatorname{tg}\frac{\alpha}{2} \end{pmatrix}, \quad \begin{array}{l} a).\ \text{or}\ \operatorname{ch}\phi = \frac{b}{\operatorname{sh}\frac{\theta}{2}}, \quad \operatorname{sh}\phi = -b\operatorname{tg}\frac{\alpha}{2}, \\ b).\ \text{or}\ \operatorname{sh}\phi = -\frac{b}{\operatorname{sh}\frac{\theta}{2}}, \quad \operatorname{ch}\phi = \ b\operatorname{tg}\frac{\alpha}{2}. \end{array} \tag{5.12}$$

In the case a)., the coordinate axis (1,0,0) is fixed, and in the case b)., the axis (0,0,1) is fixed. If $1/\operatorname{sh}\frac{\theta}{2} = \pm\operatorname{tg}\frac{\alpha}{2}$ *then it is impossible to simplify the initial superposition with the help of any pseudo-orthogonal rotation of the basis only.*

2. *Let's consider the composition of two boosts for the space* $\mathbb{R}^3_{(1,2)}$

$$\begin{pmatrix} \operatorname{ch}\theta & -\operatorname{sh}\theta & 0 \\ \operatorname{sh}\theta & -\operatorname{ch}\theta & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \operatorname{ch}\psi & 0 & -\operatorname{sh}\psi \\ 0 & 1 & 0 \\ \operatorname{sh}\psi & 0 & -\operatorname{ch}\psi \end{pmatrix} = \begin{pmatrix} \operatorname{ch}\theta\operatorname{ch}\psi & -\operatorname{sh}\theta & -\operatorname{sh}\psi\operatorname{ch}\theta \\ \operatorname{sh}\theta\operatorname{ch}\psi & -\operatorname{ch}\theta & -\operatorname{sh}\psi\operatorname{sh}\theta \\ \operatorname{sh}\psi & 0 & -\operatorname{ch}\psi \end{pmatrix}. \tag{5.13}$$

Let this composition will leave motionless the vector (a,b,c)

$$\left\{ \begin{array}{rcl} a\operatorname{ch}\theta\operatorname{ch}\psi - b\operatorname{sh}\theta - c\operatorname{sh}\psi\operatorname{ch}\theta & = & a \\ a\operatorname{sh}\theta\operatorname{ch}\psi - b\operatorname{ch}\theta - c\operatorname{sh}\psi\operatorname{sh}\theta & = & b \\ a\operatorname{sh}\psi - c\operatorname{ch}\psi & = & c \end{array} \right., \quad \begin{array}{l} \frac{c}{a} = \operatorname{th}\frac{\psi}{2}, \quad \frac{b}{a} = \operatorname{th}\frac{\theta}{2}, \\ \\ r = (a, a\operatorname{th}\frac{\theta}{2}, a\operatorname{th}\frac{\psi}{2}). \end{array} \tag{5.14}$$

Let's spend the basis transformation

$$\begin{pmatrix} \operatorname{ch}\frac{\theta}{2} & -\operatorname{sh}\frac{\theta}{2} & 0 \\ -\operatorname{sh}\frac{\theta}{2} & \operatorname{ch}\frac{\theta}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a \\ a\operatorname{th}\frac{\theta}{2} \\ a\operatorname{th}\frac{\psi}{2} \end{pmatrix} = \begin{pmatrix} a/\operatorname{ch}\frac{\theta}{2} \\ 0 \\ a\operatorname{th}\frac{\psi}{2} \end{pmatrix}. \tag{5.15}$$

As now $|r|^2 = a^2(1/\operatorname{ch}^2\frac{\theta}{2} - \operatorname{th}^2\frac{\psi}{2})$*, we will set* $a := \pm\frac{1}{\sqrt{|1/\operatorname{ch}^2\frac{\theta}{2} - \operatorname{th}^2\frac{\psi}{2}|}}$*. Then there is the basis transformation*

$$\begin{pmatrix} \operatorname{ch}\phi & 0 & \operatorname{sh}\phi \\ 0 & 1 & 0 \\ \operatorname{sh}\phi & 0 & \operatorname{ch}\phi \end{pmatrix} \begin{pmatrix} a/\operatorname{ch}\frac{\theta}{2} \\ 0 \\ a\operatorname{th}\frac{\psi}{2} \end{pmatrix}, \quad \begin{array}{l} a).\ \textit{or}\ \operatorname{ch}\phi = \frac{a}{\operatorname{ch}\frac{\theta}{2}}, \quad \operatorname{sh}\phi = -a\operatorname{th}\frac{\psi}{2}, \\ b).\ \textit{or}\ \operatorname{sh}\phi = -\frac{a}{\operatorname{ch}\frac{\theta}{2}}, \quad \operatorname{ch}\phi = a\operatorname{th}\frac{\psi}{2}. \end{array} \tag{5.16}$$

In the case a)., the coordinate axis (1,0,0) is fixed, and in the case b)., the axis (0,0,1) is fixed. If $1/\operatorname{ch}^2\frac{\theta}{2} = \pm\operatorname{th}^2\frac{\psi}{2}$*, then it is impossible to simplify the initial superposition with the help of any pseudo-orthogonal rotation of the basis only.*

Example 5.2. *Let's consider a superposition of elementary transformations from the group* $O(1,2)$

$$\begin{pmatrix} \operatorname{ch}\phi & 0 & \operatorname{sh}\phi \\ 0 & 1 & 0 \\ \operatorname{sh}\phi & 0 & \operatorname{ch}\phi \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\alpha & \sin\alpha \\ 0 & -\sin\alpha & \cos\alpha \end{pmatrix} \begin{pmatrix} \operatorname{ch}\psi & \operatorname{sh}\psi & 0 \\ \operatorname{sh}\psi & \operatorname{ch}\psi & 0 \\ 0 & 0 & 1 \end{pmatrix} =$$

$$= \begin{pmatrix} \operatorname{ch}\tilde\phi & 0 & \operatorname{sh}\tilde\phi \\ 0 & 1 & 0 \\ \operatorname{sh}\tilde\phi & 0 & \operatorname{ch}\tilde\phi \end{pmatrix} \begin{pmatrix} \operatorname{ch}\tilde\psi & \operatorname{sh}\tilde\psi & 0 \\ \operatorname{sh}\tilde\psi & \operatorname{ch}\tilde\psi & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\tilde\alpha & \sin\tilde\alpha \\ 0 & -\sin\tilde\alpha & \cos\tilde\alpha \end{pmatrix}, \tag{5.17}$$

$$\begin{pmatrix} \operatorname{ch}(\phi-\tilde\phi)\operatorname{ch}\psi - \operatorname{sh}(\phi-\tilde\phi)\operatorname{sh}\psi\sin\alpha & \operatorname{ch}(\phi-\tilde\phi)\operatorname{sh}\psi - \operatorname{sh}(\phi-\tilde\phi)\operatorname{ch}\psi\sin\alpha & \operatorname{sh}(\phi-\tilde\phi)\cos\alpha \\ \cos\alpha\operatorname{sh}\psi & \cos\alpha\operatorname{ch}\psi & \sin\alpha \\ \operatorname{sh}(\phi-\tilde\phi)\operatorname{ch}\psi - \operatorname{ch}(\phi-\tilde\phi)\operatorname{sh}\psi\sin\alpha & \operatorname{sh}(\phi-\tilde\phi)\operatorname{sh}\psi - \operatorname{ch}(\phi-\tilde\phi)\operatorname{ch}\psi\sin\alpha & \operatorname{ch}(\phi-\tilde\phi)\cos\alpha \end{pmatrix} =$$

$$= \begin{pmatrix} \operatorname{ch}\tilde\psi & \operatorname{sh}\tilde\psi\cos\tilde\alpha & \operatorname{sh}\tilde\psi\sin\tilde\alpha \\ \operatorname{sh}\tilde\psi & \operatorname{ch}\tilde\psi\cos\tilde\alpha & \operatorname{ch}\tilde\psi\sin\tilde\alpha \\ 0 & -\sin\tilde\alpha & \cos\tilde\alpha \end{pmatrix}$$

The solution of this system will be

$$\begin{array}{l} \sin\alpha = \operatorname{ch}\tilde\psi\sin\tilde\alpha, \quad \operatorname{sh}\psi = \frac{\operatorname{sh}\tilde\psi}{\sqrt{1-\operatorname{ch}^2\tilde\psi\sin^2\tilde\alpha}}, \quad \operatorname{sh}(\phi-\tilde\phi) = \frac{\operatorname{sh}\tilde\psi\sin\tilde\alpha}{\sqrt{1-\operatorname{ch}^2\tilde\psi\sin^2\tilde\alpha}}. \\ \sin\tilde\alpha = \frac{\sin\alpha}{\sqrt{1+\operatorname{sh}^2\psi\cos^2\alpha}}, \quad \operatorname{sh}\tilde\psi = \operatorname{sh}\psi\cos\alpha, \quad \operatorname{sh}(\phi-\tilde\phi) = \frac{\operatorname{sh}\psi\sin\alpha}{\sqrt{1+\operatorname{sh}^2\psi\cos^2\alpha}}. \end{array} \tag{5.18}$$

Besides, an unusual case is $\operatorname{ch}^2\tilde\psi\sin^2\tilde\alpha \geq 1$*. Therefore, it is not always one can rearrange elementary transformations in a superposition unlike the orthogonal group* $O(3)$*.*

Example 5.3. *Let's consider transformations from the group $O(2,2)$. Any transformation from this group is presented by means of four transformations (5.2) according to the Cartan-Dieudonne theorem [10, v. 2, p. 33 (rus)]*

$$\pm\, r_{\tilde{\Lambda}} r^{\tilde{\Psi}} - \delta_{\tilde{\Lambda}}{}^{\tilde{\Psi}}, \quad r_{\tilde{\Lambda}} r^{\tilde{\Lambda}} = \pm 2. \tag{5.19}$$

However, always it is possible to pick up the decomposition so that two of such the transformations will have a diagonal matrix. Elements of the main diagonal of the matrix $I_{\tilde{\Lambda}}{}^{\tilde{\Psi}}$ on the module are equal to 1. Therefore, (5.2) can be copied as

$$\begin{gathered}(I_1)_{\tilde{\Phi}}{}^{\tilde{\Lambda}}(\pm(r_1)_{\tilde{\Lambda}}(r_1)^{\tilde{\Psi}} - \delta_{\tilde{\Lambda}}{}^{\tilde{\Psi}})(I_2)_{\tilde{\Psi}}{}^{\tilde{\Theta}}(\pm(r_2)_{\tilde{\Theta}}(r_2)^{\tilde{\Omega}} - \delta_{\tilde{\Theta}}{}^{\tilde{\Omega}}),\\ (r_1)^{\tilde{\Psi}}(I_2)_{\tilde{\Psi}}{}^{\tilde{\Theta}}(r_2)_{\tilde{\Theta}} = 0, \quad (r_2)^{\tilde{\Psi}}(I_1)_{\tilde{\Psi}}{}^{\tilde{\Theta}}(r_1)_{\tilde{\Theta}} = 0.\end{gathered} \tag{5.20}$$

There are significantly the various 3 mutually exclusive variants.

1. *In this case, a pseudo-orthogonal transformation can have the block-diagonal structure*

$$\begin{gathered}
I_1 \begin{pmatrix} \operatorname{ch}\theta & -\operatorname{sh}\theta & 0 & 0 \\ \operatorname{sh}\theta & -\operatorname{ch}\theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} I_2 \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \operatorname{ch}\phi & -\operatorname{sh}\phi \\ 0 & 0 & \operatorname{sh}\phi & -\operatorname{ch}\phi \end{pmatrix} = \\[2ex]
I_1 \left(\underbrace{\begin{pmatrix} \sqrt{2}\operatorname{sh}\frac{\theta}{2} \\ \sqrt{2}\operatorname{ch}\frac{\theta}{2} \\ 0 \\ 0 \end{pmatrix}}_{(r_1)_{\tilde{\Theta}}} \underbrace{\begin{pmatrix} \sqrt{2}\operatorname{sh}\frac{\theta}{2} & -\sqrt{2}\operatorname{ch}\frac{\theta}{2} & 0 & 0 \end{pmatrix}}_{(r_1)^{\tilde{\Omega}}} + \underbrace{\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\delta_{\tilde{\Theta}}{}^{\tilde{\Omega}}} \right) \\[2ex]
I_2 \left(\underbrace{\begin{pmatrix} 0 \\ 0 \\ \sqrt{2}\operatorname{sh}\frac{\phi}{2} \\ \sqrt{2}\operatorname{ch}\frac{\phi}{2} \end{pmatrix}}_{(r_2)_{\tilde{\Theta}}} \underbrace{\begin{pmatrix} 0 & 0 & \sqrt{2}\operatorname{sh}\frac{\phi}{2} & -\sqrt{2}\operatorname{ch}\frac{\phi}{2} \end{pmatrix}}_{(r_2)^{\tilde{\Omega}}} + \underbrace{\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\delta_{\tilde{\Theta}}{}^{\tilde{\Omega}}} \right), \\[2ex]
(r_1)^{\tilde{\Psi}}(r_1)_{\tilde{\Psi}} = -2, \quad (r_2)^{\tilde{\Psi}}(r_2)_{\tilde{\Psi}} = -2.
\end{gathered} \tag{5.21}$$

2. *In this case, a pseudo-orthogonal transformation can have the block-diagonal structure*

$$I_1 \begin{pmatrix} \cos\theta & \sin\theta & 0 & 0 \\ \sin\theta & -\cos\theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} I_2 \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -\cos\phi & -\sin\phi \\ 0 & 0 & -\sin\phi & \cos\phi \end{pmatrix} = \tag{5.22}$$

$$I_1\left(\underbrace{\begin{pmatrix}\sqrt{2}\cos\frac{\theta}{2}\\ \sqrt{2}\sin\frac{\theta}{2}\\ 0\\ 0\end{pmatrix}}_{(r_1)_{\tilde{\Theta}}}\underbrace{\begin{pmatrix}\sqrt{2}\cos\frac{\theta}{2} & \sqrt{2}\sin\frac{\theta}{2} & 0 & 0\end{pmatrix}}_{(r_1)^{\tilde{\Omega}}}-\underbrace{\begin{pmatrix}1&0&0&0\\0&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}}_{\delta_{\tilde{\Theta}}{}^{\tilde{\Omega}}}\right)$$

$$I_2\left(\underbrace{\begin{pmatrix}0\\ 0\\ -\sqrt{2}\cos\frac{\phi}{2}\\ -\sqrt{2}\sin\frac{\phi}{2}\end{pmatrix}}_{(r_2)_{\tilde{\Theta}}}\underbrace{\begin{pmatrix}0 & 0 & \sqrt{2}\cos\frac{\phi}{2} & \sqrt{2}\sin\frac{\phi}{2}\end{pmatrix}}_{(r_2)^{\tilde{\Omega}}}+\underbrace{\begin{pmatrix}1&0&0&0\\0&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}}_{\delta_{\tilde{\Theta}}{}^{\tilde{\Omega}}}\right),$$

$$(r_1)^{\tilde{\Psi}}(r_1)_{\tilde{\Psi}}=2,\quad (r_2)^{\tilde{\Psi}}(r_2)_{\tilde{\Psi}}=-2.$$

3. *In this case, a pseudo-orthogonal transformation will not have a block-diagonal structure, but can be presented as*

$$\begin{pmatrix}1&0&0&0\\0&1+\frac{a^2}{2}&a&-\frac{a^2}{2}\\0&a&1&-a\\0&\frac{a^2}{2}&a&1-\frac{a^2}{2}\end{pmatrix}\begin{pmatrix}-1&-b&0&b\\b&-1+\frac{b^2}{2}&0&-\frac{b^2}{2}\\0&0&-1&0\\b&\frac{b^2}{2}&0&-1-\frac{b^2}{2}\end{pmatrix}=$$

$$=\underbrace{\begin{pmatrix}1&0&0&0\\0&1&0&0\\0&0&-1&0\\0&0&0&1\end{pmatrix}}_{(I_1)_{\tilde{\Phi}}{}^{\tilde{\Lambda}}}\left(\underbrace{\begin{pmatrix}0\\ \frac{a}{\sqrt{2}}\\ -\sqrt{2}\\ \frac{a}{\sqrt{2}}\end{pmatrix}}_{(r_1)_{\tilde{\Lambda}}}\underbrace{\begin{pmatrix}0 & \frac{a}{\sqrt{2}} & \sqrt{2} & -\frac{a}{\sqrt{2}}\end{pmatrix}}_{(r_1)^{\tilde{\Psi}}}+\underbrace{\begin{pmatrix}1&0&0&0\\0&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}}_{\delta_{\tilde{\Lambda}}{}^{\tilde{\Psi}}}\right)$$

$$\underbrace{\begin{pmatrix}-1&0&0&0\\0&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}}_{(I_2)_{\tilde{\Psi}}{}^{\tilde{\Theta}}}\left(\underbrace{\begin{pmatrix}\sqrt{2}\\ \frac{b}{\sqrt{2}}\\ 0\\ \frac{b}{\sqrt{2}}\end{pmatrix}}_{(r_2)_{\tilde{\Theta}}}\underbrace{\begin{pmatrix}\sqrt{2} & \frac{b}{\sqrt{2}} & 0 & -\frac{b}{\sqrt{2}}\end{pmatrix}}_{(r_2)^{\tilde{\Omega}}}-\underbrace{\begin{pmatrix}1&0&0&0\\0&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}}_{\delta_{\tilde{\Theta}}{}^{\tilde{\Omega}}}\right),$$

$$(r_1)^{\tilde{\Psi}}(r_1)_{\tilde{\Psi}}=-2,\quad (r_2)^{\tilde{\Psi}}(r_2)_{\tilde{\Psi}}=2. \tag{5.23}$$

According to the same Cartan-Dieudonne theorem, any transformation from the group $O(2n,2n)$ in some basis will have the block-diagonal structure consisting of blocks $O(2,2)$. This means, that (5.2) is executed

$$S_{\tilde{\Lambda}_1}{}^{\tilde{\Lambda}_{J+1}}=\pm\prod_{I=1}^{J}(\pm(r_I)_{\tilde{\Lambda}_I}(r_I)^{\tilde{\Lambda}_{I+1}}-\delta_{\tilde{\Lambda}_I}{}^{\tilde{\Lambda}_{I+1}}). \tag{5.24}$$

Let now the canonical complex structure be set in the space $\mathbb{R}^{2n}_{(n,n)}$ then in the decomposition (5.2), one can allocate $J/2$ factors so that the relations

$$\begin{gathered} f_{\tilde{\Lambda}_{2I}}{}^{\tilde{\Lambda}_I}((r_I)_{\tilde{\Lambda}_I}(r_I)^{\tilde{\Lambda}_{I+1}} - \delta_{\tilde{\Lambda}_I}{}^{\tilde{\Lambda}_{I+1}})f_{\tilde{\Lambda}_{I+1}}{}^{\tilde{\Lambda}_{2I+1}} = (r_{2I})_{\tilde{\Lambda}_{2I}}(r_{2I})^{\tilde{\Lambda}_{2I+1}} + \delta_{\tilde{\Lambda}_{2I}}{}^{\tilde{\Lambda}_{2I+1}}, \\ (r_I)_{\tilde{\Lambda}}(r_I)^{\tilde{\Lambda}} = 2, \quad (r_{2I})_{\tilde{\Lambda}}(r_{2I})^{\tilde{\Lambda}} = -2, \quad \forall I = \overline{1, J/2} \end{gathered} \tag{5.25}$$

will be executed. For any pseudo-orthogonal transformation $S_{\tilde{\Lambda}}{}^{\tilde{\Psi}}$, one can define the parities

$$\begin{gathered} S_{\Lambda}{}^{\Psi} := m_{\Lambda}{}^{\tilde{\Lambda}} S_{\tilde{\Lambda}}{}^{\tilde{\Psi}} m^{\Psi}{}_{\tilde{\Psi}}, \quad S_{\Lambda}{}^{\Psi'} := m_{\Lambda}{}^{\tilde{\Lambda}} S_{\tilde{\Lambda}}{}^{\tilde{\Psi}} \bar{m}^{\Psi'}{}_{\tilde{\Psi}} = 0, \\ \bar{S}_{\Lambda'}{}^{\Psi} := \bar{m}_{\Lambda'}{}^{\tilde{\Lambda}} S_{\tilde{\Lambda}}{}^{\tilde{\Psi}} m^{\Psi}{}_{\tilde{\Psi}} = 0, \quad \bar{S}_{\Lambda'}{}^{\Psi'} := \bar{m}_{\Lambda'}{}^{\tilde{\Lambda}} S_{\tilde{\Lambda}}{}^{\tilde{\Psi}} \bar{m}^{\Psi'}{}_{\tilde{\Psi}}, \\ -f_{\tilde{\Omega}}{}^{\tilde{\Lambda}} S_{\tilde{\Lambda}}{}^{\tilde{\Psi}} f_{\tilde{\Psi}}{}^{\tilde{\Theta}} = S_{\tilde{\Omega}}{}^{\tilde{\Theta}}. \end{gathered} \tag{5.26}$$

Thus, a complex symmetry will be defined as

$$\begin{gathered} m_{\Theta}{}^{\tilde{\Theta}}(r_{\tilde{\Theta}} r^{\tilde{\Sigma}} - \delta_{\tilde{\Theta}}{}^{\tilde{\Sigma}}) f_{\tilde{\Sigma}}{}^{\tilde{\Psi}} (r_{\tilde{\Psi}} r^{\tilde{\Omega}} - \delta_{\tilde{\Psi}}{}^{\tilde{\Omega}}) f_{\tilde{\Omega}}{}^{\tilde{\Xi}} m^{\Xi}{}_{\tilde{\Xi}} = \\ = m_{\Theta}{}^{\tilde{\Theta}}(-r_{\tilde{\Theta}} r^{\tilde{\Sigma}} f_{\tilde{\Sigma}}{}^{\tilde{\Psi}} f_{\tilde{\Psi}}{}^{\tilde{\Xi}} - f_{\tilde{\Theta}}{}^{\tilde{\Psi}} r_{\tilde{\Psi}} r^{\tilde{\Omega}} f_{\tilde{\Omega}}{}^{\tilde{\Xi}} + f_{\tilde{\Theta}}{}^{\tilde{\Psi}} f_{\tilde{\Psi}}{}^{\tilde{\Xi}}) m^{\Xi}{}_{\tilde{\Xi}} = \\ = (2 r_{\Theta} r^{\Xi} - \delta_{\Theta}{}^{\Xi}), \end{gathered} \tag{5.27}$$

Besides,

$$r_{\Lambda} r^{\Lambda} = m_{\Lambda}{}^{\tilde{\Psi}} r_{\tilde{\Psi}} m^{\Lambda}{}_{\tilde{\Phi}} r^{\tilde{\Phi}} = \frac{1}{2}(\delta_{\tilde{\Phi}}{}^{\tilde{\Psi}} + i f_{\tilde{\Phi}}{}^{\tilde{\Psi}}) r_{\tilde{\Psi}} r^{\tilde{\Phi}} = \frac{1}{2} r_{\tilde{\Psi}} r^{\tilde{\Psi}} = 1, \tag{5.28}$$

that definitively confirms the existence of the decomposition

$$S_{\Lambda_1}{}^{\Lambda_{J+1}} = \pm \prod_{I=1}^{J} ((r_I)_{\Lambda_I}(r_I)^{\Lambda_{I+1}} - \delta_{\Lambda_I}{}^{\Lambda_{I+1}}), \quad (r_I)_{\Lambda_I}(r_I)^{\Lambda_I} = 2. \tag{5.29}$$

However, it is necessary to notice that not always $(r_I)_{\Lambda}(r_J)^{\Lambda} = 0, \ (I \neq J)$ in view of the existence of isotropic vectors as this has been shown in the previous example. As to the block structure of transformations from the group $O(n, \mathbb{C})$ that the dimension of the blocks will be equal to 4 too.

Example 5.4. *Let the three-dimensional pseudo-Euclidian space $\mathbb{R}^3_{(1,2)}$ be given. We will consider the transformation*

$$S_i{}^j = \begin{pmatrix} 1 + \frac{a^2}{2} & a & -\frac{a^2}{2} \\ a & 1 & -a \\ \frac{a^2}{2} & a & 1 - \frac{a^2}{2} \end{pmatrix}. \tag{5.30}$$

Let's define the inclusion operator

$$H^i{}_{\Lambda} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -i & 0 \\ 0 & 0 & -i \end{pmatrix}, \quad H_i{}^{\Lambda} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & i & 0 \\ 0 & 0 & i \end{pmatrix}. \tag{5.31}$$

It will define the complexification of our transformation

$$S_{\Lambda}{}^{\Psi} = H^i{}_{\Lambda} S_i{}^j H_j{}^{\Psi} = \begin{pmatrix} 1 + \frac{a^2}{2} & ia & -i\frac{a^2}{2} \\ -ia & 1 & -a \\ -i\frac{a^2}{2} & a & 1 - \frac{a^2}{2} \end{pmatrix} \tag{5.32}$$

with the complex Euclidian metric. To pass to the real representation, the operators

$$(m^T)_{\tilde{\varLambda}}{}^{\Lambda} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & i & 0 \\ 0 & 0 & i \\ -i & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad m_{\Lambda}{}^{\tilde{\varLambda}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 & i & 0 & 0 \\ 0 & -i & 0 & 0 & 1 & 0 \\ 0 & 0 & -i & 0 & 0 & 1 \end{pmatrix}, \tag{5.33}$$

will be required to us. Accordingly, the real representation of our transformation will look like

$$S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} = m^{\Lambda}{}_{\tilde{\varLambda}} S_{\Lambda}{}^{\Psi} m_{\Psi}{}^{\tilde{\varPsi}} + \bar{m}^{\Lambda'}{}_{\tilde{\varLambda}} S_{\Lambda'}{}^{\Psi'} \bar{m}_{\Psi'}{}^{\tilde{\varPsi}} = \begin{pmatrix} 1+\frac{a^2}{2} & a & -\frac{a^2}{2} & 0 & 0 & 0 \\ a & 1 & -a & 0 & 0 & 0 \\ \frac{a^2}{2} & a & 1-\frac{a^2}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1+\frac{a^2}{2} & a & -\frac{a^2}{2} \\ 0 & 0 & 0 & a & 1 & -a \\ 0 & 0 & 0 & \frac{a^2}{2} & a & 1-\frac{a^2}{2} \end{pmatrix}. \tag{5.34}$$

It is visible that such the representation has the block-diagonal structure, however, the complex representation has no such a structure.

Along with the transformations (5.26), it makes sense to consider the pseudo-orthogonal transformations

$$\begin{gathered} S_{\Lambda}{}^{\Psi} := m_{\Lambda}{}^{\tilde{\varLambda}} S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} m^{\Psi}{}_{\tilde{\varPsi}} = 0, \quad S_{\Lambda}{}^{\Psi'} := m_{\Lambda}{}^{\tilde{\varLambda}} S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} \bar{m}^{\Psi'}{}_{\tilde{\varPsi}}, \\ \bar{S}_{\Lambda'}{}^{\Psi} := \bar{m}_{\Lambda'}{}^{\tilde{\varLambda}} S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} m^{\Psi}{}_{\tilde{\varPsi}}, \quad S_{\Lambda'}{}^{\Psi'} := \bar{m}_{\Lambda'}{}^{\tilde{\varLambda}} S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} \bar{m}^{\Psi'}{}_{\tilde{\varPsi}} = 0, \\ f_{\tilde{\varOmega}}{}^{\tilde{\varLambda}} S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} f_{\tilde{\varPsi}}{}^{\tilde{\varTheta}} = S_{\tilde{\varOmega}}{}^{\tilde{\varTheta}}, \quad S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} S_{\tilde{\varPsi}}{}^{\tilde{\varPhi}} = \delta_{\tilde{\varLambda}}{}^{\tilde{\varPhi}}, \quad S_{\Lambda}{}^{\Psi'} \bar{S}_{\Psi'}{}^{\Phi} = \delta_{\Lambda}{}^{\Phi}, \quad S_{\Lambda}{}^{\Psi'} S_{\Phi}{}^{\Omega'} \bar{g}_{\Psi'\Omega'} = g_{\Lambda\Phi}. \end{gathered} \tag{5.35}$$

The complex representation of such the transformations will be an involution on the space $\mathbb{C}^n$ responsible for a real inclusion, therefore in the decomposition (5.2), one can allocate $J/2$ factors so that the relations

$$\begin{gathered} f_{\tilde{\varLambda}_{2I}}{}^{\tilde{\varPsi}_I} ((r_I)_{\tilde{\varPsi}_I} (r_I)^{\tilde{\varLambda}_{I+1}} - \delta_{\tilde{\varPsi}_I}{}^{\tilde{\varLambda}_{I+1}}) f_{\tilde{\varLambda}_{I+1}}{}^{\tilde{\varLambda}_{2I+1}} = -((r_{2I})_{\tilde{\varLambda}_{2I}} (r_{2I})^{\tilde{\varLambda}_{2I+1}} + \delta_{\tilde{\varLambda}_{2I}}{}^{\tilde{\varLambda}_{2I+1}}), \\ (r_I)_{\tilde{\varLambda}} (r_I)^{\tilde{\varLambda}} = 2, \quad (r_{2I})_{\tilde{\varLambda}} (r_{2I})^{\tilde{\varLambda}} = -2, \quad \forall I = \overline{1, J/2}. \end{gathered} \tag{5.36}$$

will be executed. Thus, eigenvalues of the transformation $S_{\Lambda}{}^{\Psi'}$ are equal to ±1. This automatically means, that such the transformation is orthogonally diagonalized in the complex representation, and the real representation in some basis has the block-diagonal structure

$$\begin{pmatrix} \cos\theta & \sin\theta & 0 & 0 & \ldots\ldots \\ \sin\theta & -\cos\theta & 0 & 0 & \ldots\ldots \\ 0 & 0 & -\cos\theta & -\sin\theta & \ldots\ldots \\ 0 & 0 & -\sin\theta & \cos\theta & \ldots\ldots \\ \ldots\ldots\ldots\ldots & \ldots\ldots\ldots\ldots & \ldots\ldots\ldots\ldots & \ldots\ldots\ldots\ldots & \ldots\ldots \end{pmatrix} \tag{5.37}$$

and also is diagonalized. It is necessary to address for more detailed statement to [20, pp. 79-126], [10, v. 2].

6 Clifford equation. Double covering

In this section, it is told about how to pass to the reduced Cartan spinors. This will allow to construct the one-to-one correspondence between vectors of the space $\mathbb{C}^n$ and bivectors of the spinor space $\mathbb{C}^N$ that will lead to the construction of the double covering $Spin(n,\mathbb{C})/\{\pm 1\} \cong SO(n,\mathbb{C})$. The conclusion of all results of this section is made on the basis [20], [22].

Suppose that a pseudo-Riemannian space V_{2n} is the real representation of the complex space $\mathbb{C}V_n$, where n is even. Let a set of non-degenerate operators $\gamma_{\tilde{\mathit{\Lambda}}}$ ($\tilde{\mathit{\Lambda}},\ \tilde{\mathit{\Psi}},\ ... = \overline{1,2n}$) be given at each point of V_{2n}. Moreover, we assume that each operator $\gamma_{\tilde{\mathit{\Lambda}}} := m_{\tilde{\mathit{\Lambda}}}{}^{\Lambda}\gamma_{\Lambda} + \bar{m}_{\tilde{\mathit{\Lambda}}}{}^{\Lambda'}\gamma_{\Lambda'}$ can be represented by a real matrix $(2N)^2 \times (2N)^2$, where $N = 2^{n/2-1}$. In addition, the operators must satisfy the condition (2.3)

$$\gamma_{\tilde{\mathit{\Lambda}}}\gamma_{\tilde{\mathit{\Psi}}} + \gamma_{\tilde{\mathit{\Psi}}}\gamma_{\tilde{\mathit{\Lambda}}} = G_{\tilde{\mathit{\Lambda}}\tilde{\mathit{\Psi}}}, \tag{6.1}$$

where $G_{\tilde{\mathit{\Lambda}}\tilde{\mathit{\Psi}}} = G_{\tilde{\mathit{\Psi}}\tilde{\mathit{\Lambda}}}$ is a non-degenerate metric tensor from (3.11)-(3.12) defined in the real representation of the tangent bundle V_{2n}. The operators (3.25) (if it does not matter which of the K-th operators will be selected then the number K will be omitted)

$$\gamma_{\Lambda} := \tilde{M}(m_{\Lambda}{}^{\tilde{\mathit{\Lambda}}}\gamma_{\tilde{\mathit{\Lambda}}})\tilde{\tilde{M}}^T \tag{6.2}$$

are the complex representation of the operators $\gamma_{\tilde{\mathit{\Lambda}}}$. They will satisfy the equation

$$\gamma_{\Lambda}\gamma_{\Psi} + \gamma_{\Psi}\gamma_{\Lambda} = g_{\Lambda\Psi}, \quad \bar{\gamma}_{\Lambda'}\bar{\gamma}_{\Psi'} + \bar{\gamma}_{\Psi'}\bar{\gamma}_{\Lambda'} = g_{\Lambda'\Psi'}. \tag{6.3}$$

Let r^{Λ} be a vector of the complex tangent bundle $\tau^{\mathbb{C}}$. As the base can be regarded both the complex representation $\mathbb{C}V_n$ and the real representation V_{2n}. A fiber of this bundle is isomorphic to the space $\mathbb{C}^n$, the real representation of which will be the pseudo-Euclidean space $R^{2n}_{(n,n)}$, where n is the index of the pseudo-Euclidean metric (the number of " + " on the main diagonal). Thus, a vector of the space $\mathbb{C}^n$ ($\mathbb{R}^{2n}_{(n,n)}$) can be uniquely associated to an operator from the space $\mathbb{C}^{2N}$ by the rule

$$R := r^{\Phi}\gamma_{\Phi}. \tag{6.4}$$

Because from the Clifford equation, the relation

$$tr(\gamma_{\Lambda}\gamma_{\Psi}) = N g_{\Lambda\Psi} \tag{6.5}$$

follows then each operator R can be uniquely associated to the vector of the space $\mathbb{C}^n(R^{2n}_{(n,n)})$ by means of the taking of the trace from the both parts of the conformity (6.4) with $\gamma^{\Lambda} := g^{\Lambda\Psi}\gamma_{\Psi}$

$$g^{\Lambda\Psi}tr(\gamma_{\Psi}R) = r^{\Phi}g^{\Lambda\Psi}tr(\gamma_{\Psi}\gamma_{\Phi}), \tag{6.6}$$

and then

$$\frac{1}{N}tr(\gamma^{\Psi}R) = r^{\Psi}. \tag{6.7}$$

Thus, it is obtained the one-to-one correspondence between the vectors r^{Λ} of the space $\mathbb{C}^n$ ($\mathbb{R}^{2n}_{(n,n)}$) and the operators R (special kind, of course) acting on the space $\mathbb{C}^{2N}$. Using the Clifford equation (6.3), we can write down the identity

$$R\gamma_{\Psi}R = r^{\Lambda}\gamma_{\Lambda}\gamma_{\Psi}\gamma_{\Phi}r^{\Phi} = r^{\Lambda}(g_{\Lambda\Psi} - \gamma_{\Psi}\gamma_{\Lambda})\gamma_{\Phi}r^{\Phi} = (r_{\Psi}r^{\Phi} - \frac{1}{2}(r_{\Lambda}r^{\Lambda})\delta_{\Psi}{}^{\Phi})\gamma_{\Phi}. \tag{6.8}$$

Let $r_\Lambda r^\Lambda := \pm 2$. It is known that any (pseudo-) orthogonal transformation ${S_\Psi}^\Phi$ can be represented as the finite product of the elementary transformations $S_I := (r_I)_\Psi (r_I)^\Phi - \frac{1}{2}((r_I)_\Lambda (r_I)^\Lambda){\delta_\Psi}^\Phi$ (I runs over the finite values from 1 to any finite J, for example). The previous section is devoted to the proof of this fact. Define

$$R_I := (r_I)^\Lambda \gamma_\Lambda, \quad S = \prod_{I=1}^{J} S_I = S_1 S_2 \ldots S_J, \tag{6.9}$$

$$\tilde{S} = \prod_{I=1}^{J} R_I = R_1 R_2 \ldots R_J, \quad \tilde{\tilde{S}} := \prod_{I=J}^{1} R_I = R_J \ldots R_2 R_1. \tag{6.10}$$

Here $\tilde{S}$ is the right product of the operators R_I, and $\tilde{\tilde{S}}$ is the left product of the operators R_I. Therefore, the matrixes of the operators $\tilde{S}$ and $\tilde{\tilde{S}}$ are significantly various and in any way with each other are not connected. It will allow to copy the conformity (6.8) for the orthogonal transformation ${S_\Lambda}^\Psi$ as

$${S_\Lambda}^\Psi \gamma_\Psi = \tilde{S} \gamma_\Lambda \tilde{\tilde{S}}, \quad \tilde{\tilde{S}} = \tilde{S}^{-1}. \tag{6.11}$$

The equation (6.8) is true for the real representation that will give

$${S_{\tilde{\varLambda}}}^{\tilde{\varPsi}} \gamma_{\tilde{\varPsi}} = \tilde{S} \gamma_{\tilde{\varLambda}} \tilde{\tilde{S}}, \quad {f_{\tilde{\varTheta}}}^{\tilde{\varLambda}} {S_{\tilde{\varLambda}}}^{\tilde{\varPsi}} {f_{\tilde{\varPsi}}}^{\tilde{\varPhi}} = -{S_{\tilde{\varTheta}}}^{\tilde{\varPhi}}. \tag{6.12}$$

Accordingly, for the involution, the equation (6.12) can be rewritten as

$${S_{\tilde{\varLambda}}}^{\tilde{\varPsi}} \gamma_{\tilde{\varPsi}} = \tilde{S} \gamma_{\tilde{\varLambda}} \tilde{\tilde{S}}, \quad {f_{\tilde{\varTheta}}}^{\tilde{\varLambda}} {S_{\tilde{\varLambda}}}^{\tilde{\varPsi}} {f_{\tilde{\varPsi}}}^{\tilde{\varPhi}} = {S_{\tilde{\varTheta}}}^{\tilde{\varPhi}}, \quad {S_{\tilde{\varLambda}}}^{\tilde{\varPsi}} {S_{\tilde{\varPsi}}}^{\tilde{\varPhi}} = {\delta_{\tilde{\varLambda}}}^{\tilde{\varPhi}}. \tag{6.13}$$

Then for the complex representation, we obtain the identity

$${S_\Lambda}^{\Psi'} \bar{\gamma}_{\Psi'} = \bar{\tilde{S}} \gamma_\Lambda \tilde{\tilde{S}}, \quad {S_\Lambda}^{\Psi'} {\bar{S}_{\Psi'}}{}^{\Phi} = {\delta_\Lambda}^\Phi, \quad \tilde{S} \bar{\tilde{S}} = \pm E, \quad \tilde{\tilde{S}} \bar{\tilde{\tilde{S}}} = \pm E. \tag{6.14}$$

It is now possible to complexify the real representation $\mathbb{R}^{2n}_{(n,n)}$ considering the vector $r^{\tilde{\varLambda}}$ in the same basis, but with complex coefficients. In this case, $\gamma_{\tilde{\varLambda}}$ will remain the same and ${S_{\tilde{\varLambda}}}^{\tilde{\varPsi}}$ from (6.12) will be a complex transformation. In particular, as such a transformation, we can consider the transformation $i{f_{\tilde{\varPsi}}}^{\tilde{\varPhi}}$ (see (3.14)). This means that the complex structure can be expanded as

$$i{f_{\tilde{\varLambda}}}^{\tilde{\varPsi}} \gamma_{\tilde{\varPsi}} = \tilde{I} \tilde{F} \gamma_{\tilde{\varLambda}} \tilde{\tilde{F}} \tilde{\tilde{I}}. \tag{6.15}$$

For further investigations of this correspondence, it is necessary to study the structure of the operators $\gamma_{\tilde{\varLambda}}$. Without loss of generality, we can consider a basis in which the metric tensor $G_{\tilde{\varLambda}\tilde{\varPsi}}$ on the main diagonal has ±1, and all remaining components are equal to 0

$$\gamma_{\tilde{\varLambda}} \gamma_{\tilde{\varPsi}} = -\gamma_{\tilde{\varPsi}} \gamma_{\tilde{\varLambda}}, \quad \tilde{\varLambda} \neq \tilde{\varPsi}. \tag{6.16}$$

Construct the operator

$$\gamma_0 := \prod_{\tilde{\varLambda}=1}^{2n} \gamma_{\tilde{\varLambda}} \tag{6.17}$$

according to [22, v. 2, Appendix, p. 442 (eng), (B.8)]. Then from (6.16), the relations

$$\gamma_0\gamma_{\tilde\Lambda} = -\gamma_{\tilde\Lambda}\gamma_0, \qquad (6.18)$$
$$(\gamma_0 - \lambda E)\gamma_{\tilde\Lambda} = -\gamma_{\tilde\Lambda}(\gamma_0 + \lambda E)$$

will follow. Therefor,

$$det(\gamma_0 - \lambda E) = 0 = det(\gamma_0 + \lambda E) \qquad (6.19)$$

will be the equation for the eigenvalues of γ_0. It is seen that these values are the paired: $\pm\lambda$. Then positive and negative eigenvalues correspond to the two different eigenspaces. Then there is a basis in which a block representation of the operator γ_0 (the blocks have the dimension $2N^2 \times 2N^2$)

$$\gamma_0 = \begin{pmatrix} \zeta_0 & 0 \\ 0 & \xi_0 \end{pmatrix} \qquad (6.20)$$

can exist, where ζ_0 responsible for the positive eigenvalues, and ξ_0 responsible for the negative eigenvalues. Define

$$\gamma_{\tilde\Lambda} = \begin{pmatrix} \zeta_{\tilde\Lambda} & \sigma_{\tilde\Lambda} \\ \eta_{\tilde\Lambda} & \xi_{\tilde\Lambda} \end{pmatrix}. \qquad (6.21)$$

Taking into account this definition, the condition (6.16) will lead to the relations

$$\zeta_0\zeta_{\tilde\Lambda} = -\zeta_{\tilde\Lambda}\zeta_0, \quad \xi_0\xi_{\tilde\Lambda} = -\xi_{\tilde\Lambda}\xi_0. \qquad (6.22)$$

This means that ζ_0 and ξ_0 should have pairs of the positive and negative values (even if $\zeta_{\tilde\Lambda}$ and $\xi_{\tilde\Lambda}$ are degenerate, but not equal to 0). Therefore, $\zeta_{\tilde\Lambda} = \xi_{\tilde\Lambda} \equiv 0$. Thus,

$$\gamma_{\tilde\Lambda} = \begin{pmatrix} 0 & \sigma_{\tilde\Lambda} \\ \eta_{\tilde\Lambda} & 0 \end{pmatrix}, \qquad (6.23)$$

and the equation (6.1) is equivalent to the system

$$\begin{cases} \eta_{\tilde\Lambda}\sigma_{\tilde\Psi} + \eta_{\tilde\Psi}\sigma_{\tilde\Lambda} = G_{\tilde\Lambda\tilde\Psi}, \\ \sigma_{\tilde\Lambda}\eta_{\tilde\Psi} + \sigma_{\tilde\Psi}\eta_{\tilde\Lambda} = G_{\tilde\Lambda\tilde\Psi}. \end{cases} \qquad (6.24)$$

Note that if we pass to an other basis in the space $\mathbb{R}^{(2N)^2}$ with a non-degenerate operator S from the general linear group $\tilde\gamma_{\tilde\Lambda} := S\gamma_{\tilde\Lambda}S^{-1}$ then the Clifford equation (6.1), generally speaking, will not save its form.

We now turn to the complex structure. Its square is equal to $-E$; this matrix is orthogonal. According to (6.24), for n=2, N=1, such the transformation is represented by means of the formula (6.12) as

$$\tilde F^2 = \tilde{\tilde F}^2 = \begin{pmatrix} -E & 0 \\ 0 & E \end{pmatrix}. \qquad (6.25)$$

If the complex structure $f_{\tilde\Lambda}{}^{\tilde\Psi}$ has the canonical form in the given basis then in the decomposition (6.15), $F := \tilde F = \tilde{\tilde F}$, $I := \tilde I = \tilde{\tilde I}$ has a block structure too (due to $\gamma_{\tilde\Lambda}$ having a block structure and the conformity (6.8) written for the real representation)

$$F = \begin{pmatrix} \hat F & 0 \\ 0 & \hat{\hat F}^T \end{pmatrix}, \quad \hat F^2 = -E, \quad \hat{\hat F}^2 = E, \quad I = \begin{pmatrix} iE & 0 \\ 0 & E \end{pmatrix}. \qquad (6.26)$$

With regard to the transformation I, it is determined from the relationship

$$i\gamma_\Lambda = \frac{1}{2}(i\gamma_\Lambda + (-i)(-\gamma_\Lambda)) = \underbrace{\frac{1}{2}((1+i)E + (1-i)\tilde{F}^2)}_{:=\tilde{I}}\,\gamma_\Lambda\,\underbrace{\frac{1}{2}((1+i)E + (1-i)\tilde{\tilde{F}}^2)}_{:=\tilde{\tilde{I}}}\,. \tag{6.27}$$

Whereas, the quantity of the elementary transformations, making the complex structure $f_{\tilde{\Lambda}}{}^{\tilde{\Psi}}$, is always multiple to 2 (since the dimension 2n of the real representation for even n is multiple to 4) then the block structure will be on the main diagonal. Hence, for the block components of $\gamma_{\tilde{\Lambda}}$, the relation

$$f_{\tilde{\Lambda}}{}^{\tilde{\Psi}}\sigma_{\tilde{\Psi}} = \hat{F}\sigma_{\tilde{\Lambda}}\hat{\hat{F}}^T, \quad f_{\tilde{\Lambda}}{}^{\tilde{\Psi}}\eta_{\tilde{\Psi}} = \hat{\hat{F}}^T\eta_{\tilde{\Lambda}}\hat{F} \tag{6.28}$$

will be true that defines the positive representation of (3.19) for the top sign and the negative representation of (3.19) for the lower sign. If we will set ($\tilde{A}, \tilde{B}, \tilde{C}, \tilde{D}, ... = \overline{1,4},\ \hat{A}, \hat{B}, \hat{C}, \hat{D}, ... = \overline{1,2},\ \tilde{\tilde{A}}, \tilde{\tilde{B}}, \tilde{\tilde{C}}, \tilde{\tilde{D}}, ... = \overline{1,2},\ A, B, C, D... = \overline{1,1}$)

$$\tilde{m}_{\pm\hat{A}}{}^{\tilde{B}} := \begin{pmatrix} \tilde{m}_{\pm A_1}{}^{\tilde{\tilde{B}}_1} & 0 \\ 0 & \tilde{\tilde{m}}_{\pm}{}^{A_2}{}_{\tilde{\tilde{B}}_2} \end{pmatrix}, \quad \tilde{\tilde{m}}_{\pm\hat{A}}{}^{\tilde{B}} := \begin{pmatrix} \tilde{m}_{\pm A_1}{}^{\tilde{\tilde{B}}_1} & 0 \\ 0 & \tilde{\tilde{m}}_{\pm}{}^{A_2}{}_{\tilde{\tilde{B}}_2} \end{pmatrix} \tag{6.29}$$

then the condition (3.19) will take the form

$$\left\{\begin{array}{l} (\tilde{m}_\pm)_A{}^{\tilde{\tilde{B}}}(\tilde{m}_\pm)^C{}_{\tilde{\tilde{B}}} = \delta_A{}^C, \\ \\ (\tilde{m}_\pm)_A{}^{\tilde{\tilde{B}}}(\tilde{m}_\mp)^C{}_{\tilde{\tilde{B}}} = 0, \\ \\ (\tilde{m}_\pm)_A{}^{\tilde{\tilde{B}}}(\tilde{m}_\pm)^A{}_{\tilde{\tilde{C}}} = \\ \qquad = \underbrace{\frac{1}{2}(\delta_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} \pm i\hat{F}_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}})}_{:=(\tilde{\triangle}_\pm)_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}}, \end{array}\right. \qquad \left\{\begin{array}{l} (\tilde{\tilde{m}}_\pm)_A{}^{\tilde{\tilde{B}}}(\tilde{\tilde{m}}_\pm)^C{}_{\tilde{\tilde{B}}} = \delta_A{}^C, \\ \\ (\tilde{\tilde{m}}_\pm)_A{}^{\tilde{\tilde{B}}}(\tilde{\tilde{m}}_\mp)^C{}_{\tilde{\tilde{B}}} = 0, \\ \\ (\tilde{\tilde{m}}_\pm)_A{}^{\tilde{\tilde{B}}}(\tilde{\tilde{m}}_\pm)_\pm{}^A{}_{\tilde{\tilde{C}}} = \\ \qquad = \underbrace{\frac{1}{2}(\delta_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} \pm \hat{\hat{F}}_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}})}_{:=(\tilde{\tilde{\triangle}}_\pm)_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}}\,. \end{array}\right. \tag{6.30}$$

Algorithm 6.1. *To construct the step operators (6.30) for an arbitrary even n, we use the algorithm 3.1 (*$N = 2^{n/2-1}$*,* $I = \overline{1, \frac{n}{2}}$*,* $\tilde{A}, \tilde{B}, \tilde{C}, \tilde{D}, ... = \overline{1, \frac{(2N)^2}{2^{I-1}}}$*,* $\hat{A}, \hat{B}, \hat{C}, \hat{D}, ... = \overline{1, \frac{(2N)^2}{2^I}}$*,* $\tilde{\tilde{A}}, \tilde{\tilde{B}}, \tilde{\tilde{C}}, \tilde{\tilde{D}}, ... = \overline{1, \frac{2N^2}{2^{I-1}}}$*,* $A, B, C, D... = \overline{1, \frac{2N^2}{2^I}}$*)*

$$\left\{\begin{array}{l} (\tilde{m}_{I_\pm})_A{}^{\tilde{\tilde{B}}}(\tilde{m}_{I_\pm})^C{}_{\tilde{\tilde{B}}} = \delta_A{}^C, \\ \\ (\tilde{m}_{I_\pm})_A{}^{\tilde{\tilde{B}}}(\tilde{m}_{I_\mp})^C{}_{\tilde{\tilde{B}}} = 0, \\ \\ (\tilde{m}_{I_\pm})_A{}^{\tilde{\tilde{B}}}(\tilde{m}_{I_\pm})^A{}_{\tilde{\tilde{C}}} = (\tilde{\triangle}_{I_\pm})_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}, \end{array}\right. \qquad \left\{\begin{array}{l} (\tilde{\tilde{m}}_{I_\pm})_A{}^{\tilde{\tilde{B}}}(\tilde{\tilde{m}}_{I_\pm})^C{}_{\tilde{\tilde{B}}} = \delta_A{}^C, \\ \\ (\tilde{\tilde{m}}_{I_\pm})_A{}^{\tilde{\tilde{B}}}(\tilde{\tilde{m}}_{I_\mp})^C{}_{\tilde{\tilde{B}}} = 0, \\ \\ (\tilde{\tilde{m}}_{I_\pm})_A{}^{\tilde{\tilde{B}}}(\tilde{\tilde{m}}_{I_\pm})_\pm{}^A{}_{\tilde{\tilde{C}}} = (\tilde{\tilde{\triangle}}_{I_\pm})_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}. \end{array}\right. \tag{6.31}$$

Thus,

$$(\tilde{\triangle}_{I_\pm})_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{A}}} := \frac{1}{2}(\delta_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{A}}} \pm \hat{I}_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{K}}}\hat{F}_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}), \quad (\tilde{\tilde{\triangle}}_{I_\pm})_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{A}}} := \frac{1}{2}(\delta_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{A}}} \pm \hat{\hat{I}}_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{K}}}\hat{\hat{F}}_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}). \tag{6.32}$$

It is now possible to define the operators ($\tilde{\tilde{A}}, \tilde{\tilde{B}}, \tilde{\tilde{C}}, \tilde{\tilde{D}}, ... = \overline{1, 2N^2},\ A, B, C, D... = \overline{1, N},\ K \neq J$)

$$\begin{gathered}
\tilde{M}_K := \tilde{m}_{\frac{n}{2}{z_{\frac{n}{2}}}} \tilde{m}_{\frac{n}{2}-1_{z_{\frac{n}{2}-1}}} \cdot ... \cdot \tilde{m}_{2_{z_2}} \tilde{m}_{1_{z_1}}, \\
(\tilde{M}_K)_C{}^{\tilde{\tilde{A}}} (\tilde{M}^*_K)^B{}_{\tilde{\tilde{A}}} = \delta_C{}^B, \\
(\tilde{M}_K)_C{}^{\tilde{\tilde{A}}} (\tilde{M}^*_J)^B{}_{\tilde{\tilde{A}}} = 0, \\
\sum_{K=1}^{2N} (\tilde{M}_K)_A{}^{\tilde{\tilde{C}}} (\tilde{M}^*_K)^A{}_{\tilde{\tilde{B}}} = \delta_{\tilde{\tilde{B}}}{}^{\tilde{\tilde{C}}},
\end{gathered}
\qquad
\begin{gathered}
\tilde{\tilde{M}}_K := \tilde{\tilde{m}}_{\frac{n}{2}{z_{\frac{n}{2}}}} \tilde{\tilde{m}}_{\frac{n}{2}-1_{z_{\frac{n}{2}-1}}} \cdot ... \cdot \tilde{\tilde{m}}_{2_{z_2}} \tilde{\tilde{m}}_{1_{z_1}}, \\
(\tilde{\tilde{M}}_K)_C{}^{\tilde{\tilde{A}}} (\tilde{\tilde{M}}^*_K)^B{}_{\tilde{\tilde{A}}} = \delta_C{}^B, \\
(\tilde{\tilde{M}}_K)_C{}^{\tilde{\tilde{A}}} (\tilde{\tilde{M}}^*_J)^B{}_{\tilde{\tilde{A}}} = 0, \\
\sum_{K=1}^{2N} (\tilde{\tilde{M}}_K)_A{}^{\tilde{\tilde{C}}} (\tilde{\tilde{M}}^*_K)^A{}_{\tilde{\tilde{B}}} = \delta_{\tilde{\tilde{B}}}{}^{\tilde{\tilde{C}}},
\end{gathered}
\tag{6.33}$$

where z_J ($J = \overline{1, \frac{n}{2}}$) *is equal to 0 for the sign* $\ll - \gg$ *or 1 for the sign* $\ll + \gg$ *then* $K = \sum_{J=1}^{\frac{n}{2}} z_J \cdot 2^{J-1} + 1$. *If it does not matter which of the K-th operators will be selected then the number K will be omitted. Using these operators, it is possible to define* ($\tilde{M} := \tilde{M}_C{}^{\tilde{\tilde{A}}}$, $\tilde{M}^* := \tilde{M}^{*C}{}_{\tilde{\tilde{A}}}$, $\tilde{\tilde{M}} := \tilde{\tilde{M}}_C{}^{\tilde{\tilde{A}}}$, $\tilde{\tilde{M}}^* := \tilde{\tilde{M}}^{*C}{}_{\tilde{\tilde{A}}}$)

$$\sigma_\Lambda := \tilde{M}(m_\Lambda{}^{\tilde{\Lambda}} \sigma_{\tilde{\Lambda}}) \tilde{\tilde{M}}^T, \quad \eta_\Lambda := \tilde{\tilde{M}}^*(m_\Lambda{}^{\tilde{\Lambda}} \eta_{\tilde{\Lambda}})(\tilde{M}^*)^T. \tag{6.34}$$

For further calculations, we must impose the condition

$$tr(\eta_{\tilde{\Psi}} \sigma_{\tilde{\Phi}}) = \frac{N}{2} g_{\tilde{\Psi}\tilde{\Phi}}. \tag{6.35}$$

Going to the complex representation in (6.24) with the help of the operators (6.33), we can obtain

$$\left\{ \begin{array}{c} \eta_\Lambda \sigma_\Psi + \eta_\Psi \sigma_\Lambda = g_{\Lambda\Psi}, \\ \sigma_\Lambda \eta_\Psi + \sigma_\Psi \eta_\Lambda = g_{\Lambda\Psi}, \\ tr(\eta_\Psi \sigma_\Lambda) = \frac{N}{2} g_{\Psi\Lambda}. \end{array} \right. \tag{6.36}$$

According to (6.4), this will lead to

$$R_\eta := r^\Lambda \eta_\Lambda, \quad R_\sigma := r^\Lambda \sigma_\Lambda, \quad \frac{2}{N} tr((R_\sigma)\eta_\Lambda) = r_\Lambda, \quad \frac{2}{N} tr((R_\eta)\sigma_\Lambda) = r_\Lambda, \tag{6.37}$$

$$R = \begin{pmatrix} 0 & R_\sigma \\ R_\eta & 0 \end{pmatrix}, \tag{6.38}$$

where the tensors $R_\eta,\ R_\sigma$ have the dimension $N \times N$. Hence, (6.8) is separated into the pair of the equations

$$R_\sigma \eta_\Psi R_\sigma = (r_\Psi r^\Phi - \frac{1}{2}(r_\Omega r^\Omega)\delta_\Psi{}^\Phi)\sigma_\Phi, \quad R_\eta \sigma_\Psi R_\eta = (r_\Psi r^\Phi - \frac{1}{2}(r_\Omega r^\Omega)\delta_\Psi{}^\Phi)\eta_\Phi. \tag{6.39}$$

Since

$$R_\sigma R_\eta = \frac{1}{2}(r_\Omega r^\Omega)E, \tag{6.40}$$

then one can determine the formalization of the operators η_Λ , σ_Λ as follows

$$\eta_\Lambda := \eta_\Lambda{}^{AB}, \quad \sigma_\Lambda := \sigma_{\Lambda AB} = \eta_{\Lambda BA}, \quad (A, B, ... = \overline{1, N}). \tag{6.41}$$

Then (6.11) can be rewritten as

$$I).\ S_\Psi{}^\Lambda \eta_\Lambda = \tilde{S}^T \eta_\Psi \tilde{\tilde{S}}, \quad II).\ S_\Psi{}^\Lambda \eta_\Lambda = \tilde{\tilde{S}}^T \sigma_\Psi \tilde{S}, \tag{6.42}$$

but here the operators $\tilde{\tilde{S}}$ and $\tilde{S}$ have the dimension $N \times N$. However, the decomposition of these operators for the special transformation $S_\Lambda{}^\Psi$ (det $\| S_\Lambda{}^\Psi \| = 1$) (case I) will have the form

$$\tilde{S}^T = \prod_{I=1}^{J} (R_\eta)_{2I-1}(R_\sigma)_{2I} = (R_\eta)_1 (R_\sigma)_2 (R_\eta)_3 (R_\sigma)_4 \dots (R_\eta)_{2J-1} (R_\sigma)_{2J}, \tag{6.43}$$

$$\tilde{\tilde{S}} = \prod_{I=J}^{1} (R_\sigma)_{2I}(R_\eta)_{2I-1} = (R_\sigma)_{2J} (R_\eta)_{2J-1} \dots (R_\sigma)_4 (R_\eta)_3 (R_\sigma)_2 (R_\eta)_1, \tag{6.44}$$

and for the non-special transformation $S_\Lambda{}^\Psi$ (det $\| S_\Lambda{}^\Psi \| = -1$) (case II) will have the form

$$\tilde{\tilde{S}}^T = (R_\eta)_1 \prod_{I=1}^{J} (R_\sigma)_{2I}(R_\eta)_{2I+1} = (R_\eta)_1 (R_\sigma)_2 (R_\eta)_3 \dots (R_\sigma)_{2J} (R_\eta)_{2J+1}, \tag{6.45}$$

$$\tilde{S} = (\prod_{I=J}^{1} (R_\eta)_{2I+1}(R_\sigma)_{2I})(R_\eta)_1 = (R_\eta)_{2J+1} (R_\sigma)_{2J} \dots (R_\eta)_3 (R_\sigma)_2 (R_\eta)_1. \tag{6.46}$$

Thus, the equation (6.42) defines the algebraic realization of the double covering $Spin(n, \mathbb{C})/\{\pm 1\} \cong SO(n, \mathbb{C})$. Accordingly, for the involution, the condition (6.14) determines

$$\begin{array}{ll} I). \quad S_\Psi{}^{\Lambda'} \eta_{\Lambda'} = \tilde{S}^T \eta_\Psi \tilde{\tilde{S}}, & II). \quad S_\Psi{}^{\Lambda'} \eta_{\Lambda'} = \tilde{\tilde{S}}^T \sigma_\Psi \tilde{S}, \\ \quad \tilde{S}\bar{\tilde{S}} = \pm E, \quad \tilde{\tilde{S}}\bar{\tilde{\tilde{S}}} = \pm E. & \quad \tilde{S} = \pm \bar{\tilde{S}}, \quad \tilde{\tilde{S}} = \pm \bar{\tilde{\tilde{S}}}. \end{array} \tag{6.47}$$

The bottom line of the parities are inherited by the real representation of the involution in the tangent and spinor spaces. But the real representation of an involution in the tangent space is an orthogonal transformation, the square of which has the identity matrix. Accordingly, in the real representation of the spinor space, this property inherits, and hence the square of the spinor involution in the real representation (case I) up to sign is the identity transformation too. It remains to pass to the complex representation on the rule (3.20).

Example 6.1. *2n=2, N=1. if* $g_{\tilde{\Lambda}\tilde{\Psi}} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, *then*

$$\begin{array}{c} \gamma_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}. \\ \eta_1 = \frac{1}{\sqrt{2}}, \quad \eta_2 = -\frac{i}{\sqrt{2}}, \quad \sigma_1 = \frac{1}{\sqrt{2}}, \quad \sigma_2 = \frac{i}{\sqrt{2}}. \\ tr(\eta_{\tilde{\Lambda}} \sigma_{\tilde{\Psi}}) = \frac{1}{2} \begin{pmatrix} 1 & i \\ -i & 1 \end{pmatrix}, \quad \text{- \textit{the tensor is degenerated.}} \end{array} \tag{6.48}$$

Thus, the identity $tr(\eta_{\tilde\Lambda}\sigma_{\tilde\Psi}) = \frac{1}{2}g_{\tilde\Lambda\tilde\Psi}$ *is false. Nevertheless, we can use (6.11) and (6.12). Define*

$$\begin{gathered}
(r_1)_{\tilde\Lambda} = (r_1)^{\tilde\Lambda} := \sqrt{2}(\cos\phi\ ,\ \sin\phi) \\
(S_1)_{\tilde\Lambda}{}^{\tilde\Psi} = \begin{pmatrix} 2\cos^2\phi - 1 & 2\cos\phi\sin\phi \\ 2\cos\phi\sin\phi & -1+2\sin^2\phi \end{pmatrix} = \begin{pmatrix} \cos 2\phi & \sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix} = (r_1)^{\tilde\Psi}(r_1)_{\tilde\Lambda} - \delta_{\tilde\Lambda}{}^{\tilde\Psi}, \\
R_1 = (r_1)^{\tilde\Lambda}\gamma_{\tilde\Lambda} = \cos\phi \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + \sin\phi \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}.
\end{gathered} \tag{6.49}$$

Then for any vector $x^{\tilde\Lambda} = (T, X)$*, the conformity* $X = x^{\tilde\Lambda}\gamma_{\tilde\Lambda}$*,* $(S_1)_{\tilde\Lambda}{}^{\tilde\Psi}x_{\tilde\Psi} \longleftrightarrow R_1 X R_1$ *will have the form*

$$\begin{pmatrix} \cos 2\phi & \sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix}\begin{pmatrix} T \\ X \end{pmatrix} \longleftrightarrow \begin{pmatrix} 0 & e^{i\phi} \\ e^{-i\phi} & 0 \end{pmatrix}\begin{pmatrix} 0 & T+iX \\ T-iX & 0 \end{pmatrix}\begin{pmatrix} 0 & e^{i\phi} \\ e^{-i\phi} & 0 \end{pmatrix}. \tag{6.50}$$

Define

$$\begin{gathered}
(r_2)_{\tilde\Lambda} = (r_2)^{\tilde\Lambda} := \sqrt{2}(\cos\phi,\quad -\sin\phi). \\
(S_2)_{\tilde\Lambda}{}^{\tilde\Psi} = \begin{pmatrix} \cos 2\phi & -\sin 2\phi \\ -\sin 2\phi & -\cos 2\phi \end{pmatrix} = (r_2)^{\tilde\Psi}(r_2)_{\tilde\Lambda} - \delta_{\tilde\Lambda}{}^{\tilde\Psi}, \\
R_2 = (r_2)^{\tilde\Lambda}\gamma_{\tilde\Lambda} = \cos\phi \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} - \sin\phi \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}.
\end{gathered} \tag{6.51}$$

Then for any vector $x^{\tilde\Lambda} = (T, X)$*, the conformity* $X = x^{\tilde\Lambda}\gamma_{\tilde\Lambda}$*,* $(S_2)_{\tilde\Lambda}{}^{\tilde\Psi}x_{\tilde\Psi} \longleftrightarrow R_2 X R_2$ *will have the form*

$$\begin{pmatrix} \cos 2\phi & -\sin 2\phi \\ -\sin 2\phi & -\cos 2\phi \end{pmatrix}\begin{pmatrix} T \\ X \end{pmatrix} \longleftrightarrow \begin{pmatrix} 0 & e^{-i\phi} \\ e^{i\phi} & 0 \end{pmatrix}\begin{pmatrix} 0 & T+iX \\ T-iX & 0 \end{pmatrix}\begin{pmatrix} 0 & e^{-i\phi} \\ e^{i\phi} & 0 \end{pmatrix}. \tag{6.52}$$

It is now possible to construct the special transformation $S := S_1 S_2$ *for which* $S_{\tilde\Lambda}{}^{\tilde\Psi}x_{\tilde\Psi} \longleftrightarrow R_1 R_2 X R_2 R_1$ *and* $\tilde{S} = R_1 R_2$*,* $\tilde{\tilde{S}} = R_2 R_1$*. The explicit record of the conformity will have the form*

$$\begin{pmatrix} \cos 4\phi & -\sin 4\phi \\ \sin 4\phi & \cos 4\phi \end{pmatrix}\begin{pmatrix} T \\ X \end{pmatrix} \longleftrightarrow \begin{pmatrix} e^{2i\phi} & 0 \\ 0 & e^{-2i\phi} \end{pmatrix}\begin{pmatrix} 0 & T+iX \\ T-iX & 0 \end{pmatrix}\begin{pmatrix} e^{-2i\phi} & 0 \\ 0 & e^{2i\phi} \end{pmatrix}. \tag{6.53}$$

It is obvious that $SO(2)$ *is formed by such the transformations* $S_{\tilde\Lambda}{}^{\tilde\Psi}$ *then* $\begin{pmatrix} e^{2i\phi} & 0 \\ 0 & e^{-2i\phi} \end{pmatrix}$ *represents the group, double covering the group* $\begin{pmatrix} e^{4i\phi} & 0 \\ 0 & e^{-4i\phi} \end{pmatrix} \cong U(1)$*. At the same time,* $\tilde{\tilde{S}} = \overline{\tilde{S}}$ *(*complex conjugate*). This proved that* $SO(2) \cong U(1)$*.*

Example 6.2. *2n=2, N=1. If* $g_{\tilde{\Lambda}\tilde{\Psi}} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$*, then*

$$\begin{aligned} \gamma_1 &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma_2 = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \\ \eta_1 &= \tfrac{1}{\sqrt{2}}, \quad \eta_2 = -\tfrac{1}{\sqrt{2}}, \quad \sigma_1 = \tfrac{1}{\sqrt{2}}, \quad \sigma_2 = \tfrac{1}{\sqrt{2}}. \\ tr(\eta_{\tilde{\Lambda}}\sigma_{\tilde{\Psi}}) &= \tfrac{1}{2} \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix}, \quad \text{- the tensor is degenerated.} \end{aligned} \tag{6.54}$$

Thus, the identity $tr(\eta_{\tilde{\Lambda}}\sigma_{\tilde{\Psi}}) = \frac{1}{2}g_{ij}$ *is false. The explicit record of the conformity* $S_{\tilde{\Lambda}}{}^{\tilde{\Psi}} x_{\tilde{\Psi}} \longleftrightarrow \tilde{S} X \tilde{\tilde{S}}$ *will have the form*

$$\begin{pmatrix} \text{ch } 4\phi & \text{sh } 4\phi \\ \text{sh } 4\phi & \text{ch } 4\phi \end{pmatrix} \begin{pmatrix} T \\ X \end{pmatrix} \longleftrightarrow \begin{pmatrix} e^{2\phi} & 0 \\ 0 & e^{-2\phi} \end{pmatrix} \begin{pmatrix} 0 & T+X \\ T-X & 0 \end{pmatrix} \begin{pmatrix} e^{-2\phi} & 0 \\ 0 & e^{2\phi} \end{pmatrix}, \tag{6.55}$$

that determines the conformity $SO^+(1,1) \cong R$ *(*additive*), where* $SO^+(1,1)$ *is the connected identity component with* $S_1{}^1 > 0$*.*

Example 6.3. *2n=4, N=2. If* $g_{ij} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$ *then we can construct*

$$\begin{aligned} \eta_1 &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \eta_2 = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \\ \eta_3 &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \eta_4 = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}, \\ \sigma_1 &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma_2 = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \\ \sigma_3 &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma_4 = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \end{aligned} \tag{6.56}$$

The operators γ_i *constructed in accordance with* $\gamma_i = \begin{pmatrix} 0 & \sigma_i \\ \eta_i & 0 \end{pmatrix}$*. It is easy to verify that* $tr(\eta_i\sigma_j) = tr(\sigma_i\eta_j) = g_{ij}$*. The calculations in this case are more cumbersome, but similar to the case* $n = 2$ *(and in part, will be carried out below). Just note that the fact that here the covering* $SO^+(2,2) \cong SL(2) \times SL(2)/\pm 1$ *acts. However, it would be desirable to know how to pass from the case n=4 to the case n=2.*

Example 6.4. $n = 2 \subset 2n = 4,\ N = 2.$

$$m_{\Lambda}{}^{\tilde\Lambda} = \tfrac{1}{2}\begin{pmatrix} 1 & -1 & -i & i \\ i & i & 1 & 1 \end{pmatrix}, \quad m^{\Lambda}{}_{\tilde\Lambda} = \tfrac{1}{2}\begin{pmatrix} 1 & -1 & i & -i \\ -i & -i & 1 & 1 \end{pmatrix},$$
$$g_{\Lambda\Psi} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad G_{\tilde\Lambda\tilde\Psi} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad f_{\tilde\Lambda}{}^{\tilde\Psi} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix},$$
$$x^{\tilde\Lambda} = (T, Y, Z, X), \quad x^{\Lambda} = m^{\Lambda}{}_{\tilde\Lambda} x^{\tilde\Lambda} = \tfrac{1}{2}(T - Y + i(Z - X), -i(T + Y) + Z + X) \tag{6.57}$$

then each special orthogonal transformation $S_{\tilde\Lambda}{}^{\tilde\Psi} \in SO^{+}(2,2)$ (det $\| S_{\Lambda}{}^{\Psi} \| = 1$, *and the determinant of the upper left minor* 2×2 *is positive) can be represented as a finite product of the elementary transformations. Since* $fSf = -S$ *due to the presence of the complex structure in the tangent bundle then such the transformation will be of the two kinds*

$$(S_1)_{\tilde\Lambda}{}^{\tilde\Psi} = \begin{pmatrix} \cos 2\varphi & \sin 2\varphi & 0 & 0 \\ \sin 2\varphi & -\cos 2\varphi & 0 & 0 \\ 0 & 0 & \cos 2\varphi & \sin 2\varphi \\ 0 & 0 & \sin 2\varphi & -\cos 2\varphi \end{pmatrix},$$
$$(S_2)_{\tilde\Lambda}{}^{\tilde\Psi} = \begin{pmatrix} \mathrm{ch}\ 2\psi & 0 & 0 & -\mathrm{sh}\ 2\psi \\ 0 & -\mathrm{ch}\ 2\psi & -\mathrm{sh}\ 2\psi & 0 \\ 0 & \mathrm{sh}\ 2\psi & \mathrm{ch}\ 2\psi & 0 \\ \mathrm{sh}\ 2\psi & 0 & 0 & -\mathrm{ch}\ 2\psi \end{pmatrix}. \tag{6.58}$$

The complex representation $S_{\tilde\Lambda}{}^{\tilde\Psi}$ *can be defined as* $S_{\Lambda}{}^{\Psi} = m_{\Lambda}{}^{\tilde\Lambda} S_{\tilde\Lambda}{}^{\tilde\Psi} m^{\Psi}{}_{\tilde\Psi}$

$$(S_1)_{\Lambda}{}^{\Psi} = \begin{pmatrix} -\sin 2\varphi & -i\cos 2\varphi \\ i\cos 2\varphi & \sin 2\varphi \end{pmatrix}, \quad (S_2)_{\Lambda}{}^{\Psi} = \begin{pmatrix} i\,\mathrm{sh}\ 2\psi & -i\,\mathrm{ch}\ 2\psi \\ i\,\mathrm{ch}\ 2\psi & -i\,\mathrm{sh}\ 2\psi \end{pmatrix}. \tag{6.59}$$

Therefore, $S = S_2 S_1$ *describes the group* $SO(2, \mathbb{C})$ *completely. Now let*

$$\begin{array}{lcl} (r_1)^{\tilde\Lambda} = \sqrt{2}(\cos\phi, \sin\phi, 0, 0) & , & (r_2)^{\tilde\Lambda} = \sqrt{2}(0, 0, \cos\phi, \sin\phi), \\ (r_1)_{\tilde\Lambda} = \sqrt{2}(\cos\phi, \sin\phi, 0, 0) & , & (r_2)_{\tilde\Lambda} = \sqrt{2}(0, 0, -\cos\phi, -\sin\phi). \end{array} \tag{6.60}$$

Construct $(S_1)_{\tilde\Lambda}{}^{\tilde\Theta} = ((r_1)_{\tilde\Lambda}(r_1)^{\tilde\Psi} - \delta_{\tilde\Lambda}{}^{\tilde\Psi})((r_2)_{\tilde\Psi}(r_2)^{\tilde\Theta} + \delta_{\tilde\Psi}{}^{\tilde\Theta})$. *At the same time,* $(r_1)_{\tilde\Lambda}(r_1)^{\tilde\Lambda} = -(r_2)_{\tilde\Lambda}(r_2)^{\tilde\Lambda} = 2$. *Let* $\eta_{\tilde\Lambda}$ *and* $\sigma_{\tilde\Lambda}$ *be the same as in the previous example*

$$\begin{array}{c} R_1 = (r_1)^{\tilde\Lambda}\eta_{\tilde\Lambda} = \begin{pmatrix} \cos\phi & \sin\phi \\ -\sin\phi & \cos\phi \end{pmatrix} \quad , \quad R_2 = (r_2)^{\tilde\Lambda}\sigma_{\tilde\Lambda} = \begin{pmatrix} -\cos\phi & \sin\phi \\ \sin\phi & \cos\phi \end{pmatrix}, \\ R_1 R_2 = \begin{pmatrix} -\cos 2\phi & \sin 2\phi \\ \sin 2\phi & \cos 2\phi \end{pmatrix} \quad , \quad R_2 R_1 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \\ (r_3)^{\tilde\Lambda} = \sqrt{2}(\mathrm{ch}\ \psi, 0, 0, -\mathrm{sh}\ \psi) \quad , \quad (r_4)^{\tilde\Lambda} = \sqrt{2}(0, -\mathrm{sh}\ \psi, -\mathrm{ch}\ \psi, 0), \\ (r_3)_{\tilde\Lambda} = \sqrt{2}(\mathrm{ch}\ \psi, 0, 0, \mathrm{sh}\ \psi) \quad , \quad (r_4)_{\tilde\Lambda} = \sqrt{2}(0, -\mathrm{sh}\ \psi, \mathrm{ch}\ \psi, 0). \end{array} \tag{6.61}$$

Construct $(S_2)_{\tilde\Lambda}{}^{\tilde\Theta} = ((r_3)_{\tilde\Lambda}(r_3)^{\tilde\Psi} - \delta_{\tilde\Lambda}{}^{\tilde\Psi})((r_4)_{\tilde\Psi}(r_4)^{\tilde\Theta} + \delta_{\tilde\Psi}{}^{\tilde\Theta})$. *At the same time,*

$(r_3)_{\tilde{\Lambda}}(r_3)^{\tilde{\Lambda}} = -(r_4)_{\tilde{\Lambda}}(r_4)^{\tilde{\Lambda}} = 2.$

$$
\begin{gathered}
R_3 = (r_3)^{\tilde{\Lambda}}\eta_{\tilde{\Lambda}} = \begin{pmatrix} \mathrm{ch}\ \psi & \mathrm{sh}\ \psi \\ \mathrm{sh}\ \psi & \mathrm{ch}\ \psi \end{pmatrix} \quad , \quad R_4 = (r_4)^{\tilde{\Lambda}}\sigma_{\tilde{\Lambda}} = \begin{pmatrix} \mathrm{ch}\ \psi & \mathrm{sh}\ \psi \\ -\mathrm{sh}\ \psi & -\mathrm{ch}\ \psi \end{pmatrix}, \\
R_3R_4 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad , \qquad R_4R_3 = \begin{pmatrix} \mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ -\mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix}, \\
\tilde{S}^T = R_1R_2R_3R_4 = \begin{pmatrix} -\cos 2\phi & -\sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix}, \\
\tilde{\tilde{S}} = R_4R_3R_2R_1 = \begin{pmatrix} -\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ \mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix}.
\end{gathered}
\tag{6.62}
$$

Since $x^{\tilde{\Lambda}} = (T, Y, Z, X)$, $x^{\Lambda} = \frac{1}{2}(T - Y + i(Z - X), -i(T + Y) + Z + X)$ *then*

$$
\begin{gathered}
x^{\Lambda}S_{\Lambda}{}^{\Psi} \longleftrightarrow \tilde{S}^T X_{\eta}\tilde{\tilde{S}} = \\
= \begin{pmatrix} -\cos 2\phi & -\sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix} \begin{pmatrix} T + Z & -X + Y \\ -X - Y & T - Z \end{pmatrix} \begin{pmatrix} -\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ \mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix}.
\end{gathered}
\tag{6.63}
$$

Thus, $SO(2, \mathbb{C}) \cong U(1) \times R$*(additive), where* $\begin{pmatrix} -\cos 2\phi & -\sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix}$ *form the group* $U(1)$ *and* $\begin{pmatrix} -\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ \mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix}$ *form the additive group* R*. If to make the substitution* η *on* σ *in the relevant equations then*

$$
\begin{gathered}
x^{\Lambda}S_{\Lambda}{}^{\Psi} \longleftrightarrow \tilde{\tilde{S}}'^T X_{\sigma}\tilde{S}' = \\
= \begin{pmatrix} -\mathrm{ch}\ 2\psi & -\mathrm{sh}\ 2\psi \\ -\mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix} \begin{pmatrix} T - Z & X - Y \\ X + Y & T + Z \end{pmatrix} \begin{pmatrix} -\cos 2\phi & \sin 2\phi \\ -\sin 2\phi & \cos 2\phi \end{pmatrix}.
\end{gathered}
\tag{6.64}
$$

Example 6.5. $n = 2 \subset 2n = 4$, $N = 2$. *Let the involution tensor in the tangent fiber for the real representation* V_4 *be the tensor (the left inclusion of the metric of index equal to 2, the right inclusion of the metric of index equal to 1)*

$$
\begin{gathered}
S_{\tilde{\Lambda}}{}^{\tilde{\Psi}} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad S_{\tilde{\Lambda}}{}^{\tilde{\Psi}} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \\
G_{\tilde{\Lambda}\tilde{\Psi}} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix},
\end{gathered}
\tag{6.65}
$$

and all the identities of the previous examples are executed. Then for the complex representation $\mathbb{C}V_2$*, it will have the form* $S_{\Lambda}{}^{\Psi'} = m_{\Lambda}{}^{\tilde{\Lambda}}S_{\tilde{\Lambda}}{}^{\tilde{\Psi}}\bar{m}^{\Lambda'}{}_{\tilde{\Psi}}$

$$
S_{\Lambda}{}^{\Psi'} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad S_{\Lambda}{}^{\Psi'} = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \tag{6.66}
$$

On the other hand, there is an image of the involution (as letters T denotes the transposition), and in some basis $S_{\tilde{\Lambda}}{}^{\tilde{\Psi}}\eta_{\tilde{\Psi}} = \tilde{S}^T\eta_{\tilde{\Psi}}\tilde{S}$

$$\tilde{S} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \qquad \tilde{S} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \tag{6.67}$$

In the first case (index equal to 2)

$$\begin{gathered}
\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}\begin{pmatrix} -\cos 2\phi & -\sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix}\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} -\cos 2\phi & -\sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix}, \\
\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}\begin{pmatrix} -\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ \mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix}\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} -\mathrm{ch}\ 2\psi & -\mathrm{sh}\ 2\psi \\ -\mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix}, \\
H_i{}^{\Lambda} = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 1 & -i \\ -1 & -i \end{pmatrix}, \quad H^i{}_{\Lambda} = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 1 & i \\ -1 & i \end{pmatrix}, \quad g_{ij} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad g_{\Lambda\Psi} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.
\end{gathered} \tag{6.68}$$

Therefor, $\psi = 0$

$$(S_1)_i{}^j = \begin{pmatrix} \cos 2\phi & \sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix}, \quad (S_2)_i{}^j = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \tag{6.69}$$

and we again come to the isomorphism $SO(2) \cong U(1)$

$$\begin{gathered}
\begin{pmatrix} T, & Y \end{pmatrix}\begin{pmatrix} \cos 2\phi & \sin 2\phi \\ -\sin 2\phi & \cos 2\phi \end{pmatrix} \longleftrightarrow \\
\longleftrightarrow \left(\cos 2\phi \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \sin 2\phi \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}\right)\begin{pmatrix} T & Y \\ -Y & T \end{pmatrix}.
\end{gathered} \tag{6.70}$$

In the second case (index 1),

$$\begin{gathered}
\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}\begin{pmatrix} -\cos 2\phi & -\sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix}\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} -\cos 2\phi & \sin 2\phi \\ -\sin 2\phi & -\cos 2\phi \end{pmatrix}, \\
\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}\begin{pmatrix} -\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ \mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix}\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} -\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ \mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix}, \\
H_i{}^{\Lambda} = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 1 & i \\ i & 1 \end{pmatrix}, \quad H^i{}_{\Lambda} = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 1 & -i \\ -i & 1 \end{pmatrix}, \\
g_{ij} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad g_{\Lambda\Psi} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.
\end{gathered} \tag{6.71}$$

Therefor, $2\phi = 0, \pi$

$$(S_1)_i{}^j = \pm\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (S_2)_i{}^j = \begin{pmatrix} -\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ -\mathrm{sh}\ 2\psi & \mathrm{ch}\ 2\psi \end{pmatrix}, \tag{6.72}$$

and we again come to the isomorphism $SO^+(1,1) \cong R(additive)$

$$\begin{gathered}
\begin{pmatrix} T, & X \end{pmatrix}\begin{pmatrix} \mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ \mathrm{sh}\ 2\phi & \mathrm{ch}\ 2\phi \end{pmatrix} \longleftrightarrow \\
\longleftrightarrow \begin{pmatrix} T & -X \\ -X & T \end{pmatrix}\left(\mathrm{ch}\ 2\psi \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \mathrm{sh}\ 2\psi \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}\right).
\end{gathered} \tag{6.73}$$

The following example is worthy of a separate section.

7 Quaternions

In this section, it is told about how to contact the quaternion algebra with the Clifford algebra and how to construct explicitly the double covering corresponding to this case. The conclusion of all results of this section is made on the basis [18].

It is known that the Clifford algebra $CL(g^n)$ can be represented as

$$\begin{aligned} CL(g^n) &: AI + B^{\Lambda}e_{\Lambda} + C^{\Lambda\Psi}e_{\Lambda}e_{\Psi} + D^{\Lambda\Psi\Phi}e_{\Lambda}e_{\Psi}e_{\Phi} + ... = \\ &= \begin{cases} CL^0(g^n)(\text{even}) : AI + C^{\Lambda\Psi}e_{\Lambda}e_{\Psi} + ..., \\ CL^1(g^n)(\text{odd}) : B^{\Lambda}e_{\Lambda} + D^{\Lambda\Psi\Phi}e_{\Lambda}e_{\Psi}e_{\Phi} + ..., \end{cases} \end{aligned} \tag{7.1}$$

that is called Z_2 *grading*. In this case, $CL^0(g^n)$ is a subalgebra of $CL(g^n)$. If we consider n=4 with the 8-dimensional Clifford algebra $CL^0(g^n)$ then the basis of this algebra has the form

$$1, \quad e_1e_2, \quad e_1e_3, \quad e_2e_3, \quad e_1e_4, \quad e_2e_4, \quad e_3e_4, \quad e_1e_2e_3e_4. \tag{7.2}$$

From the Clifford equation (2.8), the relation

$$e_{\Lambda}e_{\Psi} = 1, \quad \Lambda = \Psi; \quad e_{\Lambda}e_{\Psi} = -e_{\Psi}e_{\Lambda}, \quad \Lambda \neq \Psi \tag{7.3}$$

will follow in some basis

$$\begin{array}{ll} I = 1, & i := e_1e_2, \\ j =: e_1e_3, & k := e_2e_3, \\ e := e_3e_4, & f := ie = e_1e_2e_3e_4, \\ g := je = e_1e_3e_3e_4 = e_1e_4, & h := ke = e_2e_3e_3e_4 = e_2e_4, \end{array} \tag{7.4}$$

Consider the inclusion $\mathbb{R}^4 \subset \mathbb{C}^4$. Then we can find the representation of this basis with the operators $\gamma_i = H_i{}^{\Lambda}\gamma_{\Lambda}$ constructed above. But the basis, in turn, is the direct sum of two quaternionic bases; for this, the isomorphism $CL^0(g^n_{(n,0)}) \cong \mathbb{H} \oplus \mathbb{H}$ is responsible. In order to construct an appropriate representation of the specified basis, it is necessary to demand the performance of the condition $tr(\eta_i\sigma_j) = \frac{N}{2}g_{ij}$ for the parts of the operators γ_i. And then, indeed, it may be restricted by the quaternion basis (just having 2 generators i and j (k = ij) and the quaternion identity I)

$$\begin{aligned} &\tfrac{1}{\sqrt{2}}I := \eta_1 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \tfrac{1}{\sqrt{2}}i := \eta_2 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \\ &\tfrac{1}{\sqrt{2}}j := \eta_3 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \tfrac{1}{\sqrt{2}}k := \eta_4 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \\ &\sigma_1 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma_2 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix}, \\ &\sigma_3 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \sigma_4 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}. \end{aligned} \tag{7.5}$$

Now suppose that in the space $\mathbb{R}^4 \subset \mathbb{C}^4$, the vectors

$$\begin{aligned} &r_1 = (\cos\phi_1, \sin\phi_1, 0, 0), \quad r_2 = (\cos\phi_2, 0, \sin\phi_2, 0), \quad r_3 = (0, \cos\phi_3, \sin\phi_3, 0), \\ &r_4 = (\cos\phi_4, 0, 0, \sin\phi_4), \quad r_5 = (0, \cos\phi_5, 0, \sin\phi_5), \quad r_6 = (0, 0, \cos\phi_6, \sin\phi_6), \end{aligned} \tag{7.6}$$

generating the elementary orthogonal transformations, are defined. Here, as before, the transformations $(r_I)^i(r_I)_j - \delta_j^{\ i}$ $(I = \overline{1,6})$ generate all rotations of the group O(4). Accordingly, on the space $\mathbb{C}^2$ $(N = 2^{2/2-1} = 2)$, their images

$$\begin{gathered}
R_1 := r_1^{\ i}\eta_i = \begin{pmatrix} \cos\phi_1 & i\sin\phi_1 \\ i\sin\phi_1 & \cos\phi_1 \end{pmatrix}, \quad R_2 := r_2^{\ i}\sigma_i = \begin{pmatrix} \cos\phi_2 & \sin\phi_2 \\ -\sin\phi_2 & \cos\phi_2 \end{pmatrix}, \\
R_3 := r_3^{\ i}\eta_i = \begin{pmatrix} 0 & i\cos\phi_3 - \sin\phi_3 \\ i\cos\phi_3 + \sin\phi_3 & 0 \end{pmatrix}, \\
R_4 := r_4^{\ i}\sigma_i = \begin{pmatrix} \cos\phi_4 - i\sin\phi_4 & 0 \\ 0 & \cos\phi_4 + i\sin\phi_4 \end{pmatrix}, \\
R_5 := r_5^{\ i}\eta_i = \begin{pmatrix} i\sin\phi_5 & i\cos\phi_5 \\ i\cos\phi_5 & -i\sin\phi_5 \end{pmatrix}, \quad R_6 := r_6^{\ i}\sigma_i = \begin{pmatrix} -i\sin\phi_6 & \cos\phi_6 \\ -\cos\phi_6 & i\sin\phi_6 \end{pmatrix}
\end{gathered} \tag{7.7}$$

will be induced. To simplify calculations, we will be restricted to the case when $\phi_1 = \phi_2 = \phi_4 = \frac{\pi}{2}$. Then

$$\begin{gathered}
R_1R_2R_3R_4R_5R_6 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}\begin{pmatrix} 0 & ie^{i\phi_3} \\ ie^{-i\phi_3} & 0 \end{pmatrix}\begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} \\
\begin{pmatrix} i\sin\phi_5 & i\cos\phi_5 \\ i\cos\phi_5 & -i\sin\phi_5 \end{pmatrix}\begin{pmatrix} -i\sin\phi_6 & \cos\phi_6 \\ -\cos\phi_6 & i\sin\phi_6 \end{pmatrix} = \begin{pmatrix} 0 & ie^{i\phi_3} \\ ie^{-i\phi_3} & 0 \end{pmatrix} \\
\begin{pmatrix} \overbrace{\sin\phi_5\sin\phi_6 - i\cos\phi_5\cos\phi_6}^{=:\bar{a}} & \overbrace{i\sin\phi_5\cos\phi_6 - \cos\phi_5\sin\phi_6}^{=:-\bar{b}} \\ \underbrace{i\sin\phi_5\cos\phi_6 + \cos\phi_5\sin\phi_6}_{=:b} & \underbrace{\sin\phi_5\sin\phi_6 + i\cos\phi_5\cos\phi_6}_{=:a} \end{pmatrix} = \\
= \begin{pmatrix} ie^{i\phi_3}b & ie^{i\phi_3}a \\ ie^{-i\phi_3}\bar{a} & -ie^{-i\phi_3}\bar{b} \end{pmatrix} =: A_1.
\end{gathered} \tag{7.8}$$

$$\begin{gathered}
R_6R_5R_4R_3R_2R_1 = \begin{pmatrix} -i\sin\phi_6 & \cos\phi_6 \\ -\cos\phi_6 & i\sin\phi_6 \end{pmatrix}\begin{pmatrix} i\sin\phi_5 & i\cos\phi_5 \\ i\cos\phi_5 & -i\sin\phi_5 \end{pmatrix} \\
\begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}\begin{pmatrix} 0 & ie^{i\phi_3} \\ ie^{-i\phi_3} & 0 \end{pmatrix}\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}\begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} = \\
= \begin{pmatrix} \overbrace{\sin\phi_5\sin\phi_6 + i\cos\phi_5\cos\phi_6}^{=:a} & \overbrace{-i\sin\phi_5\cos\phi_6 + \cos\phi_5\sin\phi_6}^{=:\bar{b}} \\ \underbrace{-i\sin\phi_5\cos\phi_6 - \cos\phi_5\sin\phi_6}_{=:-b} & \underbrace{\sin\phi_5\sin\phi_6 - i\cos\phi_5\cos\phi_6}_{=:\bar{a}} \end{pmatrix} \\
\begin{pmatrix} 0 & -ie^{i\phi_3} \\ -ie^{-i\phi_3} & 0 \end{pmatrix} = \begin{pmatrix} -ie^{-i\phi_3}\bar{b} & -ie^{i\phi_3}a \\ -ie^{-i\phi_3}\bar{a} & ie^{-i\phi_3}b \end{pmatrix} =: \bar{A_1}^T = A_1^{\ *}.
\end{gathered} \tag{7.9}$$

By this, the double covering $SO(3) \cong SU(2)/\{\pm 1\}$ is constructed, and therefore $Spin(3) \cong SU(2)$

$$S_{1i}^{\ \ j}\eta_j = A_1^{\ T}\eta_i A_1^*. \tag{7.10}$$

If we now set $\phi_3 = \phi_5 = \phi_6 = \frac{\pi}{2}$ and carry out similar calculations, we obtain

$$
\begin{gathered}
R_1R_2R_3R_4R_5R_6 = \\
\begin{pmatrix} \cos\phi_1 & i\sin\phi_1 \\ i\sin\phi_1 & \cos\phi_1 \end{pmatrix}
\begin{pmatrix} \cos\phi_2 & \sin\phi_2 \\ -\sin\phi_2 & \cos\phi_2 \end{pmatrix} \\
\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}
\begin{pmatrix} e^{-i\phi_4} & \\ 0 & e^{i\phi_4} \end{pmatrix}
\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}
\begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} = \\
= \left(\overbrace{\underbrace{\begin{matrix} \cos\phi_1\cos\phi_2 - i\sin\phi_1\sin\phi_2 \\ i\sin\phi_1\cos\phi_2 - \cos\phi_1\sin\phi_2 \end{matrix}}_{:=-\bar{b}}}^{:=\bar{a}}
\quad
\overbrace{\underbrace{\begin{matrix} \cos\phi_1\sin\phi_2 + i\sin\phi_1\cos\phi_2 \\ i\sin\phi_1\sin\phi_1 + \cos\phi_1\cos\phi_2 \end{matrix}}_{=:a}}^{=:b} \right) \\
\begin{pmatrix} 0 & -e^{i\phi_4} \\ e^{-i\phi_4} & 0 \end{pmatrix} = \begin{pmatrix} e^{-i\phi_4}b & -e^{i\phi_4}\bar{a} \\ e^{-i\phi_4}a & e^{i\phi_4}\bar{b} \end{pmatrix} =: {A_2}^T.
\end{gathered}
\tag{7.11}
$$

$$
\begin{gathered}
R_6R_5R_4R_3R_2R_1 = \\
\begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}
\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}
\begin{pmatrix} e^{-i\phi_4} & 0 \\ 0 & e^{i\phi_4} \end{pmatrix}
\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \\
\begin{pmatrix} \cos\phi_2 & \sin\phi_2 \\ -\sin\phi_2 & \cos\phi_2 \end{pmatrix}
\begin{pmatrix} \cos\phi_1 & i\sin\phi_1 \\ i\sin\phi_1 & \cos\phi_1 \end{pmatrix} =
\begin{pmatrix} 0 & -e^{-i\phi_4} \\ e^{i\phi_4} & 0 \end{pmatrix} \\
\left(\overbrace{\underbrace{\begin{matrix} \cos\phi_1\cos\phi_2 + i\sin\phi_1\sin\phi_2 \\ i\sin\phi_1\cos\phi_2 - \cos\phi_1\sin\phi_2 \end{matrix}}_{:=-\bar{b}}}^{:=a}
\quad
\overbrace{\underbrace{\begin{matrix} \cos\phi_1\sin\phi_2 + i\sin\phi_1\cos\phi_2 \\ -i\sin\phi_1\sin\phi_1 + \cos\phi_1\cos\phi_2 \end{matrix}}_{=:\bar{a}}}^{:=b} \right) = \\
= \begin{pmatrix} e^{-i\phi_4}\bar{b} & -e^{-i\phi_4}\bar{a} \\ e^{i\phi_4}a & e^{i\phi_4}b \end{pmatrix} =: {A_2}^\sharp.
\end{gathered}
\tag{7.12}
$$

Therefor,

$$
{S_2}_i{}^j\eta_j = {A_2}^T\eta_i A_2^\sharp. \tag{7.13}
$$

The transformations A_1, A_2 are the representatives of $SU(2)$ and fully describe the group. Gathering the two equations together, we find that

$$
S_i{}^j\eta_j := (S_2S_1)_i{}^j\eta_j = {A_1}^T{A_2}^T\eta_j A_2^\sharp A_1^* := \tilde{S}^T\eta_j\tilde{\tilde{S}}. \tag{7.14}
$$

Generally speaking, knowing the transformation ${A_1}^T{A_2}^T$, we can say nothing about $A_2^\sharp A_1^*$. Therefore, the transformations $\tilde{S}^T$ and $\tilde{\tilde{S}}$ are distinct. This means that there is the double covering $SO(4) \cong SU(2) \times SU(2)/\{\pm 1\}$. And so $Spin(4) \cong SU(2) \times SU(2)$.

8 Particular solutions of the Clifford equation

In this section it is told about how to construct particular solutions of the Clifford equation with the help of which can uniquely construct the double covering $Spin(n, \mathbb{C})/\{\pm\ 1\} \cong SO(n, \mathbb{C})$. Such the solutions will give the chance uniquely to prolong the Riemannian torsion-free connection into the spinor bundle. The conclusion of all results of this section is made on the basis of previous calculations.

In order to find some particular solutions of the Clifford equation (6.36), we rewrite it as

$$
{\eta_\Lambda}^{AB}\eta_{\Psi CB} + {\eta_\Psi}^{AB}\eta_{\Lambda CB} = g_{\Lambda\Psi}{\delta_C}^A. \tag{8.1}
$$

Example 8.1. *n=4, N=2.*
In this case, we use the isomorphism $\mathbb{C}^4 \cong \mathbb{C}(2)$ *which is constructed by means of the connecting operators* ${\eta^\Lambda}_{AB}$ *(6.37)*

$$r^\Lambda = \frac{1}{2}{\eta^\Lambda}_{AB}R^{AB}. \tag{8.2}$$

Let the metric in $\mathbb{C}^4$ *have the form*

$$g_{\Lambda\Psi} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \tag{8.3}$$

It is obvious that any tensor R^{AB} *can be decomposed into the three symmetrical and one antisymmetrical parts*

$$\begin{gathered} R^{AB} = R_1 \underbrace{\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}}_{(\varepsilon_1)^{AB}} + R_2 \underbrace{\begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix}}_{(\varepsilon_2)^{AB}} + R_3 \underbrace{\begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}}_{(\varepsilon_3)^{AB}} + R_{\tilde 1} \underbrace{\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}}_{(\varepsilon_{\tilde 1})^{AB}}, \\ R_{AB} = R_1 \underbrace{\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}}_{(\varepsilon_1)_{AB}} + R_2 \underbrace{\begin{pmatrix} -i & 0 \\ 0 & -i \end{pmatrix}}_{(\varepsilon_2)_{AB}} + R_3 \underbrace{\begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}}_{(\varepsilon_3)_{AB}} + R_{\tilde 1} \underbrace{\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}}_{(\varepsilon_{\tilde 1})_{AB}} \end{gathered} \tag{8.4}$$

with the metric spin-tensor $\varepsilon_{AB} = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$.

Example 8.2. *n=6, N=4.*
Let's construct now ($A, B, ... = \overline{1, N}$, $a, b, ... = \overline{1, N/2}$, $I = \overline{1, 2, 3, \tilde 1}$)

$$\begin{gathered} (\varepsilon_I)^{AB} = \begin{pmatrix} (\varepsilon_I)^{ab} & 0 \\ 0 & -(\varepsilon_I)^T_{ab} \end{pmatrix}, \\ (\varepsilon_4)^{AB} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad (\varepsilon_{\tilde 2})_{AB} = \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \\ -i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{pmatrix}, \end{gathered} \tag{8.5}$$

where $(\varepsilon_I)^{ab}$ *are taken from the previous example (n=4, N=2). It is one of possible variants (the 4 symmetrical and 2 antisymmetric parts). If all received* $(\varepsilon_I)^{AB}$ ($I = \overline{1, 2, 3, 4, \tilde 1, \tilde 2}$) *multiply by*

$$\begin{pmatrix} 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \end{pmatrix} \tag{8.6}$$

on the right side then one can construct the new tensors (the 6 antisymmetric parts)

$$(\varepsilon_{\tilde{1}})^{AB} = \begin{pmatrix} 0 & 0 & 0 & i \\ 0 & 0 & i & 0 \\ 0 & -i & 0 & 0 \\ -i & 0 & 0 & 0 \end{pmatrix}, \quad (\varepsilon_{\tilde{2}})^{AB} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix},$$
$$(\varepsilon_{\tilde{3}})^{AB} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}, \quad (\varepsilon_{\tilde{4}})^{AB} = \begin{pmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & i \\ i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{pmatrix}, \tag{8.7}$$
$$(\varepsilon_{\tilde{5}})^{AB} = \begin{pmatrix} 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \end{pmatrix}, \quad (\varepsilon_{\tilde{6}})^{AB} = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix},$$

which will enable to define

$$R^{AB} = -R^{BA} = R_{\tilde{1}}(\varepsilon_{\tilde{1}})^{AB} + R_{\tilde{2}}(\varepsilon_{\tilde{2}})^{AB} + R_{\tilde{3}}(\varepsilon_{\tilde{3}})^{AB} + R_{\tilde{4}}(\varepsilon_{\tilde{4}}^{AB}) + R_{\tilde{5}}(\varepsilon_{\tilde{5}})^{AB} + R_{\tilde{6}}(\varepsilon_{\tilde{6}}^{AB}). \tag{8.8}$$

In order to obtain a similar construction for any even n, we need in the following definition.

Definition 8.1. *Let in the space $\mathbb{C}^n$ $(n \geq 4)$, n orthogonal vectors $(\eta_I)^\Lambda$*

$$(\eta_I)^\Lambda (\eta_J)_\Lambda = 2\delta_{IJ}, \quad \Lambda,\ \Psi,\ ... = \overline{1, n} \tag{8.9}$$

be given. Then for the spinor space, the vectors are uniquely associated to the n tensors

$$(\varepsilon_I)_{AB} := (\eta_I)^\Lambda \eta_{\Lambda AB}. \tag{8.10}$$

Therefore, among the n vectors $(\eta_I)^\Lambda$, some $\frac{n}{2} + q$ vectors

$$\eta^\Psi{}_{(AB)} = \frac{1}{2} \sum_{Q=1}^{\frac{n}{2}+q} (\eta_Q)^\Psi (\varepsilon_Q)_{AB}, \quad (\varepsilon_Q)_{AB} = (\varepsilon_Q)_{BA} \tag{8.11}$$

and some $\frac{n}{2} + \tilde{q}$ vectors

$$\eta^\Psi{}_{[AB]} = \frac{1}{2} \sum_{\tilde{Q}=1}^{\frac{n}{2}+\tilde{q}} (\eta_{\tilde{Q}})^\Psi (\varepsilon_{\tilde{Q}})_{AB}, \quad (\varepsilon_{\tilde{Q}})_{AB} = -(\varepsilon_{\tilde{Q}})_{BA}, \quad q + \tilde{q} = 0 \tag{8.12}$$

must be found. In addition, we require that the real representation of these operators satisfy similar relations.

Corollary 8.1. *From the relation*

$$\tilde{T}_A{}^B (\varepsilon_I)_{BC} + \tilde{\tilde{T}}_C{}^B (\varepsilon_I)_{AB} = 0, \tag{8.13}$$

performed for all $I = \overline{1, n}$, the parities $\tilde{T}_A{}^B = \tilde{\tilde{T}}_A{}^B = 0$ follow.

Proof. We use the identity

$$\begin{gathered}\eta_{[\Gamma}{}^{CN}\eta_{\Omega]MN}\varepsilon_{CD}{}^{ML}=\tfrac{1}{2}\sum_{I}\eta_{[\Gamma}{}^{CN}\eta_{\Omega]MN}(\varepsilon_I)_{CD}(\varepsilon_I)^{ML}=\\ =\tfrac{1}{2}\sum_{I}(-\eta_{[\Gamma|CD}(\varepsilon_{I|})^{CN}+(\eta_I)_{[\Gamma}\delta_{|D|}{}^{N})\eta_{\Omega]MN}(\varepsilon_I)^{ML}=\\ =\tfrac{1}{2}\sum_{I}(-\eta_{[\Gamma|CD|})(-\eta_{\Omega]}{}^{CL}+(\eta_{|I|})_{\Omega]}(\varepsilon_I)^{CL})+(\eta_I)_{[\Gamma}\eta_{\Omega]MD}(\varepsilon_I)^{ML}=\tfrac{n-4}{2}\eta_{[\Gamma|CD|}\eta_{\Omega]}{}^{CL}.\end{gathered}\tag{8.14}$$

Then

$$\tilde{T}_A{}^B(\varepsilon_I)_{BC}+\tilde{\tilde{T}}_C{}^B(\varepsilon_I)_{AB}=0,$$

$$\left\{\begin{array}{l}\frac{n}{2}\tilde{T}_A{}^D+\tilde{\tilde{T}}_C{}^B\varepsilon_{AB}{}^{DC}=0,\\ \frac{n}{2}\tilde{\tilde{T}}_A{}^D+\tilde{T}_C{}^B\varepsilon_{AB}{}^{DC}=0,\end{array}\right.\qquad\Rightarrow\tilde{\tilde{T}}_A{}^A=\tilde{T}_A{}^A=0,$$

$$\left\{\begin{array}{l}\frac{n}{2}\tilde{T}_A{}^D\eta_{\tilde{\Lambda}}{}^{AK}\eta_{\tilde{\Psi}DK}-\frac{n-4}{2}\tilde{\tilde{T}}_A{}^D\eta_{\tilde{\Lambda}}{}^{KA}\eta_{\tilde{\Psi}KD}=0,\\ \frac{n}{2}\tilde{\tilde{T}}_A{}^D\eta_{\tilde{\Lambda}}{}^{AK}\eta_{\tilde{\Psi}DK}-\frac{n-4}{2}\tilde{T}_A{}^D\eta_{\tilde{\Lambda}}{}^{KA}\eta_{\tilde{\Psi}KD}=0,\end{array}\right.$$

$$\left\{\begin{array}{l}\frac{n}{2}\tilde{T}_A{}^D\varepsilon_{PL}{}^{AK}\varepsilon^{QL}{}_{DK}-\frac{n-4}{2}\tilde{\tilde{T}}_A{}^D\varepsilon_{PL}{}^{KA}\varepsilon^{QL}{}_{KD}=0,\\ \frac{n}{2}\tilde{\tilde{T}}_A{}^D\varepsilon_{LP}{}^{AK}\varepsilon^{LQ}{}_{DK}-\frac{n-4}{2}\tilde{T}_A{}^D\varepsilon_{LP}{}^{KA}\varepsilon^{LQ}{}_{KD}=0,\end{array}\right.\quad\Rightarrow\left\{\begin{array}{l}\tilde{T}_A{}^D\varepsilon_{PL}{}^{AK}\varepsilon^{QL}{}_{DK}=0,\\ \tilde{\tilde{T}}_A{}^D\varepsilon_{PL}{}^{AK}\varepsilon^{QL}{}_{DK}=0.\end{array}\right.\tag{8.15}$$

Consequently, $\tilde{\tilde{T}}_A{}^D=\tilde{T}_A{}^D=0$. □

Thus, on the generating tensors $(\varepsilon_I)_{AB}$, the condition $(I\neq J)$

$$\begin{gathered}(\varepsilon_J)_{AB}=-(\varepsilon_I)_{CB}(\varepsilon_J)^{CD}(\varepsilon_I)_{AD},\quad\eta_\Lambda{}^{AB}(\varepsilon_I)_{AD}(\varepsilon_I)_{CB}=-\eta_{\Lambda CD}+(\eta_I)_\Lambda(\varepsilon_I)_{CD},\\ (\eta_I)^\Lambda=\tfrac{2}{N}\eta^{\Lambda AB}(\varepsilon_I)_{AB},\quad(\varepsilon_I)_{AB}=(\eta_I)^\Lambda\eta_{\Lambda AB},\quad(\varepsilon_I)_{AB}(\varepsilon_I)^{AC}=\delta_B{}^C\end{gathered}\tag{8.16}$$

is imposed by the Clifford equation (8.1).

Definition 8.2. *In the bivector space, the 4-valent tensor*

$$\begin{aligned}&\varepsilon_{ABCD}:=\eta_{\Lambda AB}\eta^\Lambda{}_{CD}, &&\bar{\varepsilon}_{A'B'C'D'}:=\bar{\eta}_{\Lambda'A'B'}\bar{\eta}^{\Lambda'}{}_{C'D'},\\ &g_{\Lambda\Psi}=\left(\tfrac{2}{N}\right)^2\eta^{\Lambda AB}\eta^{\Psi CD}\varepsilon_{ABCD}, &&\bar{g}_{\Lambda'\Psi'}=\left(\tfrac{2}{N}\right)^2\bar{\eta}^{\Lambda'A'B'}\bar{\eta}^{\Psi'C'D'}\bar{\varepsilon}_{A'B'C'D'}\end{aligned}\tag{8.17}$$

with the help of which we can raise and lower pair indices is defined.

Note 8.1. *It should be noted that for a real inclusion the condition $(\eta_I)^i(\eta_J)_i=2\delta_{IJ}$ is not always feasible. It should be modified to $(\eta_I)^i(\eta_J)_i=2g_{IJ}$, where $g_{IJ}=0,\ I\neq J$ and $g_{II}=\pm1$, while the sign is chosen in depending on the metric index and the number of the vector. This literally means that the metric in some basis has the diagonal form with ±1 on the main diagonal. Then the equation (8.16) will take the form $(I\neq J)$*

$$\begin{gathered}(\varepsilon_J)_{AB}=-g_{II}(\varepsilon_I)_{CB}(\varepsilon_J)^{CD}(\varepsilon_I)_{AD},\quad g_{II}\eta_i{}^{AB}(\varepsilon_I)_{AD}(\varepsilon_I)_{CB}=-\eta_{iCD}+(\eta_I)_i(\varepsilon_I)_{CD},\\ (\eta_I)^i=\tfrac{2}{N}\eta^{iAB}(\varepsilon_I)_{AB},\quad g_{II}(\varepsilon_I)_{AB}=(\eta_I)^i\eta_{iAB},\quad g_{II}(\varepsilon_I)_{AB}(\varepsilon_I)^{AC}=\delta_B{}^C.\end{gathered}\tag{8.16$'$}$$

In the case of the complex space, we can always restrict ourselves to +1 with the Euclidean metric δ_{IJ}. This will be enough to prove the main identities. In addition, one can pass to the real spaces using the corresponding inclusion operator.

Therefore, it is possible the two significantly different ways depending on multiplicity n

$$I).\quad n\ mod\ 4 = 0,\qquad II).\quad n\ mod\ 4 = 2 \tag{8.18}$$

which determine either the pair of the metric tensors $\varepsilon_{AB} = -\varepsilon_{BA}$, $\tilde{\varepsilon}_{AB} = \tilde{\varepsilon}_{BA}$ in Case I). or the pair of the involutions ${E_A}^B$, $\tilde{E}_A$ in Case II). (the proof is given in Appendix)

$$\begin{gathered}
I).\quad \varepsilon_{A_1 A_{(n/2+q+1)}} := (\varepsilon_{(n/2+q)})_{A_{(n/2+q)}A_{(n/2+q+1)}} \prod_{Q=1}^{(n/2+q-1)/2} (\varepsilon_{2Q-1})_{A_{2Q-1}A_{2Q}} (\varepsilon_{2Q})^{A_{2Q+1}A_{2Q}},\\
\varepsilon_{AB}\varepsilon^{CB} = {\delta_A}^C,\ \left[\begin{array}{ll} \varepsilon_{AB} = -\varepsilon_{BA}, & n/2+q > 2,\\ \varepsilon_{AB} = \varepsilon_{BA}, & n/2+q = 1, \end{array}\right],\ \eta_{\Lambda AB} = {\eta_\Lambda}^{CD}\varepsilon_{CA}\varepsilon_{DB},\\
\\
\tilde{\varepsilon}_{A_1 A_{(n/2+\tilde{q})}} := (\varepsilon_{\widetilde{(n/2+\tilde{q})}})_{A_{(n/2+\tilde{q})}A_{(n/2+\tilde{q}+1)}} \prod_{\tilde{Q}=1}^{(n/2+\tilde{q}-1)/2} (\varepsilon_{2\tilde{Q}-1})_{A_{2\tilde{Q}-1}A_{2\tilde{Q}}} (\varepsilon_{2\tilde{Q}})^{A_{2\tilde{Q}}A_{2\tilde{Q}+1}},\\
\tilde{\varepsilon}_{AB}\tilde{\varepsilon}^{CB} = {\delta_A}^C,\ \left[\begin{array}{ll} \tilde{\varepsilon}_{AB} = \tilde{\varepsilon}_{BA}, & n/2+q > 2,\\ \tilde{\varepsilon}_{AB} = -\tilde{\varepsilon}_{BA}, & n/2+q = 1, \end{array}\right],\ \eta_{\Lambda AB} = {\eta_\Lambda}^{CD}\tilde{\varepsilon}_{CA}\tilde{\varepsilon}_{DB},
\end{gathered} \tag{8.19}$$

$$\begin{gathered}
II).\quad {E_{A_1}}^{A_{(n/2+q+1)}} := \prod_{Q=1}^{(n/2+q)/2} (\varepsilon_{2Q-1})_{A_{2Q-1}A_{2Q}} (\varepsilon_{2Q})^{A_{2Q+1}A_{2Q}},\\
{E_A}^B {E_B}^C = -{\delta_A}^C,\quad {\eta_\Lambda}^{AB} = -{\eta_\Lambda}^{CD}{E_D}^A{E_C}^B,\\
{\tilde{E}_{A_1}}{}^{A_{(n/2+\tilde{q}+1)}} := \prod_{\tilde{Q}=1}^{(n/2+\tilde{q})/2} (\varepsilon_{2\tilde{Q}-1})_{A_{2\tilde{Q}-1}A_{2\tilde{Q}}} (\varepsilon_{2\tilde{Q}})^{A_{2\tilde{Q}+1}A_{2\tilde{Q}}},\\
{\tilde{E}_A}{}^B {\tilde{E}_B}{}^C = -{\delta_A}^C,\quad {\eta_\Lambda}^{AB} = {\eta_\Lambda}^{CD}{\tilde{E}_D}{}^A{\tilde{E}_C}{}^B,
\end{gathered} \tag{8.19'}$$

Note 8.2. *For real inclusion, the tensors from (8.19) multiply by* $g := (\prod_{P=1}^{(n/2+q)} \sqrt{g_{PP}})$, *where either of the two versions* $\pm i$ *for* $g_{\tilde{P}\tilde{P}} = -1$ *can be chosen as the root* $\sqrt{g_{\tilde{P}\tilde{P}}}$.

We will show that such the operators exist. For n = 4, one can construct the representation as described in Example 8.1. Suppose that we have constructed some representation ${\eta_\alpha}^{ab}$ ($\alpha\ ,\beta\ ,... = \overline{1, n-2}; a,\ b,\ ... = \overline{1, 2^{(n-2)/2-1}}$). The operators ${\eta_\Lambda}^{AB}$ ($\Lambda\ ,\Psi\ ,... = \overline{1, n}; A,\ B,\ ... = \overline{1, 2^{n/2-1}}$) are constructed as follows. Let the tensor $g_{\alpha\beta}$ contain only +1 on the main diagonal then

$$\begin{array}{llll}
{\eta_\alpha}^{AB} = \begin{pmatrix} {\eta_\alpha}^{ab} & 0 \\ 0 & -(\eta^T)_{\alpha cd} \end{pmatrix}, & {\eta^\alpha}_{AB} = \begin{pmatrix} {\eta^\alpha}_{pk} & 0 \\ 0 & -(\eta^T)^{\alpha lm} \end{pmatrix},\\
{\eta_{n-1}}^{AB} = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 & i{\delta^a}_d \\ -i{\delta_c}^b & 0 \end{pmatrix}, & {\eta^{n-1}}_{AB} = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 & -i{\delta_p}^m \\ i{\delta^l}_k & 0 \end{pmatrix},\\
{\eta_n}^{AB} = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 & {\delta^a}_d \\ {\delta_c}^b & 0 \end{pmatrix}, & {\eta^n}_{AB} = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 & {\delta_p}^m \\ {\delta^l}_k & 0 \end{pmatrix},
\end{array} \tag{8.20}$$

Under the general scheme of the constructing (8.20) for n = 4, the equation

$$\tilde{T}_A{}^B(\varepsilon_I)_{BC} + \tilde{\tilde{T}}_C{}^B(\varepsilon_I)_{AB} = 0 \tag{8.21}$$

leads to $\tilde{T}_A{}^B = \tilde{\tilde{T}}_A{}^B = 0$ that can be verified directly. Suppose that this equation holds for some even n-2. Then for n, the equation splits into the parts

1. $$\tilde{T}_a{}^b(\varepsilon_I)_{bc} + \tilde{\tilde{T}}_c{}^b(\varepsilon_I)_{ab} = 0, \quad \tilde{T}^a{}_b(\varepsilon_I)^{bc} + \tilde{\tilde{T}}^c{}_b(\varepsilon_I)^{ab} = 0, \quad I = \overline{1, n-2}, \tag{8.22}$$

   which immediately give $\tilde{T}_a{}^b = \tilde{\tilde{T}}_a{}^b = 0, \quad \tilde{T}^a{}_b = \tilde{\tilde{T}}^a{}_b = 0.$

2. $$\begin{aligned} \tilde{T}_{ab}(\varepsilon_{n+1})^b{}_c + \tilde{\tilde{T}}_{cb}(\varepsilon_{n+1})_a{}^b = 0, \quad &\Rightarrow \tilde{T}_{ac} - \tilde{\tilde{T}}_{ca} = 0, \\ \tilde{T}^{ab}(\varepsilon_{n+1})_b{}^c + \tilde{\tilde{T}}^{cb}(\varepsilon_{n+1})^a{}_b = 0, \quad &\Rightarrow -\tilde{T}^{ac} + \tilde{\tilde{T}}^{ca} = 0. \end{aligned} \tag{8.23}$$

3. $$\begin{aligned} \tilde{T}_{ab}(\varepsilon_{n+2})^b{}_c + \tilde{\tilde{T}}_{cb}(\varepsilon_{n+2})_a{}^b = 0, \quad &\Rightarrow \tilde{T}_{ac} + \tilde{\tilde{T}}_{ca} = 0, \\ \tilde{T}^{ab}(\varepsilon_{n+2})_b{}^c + \tilde{\tilde{T}}^{cb}(\varepsilon_{n+2})^a{}_b = 0, \quad &\Rightarrow \tilde{T}^{ac} + \tilde{\tilde{T}}^{ca} = 0, \end{aligned} \tag{8.24}$$

   which immediately give $\tilde{T}_{ab} = \tilde{\tilde{T}}_{ab} = 0, \quad \tilde{T}^{ab} = \tilde{\tilde{T}}^{ab} = 0$. Thus, again $\tilde{T}_A{}^B = \tilde{\tilde{T}}_A{}^B = 0$.

Now suppose that there are the real connecting operators $\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}$. In this case, there is a set of orthogonal vectors such that a part $(\varepsilon_Q)^{\tilde{\tilde{A}}\tilde{\tilde{B}}}$ is symmetric and the remaining part $(\varepsilon_{\tilde{Q}})^{\tilde{\tilde{A}}\tilde{\tilde{B}}}$ is antisymmetric. We will show that one can move towards the complex representation of such the operators which preserve this property and therefore satisfy Definition 8.1.

Algorithm 8.1.
For the first step, the operators

$$\begin{aligned} &\underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} \delta^a{}_{\tilde{\tilde{a}}} & 0 & 0 & i\delta^a{}_{\tilde{\tilde{q}}} \\ 0 & \delta_c{}^{\tilde{\tilde{c}}} & \delta_c{}^{\tilde{\tilde{p}}} & 0 \end{pmatrix}}_{:=(\tilde{\tilde{m}}_{1_+})^A{}_{\tilde{\tilde{A}}}} \cdot \\ &\cdot \underbrace{\begin{pmatrix} \eta_\alpha{}^{\tilde{\tilde{a}}\tilde{\tilde{b}}} & \alpha\delta^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{d}}} & \gamma\delta^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{k}}} & 0 \\ \beta\delta_{\tilde{\tilde{c}}}{}^{\tilde{\tilde{b}}} & -(\eta^T)_{\alpha\,\tilde{\tilde{c}}\tilde{\tilde{d}}} & 0 & \gamma\delta_{\tilde{\tilde{c}}}{}^{\tilde{\tilde{l}}} \\ \delta\delta_{\tilde{\tilde{p}}}{}^{\tilde{\tilde{b}}} & 0 & -(\eta^T)_{\alpha\,\tilde{\tilde{p}}\tilde{\tilde{k}}} & -\alpha\delta_{\tilde{\tilde{p}}}{}^{\tilde{\tilde{l}}} \\ 0 & \delta\delta^{\tilde{\tilde{q}}}{}_{\tilde{\tilde{d}}} & -\beta\delta^{\tilde{\tilde{q}}}{}_{\tilde{\tilde{k}}} & \eta_\alpha{}^{\tilde{\tilde{q}}\tilde{\tilde{l}}} \end{pmatrix}}_{\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}} \underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} \delta_{\tilde{\tilde{b}}}{}^b & 0 \\ 0 & \delta^{\tilde{\tilde{d}}}{}_d \\ 0 & \delta^{\tilde{\tilde{k}}}{}_d \\ -i\delta_{\tilde{\tilde{l}}}{}^b & 0 \end{pmatrix}}_{:=(\tilde{m}^T_{1_+})_{\tilde{\tilde{B}}}{}^B} = \\ &= \underbrace{\begin{pmatrix} \eta_\alpha{}^{ab} & \underbrace{\frac{1}{2}((\alpha - i\beta) + (\gamma + i\delta))}_{:=\phi_1}\delta^a{}_d \\ \underbrace{\frac{i}{2}((\alpha - i\beta) - (\gamma + i\delta))}_{:=\psi_1}\delta_c{}^b & -(\eta^T)_{\alpha cd} \end{pmatrix}}_{:=\eta_{1_+\tilde{\Lambda}}{}^{AB}}. \end{aligned} \tag{8.25}$$

are used as the operators (6.24) and (6.30). As the operators $(\tilde{\tilde{m}}_{1-})^{A}{}_{\tilde{\tilde{A}}}$*, we can take* $\begin{pmatrix} \delta^{a}{}_{\tilde{\tilde{a}}} & 0 & 0 & -i\delta^{a}{}_{\tilde{\tilde{q}}} \\ 0 & \delta_{c}{}^{\tilde{\tilde{c}}} & -\delta_{c}{}^{\tilde{\tilde{p}}} & 0 \end{pmatrix}$*. If we now use the operators* $m_{\Lambda}{}^{\tilde{\Lambda}}$ *translating the subspace* $\mathbb{R}^{4}_{(2,2)}$ *in the subspace* $\mathbb{C}^2$ *and leaving unchanged the subspace* $\mathbb{R}^{2n-4}_{(n-2,n-2)}$ *then we can construct the connecting operators* $\eta_{\Lambda}{}^{ab}$ *of the corresponding space* $\mathbb{C}^2 \oplus \mathbb{R}^{2n-4}_{(n-2,n-2)}$*, while the dimension of the spinor space will be reduced by 2 times.*
2. For the second step, we will use the operators

$$\underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} \delta^{a}{}_{\tilde{\tilde{a}}} & 0 & 0 & i\delta^{a}{}_{\tilde{\tilde{q}}} & 0 & 0 & 0 & 0 \\ 0 & \delta_{c}{}^{\tilde{\tilde{c}}} & \delta_{c}{}^{\tilde{\tilde{p}}} & 0 & 0 & 0 & 0 & \\ 0 & 0 & 0 & 0 & \delta_{a_1}{}^{\tilde{\tilde{a}}_1} & 0 & 0 & i\delta_{a_1}{}^{\tilde{\tilde{p}}_1} \\ 0 & 0 & 0 & 0 & 0 & \delta^{c_1}{}_{\tilde{\tilde{c}}_1} & \delta^{c_1}{}_{\tilde{\tilde{q}}_1} & 0 \end{pmatrix}}_{:=(\tilde{\tilde{m}}_2)^{A}{}_{\tilde{\tilde{A}}}} \cdot$$

$$\underbrace{\begin{pmatrix} \eta_{\alpha}{}^{\tilde{\tilde{a}}\tilde{\tilde{b}}} & \alpha\delta^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{d}}} & \gamma\delta^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{k}}} & 0 & \phi_1\delta^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{b}}_1} & 0 & 0 & 0 \\ \beta\delta_{\tilde{\tilde{c}}}{}^{\tilde{\tilde{b}}} & -(\eta^T)_{\alpha\tilde{\tilde{c}}\tilde{\tilde{d}}} & 0 & \gamma\delta_{\tilde{\tilde{c}}}{}^{\tilde{\tilde{l}}} & 0 & \phi_1\delta_{\tilde{\tilde{c}}}{}^{\tilde{\tilde{d}}_1} & 0 & 0 \\ \delta\delta_{\tilde{\tilde{p}}}{}^{\tilde{\tilde{b}}} & 0 & -(\eta^T)_{\alpha\tilde{\tilde{p}}\tilde{\tilde{k}}} & -\alpha\delta_{\tilde{\tilde{p}}}{}^{\tilde{\tilde{l}}} & 0 & 0 & \phi_1\delta_{\tilde{\tilde{p}}}{}^{\tilde{\tilde{k}}_1} & 0 \\ 0 & \delta\delta^{\tilde{\tilde{q}}}{}_{\tilde{\tilde{d}}} & -\beta\delta^{\tilde{\tilde{q}}}{}_{\tilde{\tilde{k}}} & \eta_{\alpha}{}^{\tilde{\tilde{q}}\tilde{\tilde{l}}} & 0 & 0 & 0 & \phi_1\delta^{\tilde{\tilde{q}}}{}_{\tilde{\tilde{l}}_1} \\ \psi_1\delta_{\tilde{\tilde{a}}_1}{}^{\tilde{\tilde{b}}} & 0 & 0 & 0 & -(\eta^T)_{\alpha\tilde{\tilde{a}}_1\tilde{\tilde{b}}_1} & -\alpha\delta_{\tilde{\tilde{a}}_1}{}^{\tilde{\tilde{d}}_1} & -\gamma\delta_{\tilde{\tilde{a}}_1}{}^{\tilde{\tilde{k}}_1} & 0 \\ 0 & \psi_1\delta^{\tilde{\tilde{c}}_1}{}_{\tilde{\tilde{d}}} & 0 & 0 & -\beta\delta^{\tilde{\tilde{c}}_1}{}_{\tilde{\tilde{b}}_1} & \eta_{\alpha}{}^{\tilde{\tilde{c}}_1\tilde{\tilde{d}}_1} & 0 & -\gamma\delta^{\tilde{\tilde{a}}_1}{}_{\tilde{\tilde{l}}_1} \\ 0 & 0 & \psi_1\delta^{\tilde{\tilde{q}}_1}{}_{\tilde{\tilde{k}}} & 0 & -\delta\delta^{\tilde{\tilde{q}}_1}{}_{\tilde{\tilde{b}}_1} & 0 & \eta_{\alpha}{}^{\tilde{\tilde{q}}_1\tilde{\tilde{k}}_1} & \alpha\delta^{\tilde{\tilde{q}}_1}{}_{\tilde{\tilde{l}}_1} \\ 0 & 0 & 0 & \psi_1\delta_{\tilde{\tilde{p}}_1}{}^{\tilde{\tilde{l}}} & 0 & -\delta\delta_{\tilde{\tilde{p}}_1}{}^{\tilde{\tilde{d}}_1} & \beta\delta_{\tilde{\tilde{p}}_1}{}^{\tilde{\tilde{k}}_1} & -(\eta^T)_{\alpha\tilde{\tilde{p}}_1\tilde{\tilde{l}}_1} \end{pmatrix}}_{\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}}$$

$$\cdot\underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} \delta_{\tilde{\tilde{b}}}{}^{b} & 0 & 0 & 0 \\ 0 & \delta^{\tilde{\tilde{d}}}{}_{d} & 0 & 0 \\ 0 & \delta^{\tilde{\tilde{k}}}{}_{d} & 0 & 0 \\ -i\delta_{\tilde{\tilde{l}}}{}^{b} & 0 & 0 & 0 \\ 0 & 0 & \delta^{\tilde{\tilde{b}}_1}{}_{d_2} & 0 \\ 0 & 0 & 0 & \delta^{\tilde{\tilde{d}}_1}{}_{b_2} \\ 0 & 0 & 0 & \delta^{\tilde{\tilde{k}}_1}{}_{b_2} \\ 0 & 0 & -i\delta^{\tilde{\tilde{l}}_1}{}_{d_2} & 0 \end{pmatrix}}_{:=(\tilde{m}_2^T)_{\tilde{\tilde{B}}}{}^{B}}. \tag{8.26}$$

If we now use the operators $m_{\Lambda}{}^{\tilde{\Lambda}}$ *translating the subspace* $\mathbb{R}^{4}_{(2,2)}$ *in the subspace* $\mathbb{C}^2$ *and leaving unchanged the subspace* $\mathbb{C}^2 \oplus \mathbb{R}^{2n-8}_{(n-4,n-4)}$ *then we can construct the connecting operators* $\eta_{\Lambda}{}^{ab}$ *of the corresponding space* $\mathbb{C}^4 \oplus \mathbb{R}^{2n-8}_{(n-4,n-4)}$*, while the dimension of the spinor space will be reduced by 2 times.*
3. Proceeding similarly, we can obtain the space $\mathbb{C}^{n-2} \oplus \mathbb{R}^{4}_{(2,2)}$ *and the corresponding connecting operators. It is obvious that at each step, only those components of the connecting operators which responsible for the transition from the real representation*

of the subspace $\mathbb{R}^4_{(2,2)}$ to its complex analog $\mathbb{C}^2$ are modified.
4. The final step is carried out using the operators discussed above in Example 3.2.

Corollary 8.2. *The connecting operators from Definition 8.1 satisfy the identity (the proof is given in Appendix)*

$$\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}\eta^\Theta{}_{CB} = \frac{N}{4}(g_{\Lambda\Psi}g^{\Omega\Theta} + \delta_\Lambda{}^\Theta\delta_\Psi{}^\Omega - \delta_\Psi{}^\Theta\delta_\Lambda{}^\Omega). \tag{8.27}$$

Corollary 8.3. *For the orthogonal transformation (6.42), there are the two variants of a decomposition for each case:*

I). Special rotation.

$$S_\Lambda{}^\Psi\eta_\Psi{}^{AB} = \eta_\Lambda{}^{CD}\tilde{S}_C{}^A\tilde{\tilde{S}}_D{}^B. \tag{8.28}$$

1. n mod 4 =2

$$\tilde{\tilde{S}}_A{}^B = -E_A{}^C\tilde{S}_C{}^D E_D{}^B, \quad \tilde{\tilde{S}}_A{}^B = -\tilde{E}_A{}^C\tilde{S}_C{}^D\tilde{E}_D{}^B. \tag{8.29}$$

2. n mod 4 =0

$$\begin{aligned} \tilde{\tilde{S}}_A{}^B\tilde{\tilde{S}}_C{}^D\varepsilon_{BD} = \varepsilon_{AC}, \quad \tilde{S}_A{}^B\tilde{S}_C{}^D\varepsilon_{BD} = \varepsilon_{AC}, \\ \tilde{\tilde{S}}_A{}^B\tilde{\tilde{S}}_C{}^D\tilde{\varepsilon}_{BD} = \tilde{\varepsilon}_{AC}, \quad \tilde{S}_A{}^B\tilde{S}_C{}^D\tilde{\varepsilon}_{BD} = \tilde{\varepsilon}_{AC}. \end{aligned} \tag{8.30}$$

II). Non-special rotation.

$$S_\Lambda{}^\Psi\eta_\Psi{}^{BA} = \eta_{\Lambda CD}\tilde{S}^{CA}\tilde{\tilde{S}}^{DB}. \tag{8.31}$$

1. n mod 4 =2

$$\tilde{\tilde{S}}^{CA} = -E_K{}^A\tilde{S}^{KL}E_L{}^C, \quad \tilde{\tilde{S}}^{CA} = \tilde{E}_K{}^A\tilde{S}^{KL}\tilde{E}_L{}^C. \tag{8.32}$$

2. n mod 4 =0

$$\begin{aligned} \tilde{\tilde{S}}^{AB}\tilde{\tilde{S}}^{CD}\varepsilon_{BD} = \varepsilon^{AC}, \quad \tilde{S}^{AB}\tilde{S}^{CD}\varepsilon_{BD} = \varepsilon^{AC}, \\ \tilde{\tilde{S}}^{AB}\tilde{\tilde{S}}^{CD}\tilde{\varepsilon}_{BD} = \tilde{\varepsilon}^{AC}, \quad \tilde{S}^{AB}\tilde{S}^{CD}\tilde{\varepsilon}_{BD} = \tilde{\varepsilon}^{AC}. \end{aligned} \tag{8.33}$$

Corollary 8.4. *For the involution (6.47), there are the two variants of a decomposition:*

I). Analog of the special rotation.

$$S_\Lambda{}^{\Psi'}\bar{\eta}_{\Psi'}{}^{A'B'} = \eta_\Lambda{}^{CD}\tilde{S}_C{}^{A'}\tilde{\tilde{S}}_D{}^{B'}. \tag{8.34}$$

$$\tilde{\tilde{S}}_A{}^{B'}\bar{\tilde{\tilde{S}}}_{B'}{}^D = \pm\delta_A{}^D, \quad \tilde{S}_A{}^{B'}\bar{\tilde{S}}_{B'}{}^D = \pm\delta_A{}^D. \tag{8.35}$$

II). Analog of the non-special rotation.

$$S_\Lambda{}^{\Psi'}\bar{\eta}_{\Psi'}{}^{B'A'} = \eta_{\Lambda CD}\tilde{S}^{CA'}\tilde{\tilde{S}}^{DB'}. \tag{8.36}$$

$$\tilde{\tilde{S}}^{AB'} = \pm\bar{\tilde{\tilde{S}}}^{B'A}, \quad \tilde{S}^{AB'} = \pm\bar{\tilde{S}}^{B'A}. \tag{8.37}$$

Table 1: The matrix form of the tensor S for the real inclusions of some 2-dimensional spaces in the complex space for $r^i = (T, Z)$.

N	Space	Form of S	S	Isomorphism	Form of $\sqrt{2}r^i\eta_i$	Form of $\sqrt{2}r^i\sigma_i$
1a	R^2 $\ll ++ \gg$	$S_{AA'}$	$\begin{pmatrix} 1 \end{pmatrix}$	$U(1) \cong SO^+(2)$	$T + iZ$	$T - iZ$
1b	R^2 $\ll -- \gg$	$S_{AA'}$	$\begin{pmatrix} i \end{pmatrix}$	$U(1) \cong SO^+(2)$	$-iT - Z$	$-iT + Z$
2a	$R^2_{(1,1)}$ $\ll +- \gg$	$S_A{}^{A'}$	$\begin{pmatrix} 1 \end{pmatrix}$	$R_+ \cong SO^+(1,1)$	$T + Z$	$T - Z$
2b	$R^2_{(1,1)}$ $\ll -+ \gg$	$S_A{}^{A'}$	$\begin{pmatrix} 1 \end{pmatrix}$	$R_+ \cong SO^+(1,1)$	$-T - Z$	$T - Z$

Table 2: The matrix form of the tensor S for the real inclusions of some 4-dimensional spaces in the complex space for $r^i = (T, X, Y, Z)$.

N	Space	S	Form of $\sqrt{2}r^i\eta_i/\sqrt{2}r^i\sigma_i$
1a	R^4 $\ll ---- \gg$ $SU(2){\times}SU(2)/{\pm}1{\cong}SO^+(4)$	$\tilde{S}_A{}^{A'} = \tilde{\tilde{S}}_A{}^{A'}$ $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$	$\begin{pmatrix} -iT-Z & iX+Y \\ iX-Y & iT-Z \end{pmatrix}$ $\begin{pmatrix} -iT+Z & iX+Y \\ iX-Y & iT+Z \end{pmatrix}$
1b	R^4 $\ll ++++ \gg$ $SU(2){\times}SU(2)/{\pm}1{\cong}SO^+(4)$	$\tilde{S}_A{}^{A'} = \tilde{\tilde{S}}_A{}^{A'}$ $\begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$	$\begin{pmatrix} T+iZ & iY+X \\ -iY+X & -T+iZ \end{pmatrix}$ $\begin{pmatrix} T-iZ & iY+X \\ -iY+X & -T-iZ \end{pmatrix}$
2a	$R^4_{(3,1)}$ $\ll +-++ \gg$ $SL(2,\mathbb{C})/{\pm}1{\cong}SO^+(3,1)$	$\tilde{S}_{AA'} = \tilde{\tilde{S}}_{AA'}$ $\begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$	$\begin{pmatrix} T+Z & iY+X \\ -iY+X & -T+Z \end{pmatrix}$ $\begin{pmatrix} T-Z & iY+X \\ -iY+X & -T-Z \end{pmatrix}$
2b	$R^4_{(1,3)}$ $\ll +--- \gg$ $SL(2,\mathbb{C})/{\pm}1{\cong}SO^+(1,3)$	$\tilde{S}_{AA'} = \tilde{\tilde{S}}_{AA'}$ $\begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$	$\begin{pmatrix} T+Z & iY+X \\ iY-X & -T+Z \end{pmatrix}$ $\begin{pmatrix} T-Z & iY+X \\ iY-X & -T-Z \end{pmatrix}$
3a	$R^4_{(2,2)}$ $\ll ++-- \gg$ $SL(2,\mathbb{R}){\times}SL(2,\mathbb{R})/{\pm}1{\cong}SO^+(2,2)$	$\tilde{S}_A{}^{A'} = \tilde{\tilde{S}}_A{}^{A'}$ $\begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$	$\begin{pmatrix} T+iZ & iY+X \\ iY-X & -T+iZ \end{pmatrix}$ $\begin{pmatrix} T-iZ & iY+X \\ iY-X & -T-iZ \end{pmatrix}$
3b	$R^4_{(2,2)}$ $\ll --++ \gg$ $SL(2,\mathbb{R}){\times}SL(2,\mathbb{R})/{\pm}1{\cong}SO^+(2,2)$	$\tilde{S}_A{}^{A'} = \tilde{\tilde{S}}_A{}^{A'}$ $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$	$\begin{pmatrix} -iT-Z & iX+Y \\ -iX+Y & iT-Z \end{pmatrix}$ $\begin{pmatrix} -iT+Z & iX+Y \\ -iX+Y & iT+Z \end{pmatrix}$

Table 3: The matrix form of the tensor S for the real inclusions of some 6-dimensional spaces in the complex space for $r^i = (T, X, V, W, Y, Z)$.

N	Space	Form of S	Isomorphism
			Form of $\sqrt{2}r^i\eta_i$
			Form of $\sqrt{2}r^i\sigma_i$
1	R^6	$S_{AA'}$	$SU(4)/\pm 1 \cong SO^+(6)$
		$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} T+iZ & X+iY & V+iW & 0 \\ X-iY & -T+iZ & 0 & V+iW \\ V-iW & 0 & -T+iZ & -X-iY \\ 0 & V-iW & -X+iY & T+iZ \end{pmatrix}$
			$\begin{pmatrix} T-iZ & X+iY & V+iW & 0 \\ X-iY & -T-iZ & 0 & V+iW \\ V-iW & 0 & -T-iZ & -X-iY \\ 0 & V-iW & -X+iY & T-iZ \end{pmatrix}$
2	$R^6_{(1,5)}$	$S_A{}^{A'}$	$SL(2,\mathbb{H})/\pm 1 \cong SO^+(1,5)$
		$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} T+Z & X+iY & V+iW & 0 \\ -X+iY & -T+Z & 0 & V+iW \\ -V+iW & 0 & -T+Z & -X-iY \\ 0 & -V+iW & X-iY & T+Z \end{pmatrix}$
			$\begin{pmatrix} T-Z & X+iY & V+iW & 0 \\ -X+iY & -T-Z & 0 & V+iW \\ -V+iW & 0 & -T-Z & -X-iY \\ 0 & -V+iW & X-iY & T-Z \end{pmatrix}$
3	$R^6_{(2,4)}$	$S_{AA'}$	$SU(2,2)/\pm 1 \cong SO^+(2,4)$
		$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} T+iZ & X+iY & V+iW & 0 \\ -X+iY & -T+iZ & 0 & V+iW \\ -V+iW & 0 & -T+iZ & -X-iY \\ 0 & -V+iW & X-iY & T+iZ \end{pmatrix}$
			$\begin{pmatrix} T-iZ & X+iY & V+iW & 0 \\ -X+iY & -T-iZ & 0 & V+iW \\ -V+iW & 0 & -T-iZ & -X-iY \\ 0 & -V+iW & X-iY & T-iZ \end{pmatrix}$
4	$R^6_{(3,3)}$	$S_A{}^{A'}$	$SL(4,\mathbb{R})/\pm 1 \cong SO^+(3,3)$
		$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} T+Z & X+iY & V+iW & 0 \\ X-iY & -T+Z & 0 & V+iW \\ -V+iW & 0 & -T+Z & -X-iY \\ 0 & -V+iW & -X+iY & T+Z \end{pmatrix}$
			$\begin{pmatrix} T-Z & X+iY & V+iW & 0 \\ X-iY & -T-Z & 0 & V+iW \\ -V+iW & 0 & -T-Z & -X-iY \\ 0 & -V+iW & -X+iY & T-Z \end{pmatrix}$

We assume that the operators ${\eta_\Lambda}^{AB}(\sigma_{\Lambda AB})$ are constructed either by the scheme (8.20)$(\alpha = \overline{1, n-2})$

$$a). \quad \begin{array}{l} {\eta_\Lambda}^{AB} = \begin{pmatrix} {\eta_\alpha}^{ab} & \frac{1}{2}(i\eta_{n-1} + \eta_n){\delta^a}_d \\ \frac{1}{2}(-i\eta_{n-1} + \eta_n){\delta_c}^b & -(\eta^T)_{\alpha cd} \end{pmatrix}, \\ \sigma_{\Lambda AB} = \begin{pmatrix} (\eta^T)_{\alpha ab} & \frac{1}{2}(i\eta_{n-1} + \eta_n){\delta_a}^d \\ \frac{1}{2}(-i\eta_{n-1} + \eta_n){\delta^c}_b & -{\eta_\alpha}^{cd} \end{pmatrix} \end{array} \tag{8.38}$$

or by the scheme

$$b). \quad \begin{array}{l} {\eta_\Lambda}^{AB} = \begin{pmatrix} {\eta_\alpha}^{ab} & \frac{1}{2}(i\eta_{n-1} + \eta_n){\delta^a}_d \\ \frac{1}{2}(i\eta_{n-1} - \eta_n){\delta_c}^b & -(\eta^T)_{\alpha cd} \end{pmatrix}, \\ \sigma_{\Lambda AB} = \begin{pmatrix} (\eta^T)_{\alpha ab} & \frac{1}{2}(i\eta_{n-1} + \eta_n){\delta_a}^d \\ \frac{1}{2}(i\eta_{n-1} - \eta_n){\delta^c}_b & -{\eta_\alpha}^{cd} \end{pmatrix}. \end{array} \tag{8.38'}$$

This means that from the previous dimension, the conjugating spin-tensors(6.47)

$$\begin{array}{llll} I). & a). \; {S_A}^{A'} := \begin{pmatrix} 0 & iS_{aa'} \\ -iS^{bb'} & 0 \end{pmatrix}, & b). \; {S_A}^{A'} := \begin{pmatrix} 0 & iS_{aa'} \\ iS^{bb'} & 0 \end{pmatrix}, \\ II). & a). \; S_{AA'} := \begin{pmatrix} 0 & i{S_a}^{a'} \\ -i{S^b}_{b'} & 0 \end{pmatrix}, & b). \; S_{AA'} := \begin{pmatrix} 0 & i{S_a}^{a'} \\ i{S^b}_{b'} & 0 \end{pmatrix}, \end{array} \tag{8.39}$$

defined to within a sign, are inherited.

Example 8.3. *In the table 1, the various options for the real inclusion for n=2 are given (see Example 6.1 and Example 6.2).*

Example 8.4. *In the table 2, the various options for the real inclusion for n=4 are given (see Example 6.3).*

Example 8.5. *In the table 3, the various options for the real inclusion for n=6 are given. See Example 8.2 in the case, when* ${\tilde{\eta}_\Lambda}^{AC} := {\eta_\Lambda}^{AB}{\varepsilon_B}^C$ *is an antisymmetric tensor. Recall that* $\eta_i := {H_i}^\Lambda \eta_\Lambda$ *according to (4.4).*

9 On the structural constants of the non-division algebra. Sedenions

In this section it is told about how on the connecting operators which satisfy the Clifford equation to construct the structural constants of the sedenion algebra. Such the algebras are constructed by the inductive transition for n mod 8 = 0 based on the Bott periodicity. We consider the axioms of such the algebras. The conclusion of all results of this section is made on the basis [8], [27, pp. 192-244 (rus)].

We use the fact that for n mod 8 = 0, there are the connecting operators ${\eta_\Lambda}^{AB}$, the inductive construction of which is not the doubling procedure, but is based on the Bott periodicity. This is means by Definition 8.1 that

$${\eta_\Lambda}^{AB} + {\eta_\Lambda}^{BA} = \eta_\Lambda \varepsilon^{AB}, \quad \eta_\Lambda := (\eta_n)_\Lambda \tag{9.1}$$

and ε^{AB} is a symmetric metric tensor in the spinor space. This will give the opportunity to set

$$I).\quad {P_\Lambda}^A := {\eta_\Lambda}^{BA} X_B,\quad {P_\Lambda}^A P_{\Psi A} = g_{\Lambda\Psi},\qquad II).\quad {P_\Lambda}^A := {\eta_\Lambda}^{AB} X_B,\quad {P_\Lambda}^A P_{\Psi A} = g_{\Lambda\Psi} \tag{9.2}$$

for some X_B ($X^A X_A = 2$). Then we can define some structural constants as follows

$$I).\quad {\eta_{\Lambda\Psi}}^\Theta := \sqrt{2}{\eta_\Lambda}^{AB} P_{\Psi A} {P^\Theta}_B,\quad II).\quad {\eta_{\Lambda\Psi}}^\Theta := \sqrt{2}{\eta_\Lambda}^{BA} P_{\Psi A} {P^\Theta}_B. \tag{9.3}$$

Both variants are identical. Therefore, we will further consider Case I). Will they define *group algebra*? The answer is not quite unequivocally defined, and that's why.

1. *First axiom of multiplication* is executed. For any two elements $(r_1)^\Lambda,\ (r_2)^\Psi$, it can always only be one way to construct their *product*

$$(r_1, r_2)^\Theta := (r_1)^\Lambda (r_2)^\Psi {\eta_{\Lambda\Psi}}^\Theta. \tag{9.4}$$

2. *Second axiom of multiplication* is executed. There is uniquely defined *identity element for multiplication* $e := \frac{1}{\sqrt{2}}\eta^\Lambda$.

$$\begin{gathered}\tfrac{1}{\sqrt{2}}\eta^\Lambda {\eta_{\Lambda\Psi}}^\Theta = \varepsilon^{AB} P_{\Psi A} {P^\Theta}_B = {\delta_\Psi}^\Theta,\\ \tfrac{1}{\sqrt{2}}\eta^\Psi {\eta_{\Lambda\Psi}}^\Theta = {\eta_\Lambda}^{AB} \varepsilon_{XA} X^X {P^\Theta}_B = {P_\Lambda}^B {P^\Theta}_B = {\delta_\Lambda}^\Theta.\end{gathered} \tag{9.5}$$

3. For any non-isotropic element, there is uniquely defined *inverse element for multiplication.* For the inclusion $\mathbb{R}^n \subset \mathbb{C}^n$ ($n\ mod\ 8 = 0$), *Third axiom of multiplication* is executed for any nonzero element. Accordingly, with the help of the inclusion operators ${H_i}^\Lambda$, the structure constants ${\eta_{ij}}^k$, which are real not for all spinors X^A, are defined.

$$r^{-1} := \frac{1}{< r, r >}(2 < r, e > e - r),\quad < r_1, r_2 >:= g_{\Lambda\Psi}(r_1)^\Lambda (r_2)^\Psi. \tag{9.6}$$

$$\begin{gathered}(r^{-1})^\Psi = \tfrac{1}{r^\Phi r_\Phi} r^\Omega (\eta_\Omega \eta^\Psi - {\delta_\Omega}^\Psi),\\ r^\Lambda \tfrac{1}{r^\Phi r_\Phi} r^\Omega (\eta_\Omega \eta^\Psi - {\delta_\Omega}^\Psi) {\eta_{\Lambda\Psi}}^\Theta = \tfrac{\sqrt{2}}{r^\Phi r_\Phi} R^{AB} r^\Omega (\eta_\Omega \eta^\Psi - {\delta_\Omega}^\Psi) \eta_{\Psi XA} X^X {P^\Theta}_B =\\ = \tfrac{\sqrt{2}}{r^\Phi r_\Phi} R^{AB} R_{AX} X^X {P^\Theta}_B = \tfrac{1}{\sqrt{2}} X^B {P^\Theta}_B = \tfrac{1}{\sqrt{2}}\eta^\Theta,\\ \tfrac{1}{r^\Phi r_\Phi} r^\Omega (\eta_\Omega \eta^\Lambda - {\delta_\Omega}^\Lambda) r^\Psi {\eta_{\Lambda\Psi}}^\Theta = \tfrac{\sqrt{2}}{r^\Phi r_\Phi} R^{BA} R_{XA} X^X {P^\Theta}_B = \tfrac{1}{\sqrt{2}}\eta^\Theta.\end{gathered} \tag{9.7}$$

These identities, executed for any non-isotropic r, are equivalent to the relation

$${\eta_{\Lambda\Psi}}^\Omega + {{\eta_\Lambda}^\Omega}_\Psi = \sqrt{2}\eta_\Lambda {\delta_\Psi}^\Omega. \tag{9.8}$$

4. *Commutative axiom of multiplication* is not executed. However, there is the identity

$$\tfrac{1}{2}((r_1, r_2) + (r_2, r_1)) = - < r_1, r_2 > e + < r_1, e > r_2 + < r_2, e > r_1. \tag{9.9}$$

$$\begin{gathered}{\eta_{(\Lambda\Psi)}}^\Theta = \sqrt{2}{\eta_{(\Lambda}}^{AB} P_{\Psi)A} {P^\Theta}_B = \sqrt{2}(-{\eta_{(\Lambda}}^{AB} \eta_{\Psi)AX} + 2{\eta_{(\Lambda}}^{AB} \eta_{\Psi)(AX)}) X^X {P^\Theta}_B =\\ = -g_{\Lambda\Psi}\big(\tfrac{1}{\sqrt{2}}\eta^\Theta\big) + 2\big(\tfrac{1}{\sqrt{2}}\eta_{(\Psi}\big){\delta_{\Lambda)}}^\Theta.\end{gathered} \tag{9.10}$$

The special case of this identity

$$r^2 = - < r, r > e + 2 < r, e > r \tag{9.9'}$$

leads to (9.6) and means that there is *power-associativity*. Therefore, *Fourth axiom of multiplication* is not executed. But a replacement can be found.

5. *Associative axiom of multiplication* is not executed. However, there is *alternative-elastic identity*

$$((r_1,r_1),r_2)-(r_1,(r_1,r_2))-(r_1,(r_2,r_1))=((r_2,(r_1,r_1))-((r_2,r_1),r_1)-((r_1,r_2),r_1). \tag{9.11}$$

$$\begin{gathered}
((r_1,r_1),r_2)-(r_1,(r_1,r_2))-(r_1,(r_2,r_1))=-<r_1,r_1>r_2+2<r_1,e>(r_1,r_2)- \\
-2(r_1,-<r_1,r_2>e+<r_1,e>r_2+<r_2,e>r_1)= \\
=-<r_1,r_1>r_2+2<r_1,r_2>r_1-2<r_2,e>(-<r_1,r_1>e+2<r_1,e>r_1)= \\
=2<r_2,e><r_1,r_1>e+(2<r_1,r_2>-4<r_2,e><r_1,e>)r_1-<r_1,r_1>r_2, \\
\\
(r_2,(r_1,r_1))-((r_2,r_1),r_1))-((r_1,r_2),r_1)=-<r_1,r_1>r_2+2<r_1,e>(r_2,r_1)- \\
-2(-<r_1,r_2>e+<r_1,e>r_2+<r_2,e>r_1,r_1)= \\
=-<r_1,r_1>r_2+2<r_1,r_2>r_1-2<r_2,e>(-<r_1,r_1>e+2<r_1,e>r_1)= \\
=2<r_2,e><r_1,r_1>e+(2<r_1,r_2>-4<r_2,e><r_1,e>)r_1-<r_1,r_1>r_2,
\end{gathered} \tag{9.12}$$

Therefore, *Fifth axiom of multiplication* is not executed. But a replacement can be found.

4*. The Clifford equation gives another identity: *flexible identity* [31] (according to Definitions 8.1-8.2, $\varepsilon^{[A(BC)D]}=0$)

$$(r_1r_2)r_1=r_1(r_2r_1). \tag{9.11$'$}$$

$$\begin{gathered}
\tfrac{1}{2}\eta_{(\Lambda|\Psi}{}^{\Theta}\eta_{\Theta|\Omega)}{}^{\Gamma}=\eta_{(\Lambda}{}^{AB}P_{\Omega)C}\eta_{\Theta}{}^{CD}P_{\Psi A}P^{\Theta}{}_{B}P^{\Gamma}{}_{D}= \\
=\eta_{(\Lambda}{}^{CD}P_{\Omega)C}\eta_{\Psi}P^{\Gamma}{}_{D}-\varepsilon^{ABCD}P_{(\Omega|C|}P_{\Lambda)B}P_{\Psi A}P^{\Gamma}{}_{D}= \\
=\eta_{(\Lambda}\delta_{\Omega)}{}^{\Gamma}\eta_{\Psi}-\tfrac{1}{2}g_{\Omega\Lambda}\eta_{\Psi}\eta_{\Gamma}-\varepsilon^{ABCD}P_{(\Omega|C|}P_{\Lambda)B}P_{\Psi A}P^{\Gamma}{}_{D}, \\
\\
\tfrac{1}{2}\eta_{(\Lambda|\Theta}{}^{\Gamma}\eta_{\Psi|\Omega)}{}^{\Theta}=\eta_{(\Lambda}{}^{AB}P_{\Omega)C}\eta_{\Psi}{}^{CD}P_{\Theta A}P^{\Theta}{}_{D}P^{\Gamma}{}_{B}= \\
=\eta_{(\Lambda}P_{\Omega)C}\eta_{\Psi}{}^{CD}P^{\Gamma}{}_{D}-P_{(\Lambda|D|}P_{\Omega)C}\eta_{\Psi}{}^{CD}\eta^{\Gamma}+\eta_{\Theta}{}^{BA}\eta_{\Psi}{}^{CD}P_{(\Lambda|A|}P_{\Omega)C}P^{\Theta}{}_{D}P^{\Gamma}{}_{B}= \\
=\eta_{(\Lambda}P_{\Omega)C}\eta_{\Psi}{}^{CD}P^{\Gamma}{}_{D}-P_{(\Lambda|D|}P_{\Omega)C}\eta_{\Psi}{}^{CD}\eta^{\Gamma}+\eta_{\Psi}{}^{BA}P_{(\Lambda|A|}\eta_{\Omega)}P^{\Gamma}{}_{B}- \\
-\varepsilon^{BACD}P_{(\Lambda|A|}P_{\Omega)C}P^{\Psi}{}_{D}P^{\Gamma}{}_{B}= \\
=\eta_{(\Lambda}\delta_{\Omega)}{}^{\Gamma}\eta_{\Psi}-\tfrac{1}{2}g_{\Omega\Lambda}\eta_{\Psi}\eta_{\Gamma}-\varepsilon^{ABCD}P_{(\Omega|C|}P_{\Lambda)B}P_{\Psi A}P^{\Gamma}{}_{D}.
\end{gathered} \tag{9.12$'$}$$

From (9.9$'$), it follows that *Jordan identity* $(r_1)^k(r_2r_1)=((r_1)^kr_2)r_1,\ \forall k\in\mathbb{Z}$ is executed too.

Such the algebra is normalized in the sense of

$$g_{\Theta\Gamma}\eta_{(\Lambda|\Psi|}{}^{\Theta}\eta_{\Omega)\Phi}{}^{\Gamma}=g_{\Theta\Gamma}\eta^{\Theta}{}_{\Psi(\Lambda}\eta^{\Gamma}{}_{|\Phi|\Omega)} \tag{9.12$''$}$$

that coincides with the condition of *normalizability* (14.97) for n = 8 .

According to the general theory, the sedenion algebra has *0-divisors* for $n\neq 8$.

$$(r_1,r_2)=0,\ \ (r_1)^{\Lambda}(r_2)^{\Psi}\eta_{\Lambda\Psi}{}^{\Omega}=0,\ \ (R_1)^{AB}(R_2)_{DA}X^{D}P^{\Omega}{}_{B}P_{\Omega}{}^{C}=0. \tag{9.13}$$

0-divisors appear because the operator ${P_\Omega}^C$ is degenerate for $n > 8$. However, there exists a sedenion basis for which

$$\begin{gathered}(r_i, e) = (e, r_i), \quad e^2 = e,\\ (r_i, r_j) = -(r_j, r_i), \quad r_i \neq e, \quad r_j \neq e, \quad i \neq j,\\ (r_i)^2 = -e, \quad r_i \neq e.\end{gathered} \tag{9.14}$$

The algebras for which we can construct the basis (9.14) whose elements are not 0-divisors and satisfy *alternative identities*

$$(r_i(r_i, r_j)) = ((r_i, r_i), r_j), \quad ((r_j, r_i), r_i) = (r_j, (r_i, r_i)). \tag{9.15}$$

are the most interesting for the study. Such the algebras are called *sedenion algebras.* Of course, not for all hypercomplex algebras, such the basis exists. Therefore, the hypercomplex algebras will be classified by the maximum number of basic elements, which are not 0-divisors. The inductive construction of the higher dimension hypercomplex metric group alternative-elastic algebra looks as follows.

Algorithm 9.1. 1. *Let us know antisymmetric operators ${\eta_\alpha}^{ab}$ for n mod 8 = 6. Then the connecting operators ${\eta_\Lambda}^{AB}$ for n mod 8 = 0 are constructed according to the scheme*

$$\begin{aligned}{\eta_\Lambda}^{AB} &= \begin{pmatrix} {\eta_\alpha}^{ab} & \frac{1}{2}(i\eta_{n-1} + \eta_n){\delta^a}_d \\ \frac{1}{2}(-i\eta_{n-1} + \eta_n){\delta_c}^b & -(\eta^T)_{\alpha cd} \end{pmatrix},\\ \sigma_{\Lambda AB} &= \begin{pmatrix} (\eta^T)_{\alpha ab} & \frac{1}{2}(i\eta_{n-1} + \eta_n){\delta_a}^d \\ \frac{1}{2}(-i\eta_{n-1} + \eta_n){\delta^c}_b & -{\eta_\alpha}^{cd} \end{pmatrix},\end{aligned} \tag{9.16}$$

that would provide the existence of the symmetric metric spin-tensor $\tilde{\varepsilon}_{AB} \equiv \varepsilon_{AB}$ for n mod 8 = 0.

2. *Let us know the connecting operators ${\eta_\alpha}^{ab}$ for n mod 8 = 0. Then the connecting operators ${\eta_\Lambda}^{AB}$ for n mod 8 = 2 are constructed according to the scheme*

$$\begin{aligned}{\eta_\Lambda}^{AB} &= \begin{pmatrix} {\eta_\alpha}^{ab} & \frac{1}{2}(i\eta_{n-1} + \eta_n){\delta^a}_d \\ \frac{1}{2}(-i\eta_{n-1} + \eta_n){\delta_c}^b & -(\eta^T)_{\alpha cd} \end{pmatrix},\\ \sigma_{\Lambda AB} &= \begin{pmatrix} (\eta^T)_{\alpha ab} & \frac{1}{2}(i\eta_{n-1} + \eta_n){\delta_a}^d \\ \frac{1}{2}(-i\eta_{n-1} + \eta_n){\delta^c}_b & -{\eta_\alpha}^{cd} \end{pmatrix}.\end{aligned} \tag{9.17}$$

3. *Let us know the connecting operators ${\eta_\alpha}^{ab}$ for n mod 8 = 2. Then the connecting operators ${\eta_\Lambda}^{AB}$ for n mod 8 = 4 are constructed according to the scheme*

$$\begin{aligned}{\eta_\Lambda}^{AB} &= \begin{pmatrix} {\eta_\alpha}^{ab} & \frac{1}{2}(\eta_{n-1} + i\eta_n){\delta^a}_d \\ \frac{1}{2}(-\eta_{n-1} + i\eta_n){\delta_c}^b & (\eta^T)_{\alpha cd} \end{pmatrix},\\ \sigma_{\Lambda AB} &= \begin{pmatrix} (\eta^T)_{\alpha ab} & -\frac{1}{2}(\eta_{n-1} + i\eta_n){\delta_a}^d \\ -\frac{1}{2}(-\eta_{n-1} + i\eta_n){\delta^c}_b & {\eta_\alpha}^{cd} \end{pmatrix},\end{aligned} \tag{9.18}$$

providing for n mod 8 = 4 the existence of the antisymmetric metric spin-tensor

$$\varepsilon_{AB} := (\varepsilon_1)_{AC}(\varepsilon_2)^{DK}(\varepsilon_3)_{BK} = \begin{pmatrix} 0 & 0 & 0 & i(\varepsilon_1)_{ab} \\ 0 & 0 & -i(\varepsilon_1)^{cd} & 0 \\ 0 & i(\varepsilon_1)^{kl} & 0 & 0 \\ -i(\varepsilon_1)_{mn} & 0 & 0 & 0 \end{pmatrix}, \tag{9.19}$$

where $(\varepsilon_1)_{ab}$ is the symmetric metric spin-tensor for n mod 8 = 0.

4. *Let us know the connecting operators $\eta_\alpha{}^{ab}$ for n mod 8 = 4. Then the connecting operators $\eta_\Lambda{}^{AB}$ for n mod 8 = 6 are constructed according to the scheme*

$$\begin{aligned} \eta_\Lambda{}^{AB} &= \begin{pmatrix} \eta_\alpha{}^{ab} & \frac{1}{2}(\eta_{n-1}+i\eta_n)\delta^a{}_d \\ \frac{1}{2}(-\eta_{n-1}+i\eta_n)\delta_c{}^b & (\eta^T)_{\alpha cd} \end{pmatrix}, \\ \sigma_{\Lambda AB} &= \begin{pmatrix} (\eta^T)_{\alpha ab} & -\frac{1}{2}(\eta_{n-1}+i\eta_n)\delta_a{}^d \\ -\frac{1}{2}(-\eta_{n-1}+i\eta_n)\delta^c{}_b & \eta_\alpha{}^{cd} \end{pmatrix}. \end{aligned} \tag{9.20}$$

Let's, in addition, spend the transformation of the connecting operators according to the scheme

$$\tilde{\eta}_\Lambda{}^{AC} := \eta_\Lambda{}^{AB}\varepsilon_B{}^C, \quad \varepsilon_B{}^C := \begin{pmatrix} 0 & -i\varepsilon_{bm} \\ i\varepsilon^{dk} & 0 \end{pmatrix}, \tag{9.20$'$}$$

where ε_{ab} $(n/2+q=3)$ for n mod 8 = 4 will be the metric antisymmetric spin-tensor. Then for the next implementation n mod 8 = 6 the operators $\tilde{\eta}_\Lambda{}^{AC} = -\tilde{\eta}_\Lambda{}^{CA}$ will be completely antisymmetric $(n/2+q)=0$. It is easy to verify that the Clifford equation (8.1) is executed for such the operators. If the further construction for n mod 8 = 0 to lead on the basis of such the operators then there will exist the symmetric metric spin-tensor $\varepsilon_{AB} \equiv (\varepsilon_1)_{AB}$, $n/2+q=1$. For n mod 8 = 2, there will exist the involution at $n/2+q=2$. And again, for n mod 8 = 4, we obtain $n/2+q=3$. The circle has become isolated. The beginning has been made in Examples 8.1-8.2.

5. *Suppose there is an algebra over $\mathbb{R}^n$ with the structural constants generated from the connecting operators $\eta_\Lambda{}^{AB}$ with the metric spin-tensor ε^{XZ} and the inclusion operator $H_i{}^\Lambda$. We assume that the metric tensor $g_{\Lambda\Psi}$ on the main diagonal contains $\ll + \gg$ only. Then we can construct the antisymmetric operators for the space $\mathbb{C}^{n+6}$*

$$\eta_\alpha{}^{ab} = -\eta_\alpha{}^{ba} :=$$
$$\begin{pmatrix} 0 & 0 & 0 & \xi\varepsilon^{AQ} & 0 & \gamma\varepsilon^{AK} & -\alpha\varepsilon^{AD} & \eta_\Lambda{}^A{}_B \\ 0 & 0 & -\xi\varepsilon_{CR} & 0 & -\gamma\varepsilon_{CM} & 0 & (\eta^T)_{\Lambda C}{}^D & \beta\varepsilon_{CB} \\ 0 & \xi\varepsilon_{NY} & 0 & 0 & -\alpha\varepsilon_{NM} & (\eta^T)_{\Lambda N}{}^K & 0 & \delta\varepsilon_{NB} \\ -\xi\varepsilon^{LZ} & 0 & 0 & 0 & \eta_\Lambda{}^L{}_M & \beta\varepsilon^{LK} & -\delta\varepsilon^{LD} & 0 \\ 0 & \gamma\varepsilon_{PY} & \alpha\varepsilon_{PR} & -(\eta^T)_{\Lambda P}{}^Q & 0 & 0 & 0 & -\zeta\varepsilon_{PB} \\ -\gamma\varepsilon^{SZ} & 0 & -\eta_\Lambda{}^S{}_R & -\beta\varepsilon^{SQ} & 0 & 0 & \zeta\varepsilon^{SD} & 0 \\ \alpha\varepsilon^{XZ} & -\eta_\Lambda{}^X{}_Y & 0 & \delta\varepsilon^{XQ} & 0 & -\zeta\varepsilon^{XK} & 0 & 0 \\ -(\eta^T)_{\Lambda T}{}^Z & -\beta\varepsilon_{TY} & -\delta\varepsilon_{TR} & 0 & \zeta\varepsilon_{TM} & 0 & 0 & 0 \end{pmatrix},$$

$$\begin{aligned} \alpha &:= \tfrac{1}{2}(i\eta_{n+1}+\eta_{n+2}), & \beta &:= \tfrac{1}{2}(-i\eta_{n+1}+\eta_{n+2}), \\ \gamma &:= \tfrac{1}{2}(\eta_{n+3}+i\eta_{n+4}), & \delta &:= \tfrac{1}{2}(-\eta_{n+3}+i\eta_{n+4}), \\ \xi &:= \tfrac{1}{2}(\eta_{n+5}+i\eta_{n+6}), & \zeta &:= \tfrac{1}{2}(-\eta_{n+5}+i\eta_{n+6}). \end{aligned} \tag{9.21}$$

6. *The transition to the connecting operators of the space $\mathbb{C}^{n+8}$ is carried out as follows*

$$\eta_\Lambda{}^{AB} := \begin{pmatrix} \eta_\alpha{}^{ab} & \phi\delta^a{}_d \\ \psi\delta_c{}^b & -(\eta^T)_{\alpha cd} \end{pmatrix}, \tag{9.22}$$

$$\phi := \tfrac{1}{2}(i\eta_{n+7}+\eta_{n+8}), \quad \psi := \tfrac{1}{2}(-i\eta_{n+7}+\eta_{n+8})$$

with the metric spin-trnsor $\varepsilon^{XZ} := \begin{pmatrix} 0 & \delta^a{}_d \\ \delta_c{}^b & 0 \end{pmatrix}$, $\varepsilon_{XZ} := \begin{pmatrix} 0 & \delta_a{}^d \\ \delta^c{}_b & 0 \end{pmatrix}$. *Then we pass to the connecting operators of the space* $\mathbb{R}^{n+8} \subset \mathbb{C}^{n+8}$ *using the corresponding inclusion operator. And such the operators generate the structural constants of the algebra with dimension equal to* $n+8$.

10 Infinitesimal transformations

In this section, it is told about how on the connecting operators which satisfies the Clifford equation, to construct the infinitesimal transformations. The conclusion of all results of this section is made on the basis [22, v. 1, pp. 175-177 (eng)].

Using Corollary 8.3 for the special orthogonal transformations, the decomposition

$$\begin{array}{ll} 1). & S_\Lambda{}^\Psi \eta_\Psi{}^{AB} = -\eta_\Lambda{}^{CD} S_C{}^A E_D{}^K S_K{}^M E_M{}^B, \\ 2). & S_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \eta_\Lambda{}^{CD} S_C{}^A (\eta_I)^\Omega S_\Omega{}^\Phi \eta_\Phi{}^{KB} \varepsilon_{KL} S_X{}^L \varepsilon^{MX} (\varepsilon_I)_{MD}. \end{array} \tag{10.1}$$

is obtained. Suppose that there is *one-parameter family of orthogonal transformations*

$$S_\Lambda{}^\Psi(\lambda) S_\Omega{}^\Phi(\lambda) g_{\Psi\Phi} = g_{\Lambda\Omega}, \quad S_\Lambda{}^\Psi(0) := \delta_\Lambda{}^\Psi \tag{10.2}$$

that induces the one-parameter transformations in the spinor space

$$S_C{}^A(\lambda), \quad S_C{}^A(0) = \delta_C{}^A. \tag{10.3}$$

Infinitesimal transformation is defined as

$$\begin{array}{ll} T_\Lambda{}^\Psi = \left[\frac{d}{d\lambda} S_\Lambda{}^\Psi(\lambda)\right]\Big|_{\lambda=0}, & T_{\Lambda\Psi} = -T_{\Psi\Lambda}, \\ T_C{}^A = \left[\frac{d}{d\lambda} S_C{}^A(\lambda)\right]\Big|_{\lambda=0}, & \begin{array}{ll} 1). & T_C{}^A = -E_C{}^B T_B{}^D E_D{}^A, \\ 2). & T_C{}^A = -T^A{}_C. \end{array} \end{array} \tag{10.4}$$

Differentiating in zero (10.1), we will obtain the identity (the calculations are given in Appendix)

$$T_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \eta_\Lambda{}^{CB} T_C{}^A - \eta_\Lambda{}^{AD} (\varepsilon_I)^{KB} T_K{}^M (\varepsilon_I)_{MD} + \eta_\Lambda{}^{AD} (\varepsilon_I)_{MD} (\eta_I)^\Omega T_\Omega{}^\Phi \eta_\Phi{}^{MB}. \tag{10.5}$$

Particular solution of this equation will have the form

$$T_C{}^A = \frac{1}{2} T^{\Theta\Phi} \eta_\Phi{}^{AB} \eta_{\Theta CB}, \tag{10.6}$$

that is verified by the direct substitution. *Homogeneous solution* of the equation

$$\eta_\Lambda{}^{CB} T_C{}^A - \eta_\Lambda{}^{AD} (\varepsilon_I)^{KB} T_K{}^M (\varepsilon_I)_{MD} = 0 \tag{10.7}$$

is rewritten as (8.21) that means triviality of such the solution.

11 Complex and real connections

In this section, it is told about how using some particular solutions of the Clifford equation, to construct a prolongation of the Riemannian torsion-free connection into the spinor bundle. We consider the two different variants. This will give the opportunity to construct the spinor analogues of the Lie operators. The conclusion of all results of this section is made on the basis [9], [22, v. 1, pp. 228-231 (eng)].

For the real representation of the tangent bundle $\tau^{\mathbb{R}}(V_{2n})$, the equation of the geodesics take the form

$$\frac{d^2x^{\tilde\Omega}}{ds^2}+{\Gamma_{\tilde\Phi\tilde\Psi}}^{\tilde\Omega}\,\frac{dx^{\tilde\Phi}}{ds}\frac{dx^{\tilde\Psi}}{ds}=0. \tag{11.1}$$

For the parameterization of the atlas $w^{\Theta}=\frac{1}{\sqrt{2}}(u^{\Theta}(x^{\tilde\Theta})+iv^{\Theta}(x^{\tilde\Theta}))$, we obtain

$$\begin{gathered}\frac{d^2x^{\tilde\Omega}}{ds^2}=\frac{d}{ds}\Big(\frac{\partial x^{\tilde\Omega}}{\partial w^{\Theta}}\,\frac{dw^{\Theta}}{ds}+\frac{\partial x^{\tilde\Omega}}{\partial w^{\Theta'}}\,\frac{dw^{\Theta'}}{ds}\Big)=\\ \frac{\partial^2x^{\tilde\Omega}}{\partial w^{\Theta}ds}\,\frac{dw^{\Theta}}{ds}+\frac{\partial x^{\tilde\Omega}}{\partial w^{\Theta}}\,\frac{d^2w^{\Theta}}{ds^2}+\frac{\partial^2x^{\tilde\Omega}}{\partial w^{\Theta'}ds}\,\frac{dw^{\Theta'}}{ds}+\frac{\partial x^{\tilde\Omega}}{\partial w^{\Theta'}}\,\frac{d^2w^{\Theta'}}{ds^2},\\ \frac{d^2w^{\Theta}}{ds^2}\,\frac{\partial x^{\tilde\Omega}}{\partial w^{\Theta}}+\frac{d^2w^{\Theta'}}{ds^2}\,\frac{\partial x^{\tilde\Omega}}{\partial w^{\Theta'}}+\frac{\partial^2x^{\tilde\Omega}}{\partial w^{\Theta}ds}\,\frac{dw^{\Theta}}{ds}+\frac{\partial^2x^{\tilde\Omega}}{\partial w^{\Theta'}ds}\,\frac{dw^{\Theta'}}{ds}+\\ +{\Gamma_{\tilde\Phi\tilde\Psi}}^{\tilde\Omega}\frac{dx^{\tilde\Phi}}{ds}\,\frac{\partial x^{\tilde\Psi}}{\partial w^{\Theta}}\,\frac{dw^{\Theta}}{ds}+{\Gamma_{\tilde\Phi\tilde\Psi}}^{\tilde\Omega}\frac{dx^{\tilde\Phi}}{ds}\,\frac{\partial x^{\tilde\Psi}}{\partial w^{\Theta'}}\,\frac{dw^{\Theta'}}{ds}=0.\end{gathered} \tag{11.2}$$

We multiply the both sides by $\frac{dw^{\Lambda}}{dx^{\tilde\Omega}}={m^{\Lambda}}_{\tilde\Omega}$. Considering the identities (3.8) for the operators ${m^{\Lambda}}_{\tilde\Omega}$ and ${m^{\Lambda'}}_{\tilde\Omega}$, we will obtain

$$\begin{gathered}\frac{d^2w^{\Lambda}}{ds^2}+\Big(\frac{\partial w^{\Lambda}}{\partial x^{\tilde\Omega}}\,\frac{\partial^2x^{\tilde\Omega}}{\partial w^{\Theta}ds}\,\frac{dw^{\Theta}}{ds}+{\Gamma_{\tilde\Phi\tilde\Psi}}^{\tilde\Omega}\frac{\partial x^{\tilde\Psi}}{\partial w^{\Theta}}\frac{dx^{\tilde\Phi}}{ds}\,\frac{dw^{\Theta}}{ds}\,\frac{\partial w^{\Lambda}}{\partial x^{\tilde\Omega}}\Big)+\\ +\Big(\frac{\partial w^{\Lambda}}{\partial x^{\tilde\Omega}}\,\frac{\partial^2x^{\tilde\Omega}}{\partial w^{\Theta'}ds}\,\frac{dw^{\Theta'}}{ds}+{\Gamma_{\tilde\Phi\tilde\Psi}}^{\tilde\Omega}\frac{\partial x^{\tilde\Psi}}{\partial w^{\Theta'}}\frac{dx^{\tilde\Phi}}{ds}\,\frac{dw^{\Theta'}}{ds}\frac{\partial w^{\Lambda}}{\partial x^{\tilde\Omega}}\Big)=0.\end{gathered} \tag{11.3}$$

For the complex representation of the tangent bundle $\tau^{\mathbb{C}}(V_{2n})$, the geodesic equations take the form

$$\frac{d^2w^{\Lambda}}{ds^2}+\Big({\Gamma_{\tilde\Phi\Theta}}^{\Lambda}\,\frac{dw^{\Theta}}{ds}+{\Gamma_{\tilde\Phi\Theta'}}^{\Lambda}\,\frac{dw^{\Theta'}}{ds}\Big)\,\frac{dx^{\tilde\Phi}}{ds}=0. \tag{11.4}$$

Comparing these two equations with Definition 3.8 of the operators ${m^{\Lambda}}_{\tilde\Omega}$, ${m_{\Lambda}}^{\tilde\Omega}$, we obtain

$$\left\{\begin{array}{l}{m^{\Lambda}}_{\tilde\Omega}\ (\partial_{\tilde\Phi}{m_{\Theta}}^{\tilde\Omega}+{\Gamma_{\tilde\Phi\tilde\Psi}}^{\tilde\Omega}{m_{\Theta}}^{\tilde\Psi})={\Gamma_{\tilde\Phi\Theta}}^{\Lambda},\\ {m^{\Lambda}}_{\tilde\Omega}\ (\partial_{\tilde\Phi}{\bar m_{\Lambda'}}{}^{\tilde\Omega}+{\Gamma_{\tilde\Phi\tilde\Psi}}^{\tilde\Omega}{\bar m_{\Theta'}}{}^{\tilde\Psi})={\Gamma_{\tilde\Phi\Theta'}}^{\Lambda}\end{array}\right. \tag{11.5}$$

taking into account the identity

$$\frac{\partial^2x^{\tilde\Omega}}{\partial w^{\Theta}ds}=\frac{\partial^2x^{\tilde\Omega}}{\partial w^{\Theta}\partial x^{\tilde\Phi}}\frac{dx^{\tilde\Phi}}{ds}=\frac{dx^{\tilde\Phi}}{ds}\partial_{\tilde\Phi}\frac{\partial x^{\tilde\Omega}}{\partial w^{\Theta}}=\frac{dx^{\tilde\Phi}}{ds}\partial_{\tilde\Phi}{m_{\Theta}}^{\tilde\Omega}. \tag{11.6}$$

Other record of the system has the form

$$\begin{gathered}\left\{\begin{array}{l}{m_{\Xi}}^{\tilde\Psi}\ (-\partial_{\tilde\Phi}{m^{\Theta}}_{\tilde\Psi}+{\Gamma_{\tilde\Phi\tilde\Psi}}^{\tilde\Omega}{m^{\Theta}}_{\tilde\Omega})={\Gamma_{\tilde\Phi\Xi}}^{\Theta}\ \text{, multiply by }\ {m^{\Xi}}_{\tilde\Lambda},\\ {\bar m_{\Xi'}}{}^{\tilde\Psi}\ (-\partial_{\tilde\Phi}{m^{\Theta}}_{\tilde\Psi}+{\Gamma_{\tilde\Phi\tilde\Psi}}^{\tilde\Omega}{m^{\Theta}}_{\tilde\Omega})={\Gamma_{\tilde\Phi\Xi'}}^{\Theta}\ \text{, multiply by }\ {\bar m^{\Xi'}}{}_{\tilde\Lambda},\end{array}\right.\\ \\ \left\{\begin{array}{l}{\triangle_{\tilde\Lambda}}^{\tilde\Psi}\ (-\partial_{\tilde\Phi}{m^{\Lambda}}_{\tilde\Psi}+{\Gamma_{\tilde\Phi\tilde\Psi}}^{\tilde\Omega}{m^{\Lambda}}_{\tilde\Omega})={\Gamma_{\tilde\Phi\Xi}}^{\Lambda}{m^{\Xi}}_{\tilde\Lambda},\\ {\bar\triangle_{\tilde\Lambda}}{}^{\tilde\Psi}\ (-\partial_{\tilde\Phi}{m^{\Lambda}}_{\tilde\Psi}+{\Gamma_{\tilde\Phi\tilde\Psi}}^{\tilde\Omega}{m^{\Lambda}}_{\tilde\Omega})={\Gamma_{\tilde\Phi\Xi'}}^{\Lambda}{\bar m^{\Xi'}}{}_{\tilde\Lambda}.\end{array}\right.\end{gathered} \tag{11.7}$$

Let's add these two equations

$$\partial_{\tilde{\varPhi}} m^{\Lambda}{}_{\tilde{\varLambda}} - \Gamma_{\tilde{\varPhi}\tilde{\varLambda}}{}^{\tilde{\varOmega}} m^{\Lambda}{}_{\tilde{\varOmega}} + \Gamma_{\tilde{\varPhi}\Xi}{}^{\Lambda} m^{\Xi}{}_{\tilde{\varLambda}} + \Gamma_{\tilde{\varPhi}\Xi'}{}^{\Lambda} \bar{m}^{\Xi'}{}_{\tilde{\varLambda}} = 0. \tag{11.8}$$

We will contract it with $\bar{m}_{\Psi'}{}^{\tilde{\varLambda}}$

$$\begin{gathered} \bar{m}_{\Psi'}{}^{\tilde{\varLambda}} \partial_{\tilde{\varPhi}} m^{\Lambda}{}_{\tilde{\varLambda}} - \Gamma_{\tilde{\varPhi}\tilde{\varLambda}}{}^{\tilde{\varOmega}} m^{\Lambda}{}_{\tilde{\varOmega}} \bar{m}_{\Psi'}{}^{\tilde{\varLambda}} + \Gamma_{\tilde{\varPhi}\Psi'}{}^{\Lambda} = 0, \\ \Gamma_{\tilde{\varPhi}\Psi'}{}^{\Lambda} = \Gamma_{\tilde{\varPhi}\tilde{\varLambda}}{}^{\tilde{\varOmega}} m^{\Lambda}{}_{\tilde{\varOmega}} \bar{m}_{\Psi'}{}^{\tilde{\varLambda}} - \tfrac{i}{2} \bar{m}_{\Psi'}{}^{\tilde{\varLambda}} m^{\Lambda}{}_{\tilde{\varTheta}} \partial_{\tilde{\varPhi}} f_{\tilde{\varLambda}}{}^{\tilde{\varTheta}}. \end{gathered} \tag{11.9}$$

Therefore, the connection in the tangent bundle should be predetermined by the system

$$\begin{cases} \nabla_{\tilde{\varPhi}} m^{\Lambda}{}_{\tilde{\varLambda}} = 0, \\ \nabla_{\tilde{\varPhi}} \bar{m}^{\Lambda'}{}_{\tilde{\varLambda}} = 0. \end{cases} \tag{11.10}$$

Indeed, since $m^{\Psi}{}_{\tilde{\varOmega}} = i\ f_{\tilde{\varOmega}}{}^{\tilde{\varLambda}} m^{\Psi}{}_{\tilde{\varLambda}}$ then

$$\begin{gathered} \nabla_{\tilde{\varPhi}} f_{\tilde{\varOmega}}{}^{\tilde{\varLambda}} = 0, \\ \partial_{\tilde{\varPhi}} f_{\tilde{\varOmega}}{}^{\tilde{\varLambda}} = (\Gamma_{\tilde{\varPhi}\tilde{\varOmega}}{}^{\tilde{\varTheta}} f_{\tilde{\varTheta}}{}^{\tilde{\varLambda}} - \Gamma_{\tilde{\varPhi}\tilde{\varTheta}}{}^{\tilde{\varLambda}} f_{\tilde{\varOmega}}{}^{\tilde{\varTheta}}). \end{gathered} \tag{11.11}$$

We will multiply this equation by $\frac{i}{2} \bar{m}_{\Psi'}{}^{\tilde{\varOmega}} m^{\Lambda}{}_{\tilde{\varLambda}}$

$$\frac{i}{2} \bar{m}_{\Psi'}{}^{\tilde{\varOmega}} m^{\Lambda}{}_{\tilde{\varLambda}} \partial_{\tilde{\varPhi}} f_{\tilde{\varOmega}}{}^{\tilde{\varLambda}} = \bar{m}_{\Psi'}{}^{\tilde{\varOmega}} m^{\Lambda}{}_{\tilde{\varLambda}} \Gamma_{\tilde{\varPhi}\tilde{\varOmega}}{}^{\tilde{\varLambda}}. \tag{11.12}$$

Hence,

$$\Gamma_{\tilde{\varPhi}\Psi'}{}^{\Lambda} = 0. \tag{11.13}$$

Therefore, we can write down

$$\nabla_{\tilde{\varPhi}} m^{\Lambda}{}_{\tilde{\varLambda}} := \partial_{\tilde{\varPhi}} m^{\Lambda}{}_{\tilde{\varLambda}} - \Gamma_{\tilde{\varPhi}\tilde{\varLambda}}{}^{\tilde{\varOmega}} m^{\Lambda}{}_{\tilde{\varOmega}} + \Gamma_{\tilde{\varPhi}\Psi}{}^{\Lambda} m^{\Psi}{}_{\tilde{\varLambda}} = 0. \tag{11.14}$$

Thus, if we know the coefficients of the connection in one of the representation (real or complex) then the connection coefficients in the other representation (complex or real, respectively) are uniquely restored from this equation. If now in the real representation of the tangent bundle, the torsion-free connection, compatible with the metric

$$\nabla_{\tilde{\varLambda}} G_{\tilde{\varPsi}\tilde{\varOmega}} = 0, \tag{11.15}$$

is given then it can be prolonged into the spinor bundle with the help of the condition

$$\nabla_{\tilde{\varLambda}} \eta_{\tilde{\varPsi}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}} = 0. \tag{11.16}$$

Now let I). n mod 4 = 0, n> 2, II). n mod 4 = 2, n> 4, then

$$
\begin{aligned}
I).\qquad & \nabla_{\tilde{\Psi}}\eta_{\tilde{\Lambda}\tilde{\tilde{C}}\tilde{\tilde{D}}}=0,\\
& \nabla_{\tilde{\Psi}}\varepsilon_{\tilde{\tilde{A}}\tilde{\tilde{C}}}\varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}}=0,\\
& \varepsilon_{\tilde{\tilde{A}}\tilde{\tilde{C}}}\nabla_{\tilde{\Psi}}\varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}}+\varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}}\nabla_{\tilde{\Psi}}\varepsilon_{\tilde{\tilde{A}}\tilde{\tilde{C}}}=0,\\
& \eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}(\delta_{\tilde{\tilde{A}}}{}^{\tilde{\tilde{K}}}\varepsilon^{\tilde{\tilde{L}}\tilde{\tilde{D}}}\nabla_{\tilde{\Psi}}\varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}}+\delta_{\tilde{\tilde{B}}}{}^{\tilde{\tilde{L}}}\varepsilon^{\tilde{\tilde{K}}\tilde{\tilde{C}}}\nabla_{\tilde{\Psi}}\varepsilon_{\tilde{\tilde{A}}\tilde{\tilde{C}}})=0,\\
& \eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{K}}\tilde{\tilde{B}}}\varepsilon^{\tilde{\tilde{L}}\tilde{\tilde{D}}}\nabla_{\tilde{\Psi}}\varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}}+\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{B}}\tilde{\tilde{L}}}\varepsilon^{\tilde{\tilde{K}}\tilde{\tilde{C}}}\nabla_{\tilde{\Psi}}\varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{C}}}=0,\\
\\
II).\qquad & \nabla_{\tilde{\Psi}}\eta_{\tilde{\Lambda}\tilde{\tilde{C}}\tilde{\tilde{D}}}=0,\\
& \tilde{\nabla}_{\tilde{\Psi}}E_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{A}}}E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}=0,\\
& E_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{A}}}\tilde{\nabla}_{\tilde{\Psi}}E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}+E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}\tilde{\nabla}_{\tilde{\Psi}}E_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{A}}}=0,\\
& \eta_{\tilde{\Lambda}\tilde{\tilde{A}}\tilde{\tilde{B}}}(\delta_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}E_{\tilde{\tilde{L}}}{}^{\tilde{\tilde{C}}}\nabla_{\tilde{\Psi}}E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}+\delta_{\tilde{\tilde{L}}}{}^{\tilde{\tilde{B}}}E_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{D}}}\nabla_{\tilde{\Psi}}E_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{A}}})=0,\\
& \eta_{\tilde{\Lambda}\tilde{\tilde{K}}\tilde{\tilde{B}}}E_{\tilde{\tilde{L}}}{}^{\tilde{\tilde{C}}}\nabla_{\tilde{\Psi}}E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}+\eta_{\tilde{\Lambda}\tilde{\tilde{A}}\tilde{\tilde{L}}}E_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{D}}}\nabla_{\tilde{\Psi}}E_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{A}}}=0.
\end{aligned}
\tag{11.17}
$$

Turn to the equation (8.21). This will give

$$
I).\quad \varepsilon^{\tilde{\tilde{L}}\tilde{\tilde{D}}}\nabla_{\tilde{\Psi}}\varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}}=0,\qquad II).\quad E_{\tilde{\tilde{L}}}{}^{\tilde{\tilde{C}}}\nabla_{\tilde{\Psi}}E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}=0. \tag{11.18}
$$

Thus, we see that

$$
\begin{aligned}
I).\quad & \nabla_{\tilde{\Psi}}\varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}}=0, & II).\quad & \nabla_{\tilde{\Psi}}E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}=0,\\
& \nabla_{\tilde{\Psi}}\tilde{\varepsilon}_{\tilde{\tilde{B}}\tilde{\tilde{D}}}=0, & & \nabla_{\tilde{\Psi}}\tilde{E}_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}=0.
\end{aligned}
\tag{11.19}
$$

Therefore, the condition (11.16) imposes the restrictions

$$
\begin{aligned}
I).\quad & \nabla_{\tilde{\Psi}}\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}\varepsilon_{\tilde{\tilde{A}}\tilde{\tilde{B}}}=0, & II).\quad & \nabla_{\tilde{\Psi}}\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta_{\tilde{\Phi}\tilde{\tilde{A}}\tilde{\tilde{C}}}E_{\tilde{\tilde{B}}}{}^{\tilde{\tilde{C}}}=0,\\
& \nabla_{\tilde{\Psi}}\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}\tilde{\varepsilon}_{\tilde{\tilde{A}}\tilde{\tilde{B}}}=0, & & \nabla_{\tilde{\Psi}}\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta_{\tilde{\Phi}\tilde{\tilde{A}}\tilde{\tilde{C}}}\tilde{E}_{\tilde{\tilde{B}}}{}^{\tilde{\tilde{C}}}=0.
\end{aligned}
\tag{11.20}
$$

This kind of the connection is possible with the normalization of the spinor basis according to [22, v. 1, p. 230 (eng)]. However, we can relax the conditions (11.15) and (11.16) remembering that we consider the connection in the tangent bundle. Therefore, it is necessary to go back

$$
\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{B}}}\nabla_{\tilde{\Lambda}}\eta_{\tilde{\Omega}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}=0 \tag{11.21}
$$

which can be rewritten as

$$
\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{B}}}\partial_{\tilde{\Lambda}}\eta_{\tilde{\Omega}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}-\Gamma_{\tilde{\Lambda}\tilde{\Omega}}{}^{\tilde{\Theta}}\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta_{\tilde{\Theta}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}+\tilde{\tilde{\Gamma}}_{\tilde{\Lambda}\tilde{\tilde{D}}}{}^{\tilde{\tilde{A}}}\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta_{\tilde{\Omega}}{}^{\tilde{\tilde{D}}\tilde{\tilde{B}}}+\tilde{\Gamma}_{\tilde{\Lambda}\tilde{\tilde{D}}}{}^{\tilde{\tilde{B}}}\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta_{\tilde{\Omega}}{}^{\tilde{\tilde{A}}\tilde{\tilde{D}}}=0. \tag{11.22}
$$

Therefore, in order to prolong uniquely the constructed connection into the spinor bundle, it is necessary to demand

$$
\begin{aligned}
\tilde{\tilde{\Gamma}}_{\tilde{\Lambda}\tilde{\tilde{L}}}{}^{\tilde{\tilde{M}}} &:= \tfrac{1}{N^2}\tilde{\tilde{\Gamma}}_{\tilde{\Lambda}\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}(\eta_{\tilde{\Psi}\tilde{\tilde{L}}\tilde{\tilde{B}}}\eta_{\tilde{\Omega}}{}^{\tilde{\tilde{M}}\tilde{\tilde{B}}}\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{C}}}\eta^{\tilde{\Omega}\tilde{\tilde{K}}\tilde{\tilde{C}}}-\tfrac{(2n-2)}{4}\delta_{\tilde{\tilde{L}}}{}^{\tilde{\tilde{M}}}\delta_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}),\\
\tilde{\Gamma}_{\tilde{\Lambda}\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}} &:= -\tfrac{2}{2n-4}\tilde{\tilde{\Gamma}}_{\tilde{\Lambda}\tilde{\tilde{C}}}{}^{\tilde{\tilde{D}}}(\varepsilon_{\tilde{\tilde{D}}\tilde{\tilde{K}}}{}^{\tilde{\tilde{C}}\tilde{\tilde{A}}}-\tfrac{2n-2}{2N^2}\delta_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}\delta_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{C}}}).
\end{aligned}
\tag{11.23}
$$

Indeed,

$$
\begin{aligned}
\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta_{\tilde{\Psi}\tilde{\tilde{K}}\tilde{\tilde{B}}}\cdot(-\tfrac{2}{2n-4})(\varepsilon_{\tilde{\tilde{D}}\tilde{\tilde{A}}}{}^{\tilde{\tilde{C}}\tilde{\tilde{K}}}-\tfrac{2n-2}{2N^2}\delta_{\tilde{\tilde{A}}}{}^{\tilde{\tilde{K}}}\delta_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{C}}}) &= \eta_{[\tilde{\Lambda}}{}^{\tilde{\tilde{B}}\tilde{\tilde{C}}}\eta_{\tilde{\Psi}]\tilde{\tilde{B}}\tilde{\tilde{D}}}-\\
-\tfrac{1}{2n-4}(\tfrac{2n}{2}-\tfrac{2n-2}{2N^2}\cdot 2N^2)g_{\tilde{\Lambda}\tilde{\Psi}}\delta_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{C}}} &= \eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{B}}\tilde{\tilde{C}}}\eta_{\tilde{\Psi}\tilde{\tilde{B}}\tilde{\tilde{D}}}.
\end{aligned}
\tag{11.24}
$$

Note 11.1. *It should be noted that for $n = 4$, all conclusions remain valid and the last identity will have the form*

$$\begin{gathered}\eta_{\tilde\Lambda}{}^{\tilde{\tilde A}\tilde{\tilde C}}\eta_{\tilde\Psi\,\tilde{\tilde K}\tilde{\tilde C}} = (\eta_{\tilde\Lambda}{}^{\tilde{\tilde C}\tilde{\tilde A}} + \eta_{\tilde\Lambda}\varepsilon^{\tilde{\tilde A}\tilde{\tilde C}})\eta_{\tilde\Psi\,\tilde{\tilde K}\tilde{\tilde C}} = \eta_{\tilde\Lambda}{}^{\tilde{\tilde C}\tilde{\tilde A}}(\eta_{\tilde\Psi\,\tilde{\tilde C}\tilde{\tilde K}} + \eta_{\tilde\Psi}\varepsilon_{\tilde{\tilde K}\tilde{\tilde C}}) + \eta_{\tilde\Lambda}\eta_{\tilde\Psi\,\tilde{\tilde K}}{}^{\tilde{\tilde A}} = \\ = \eta_{\tilde\Lambda}{}^{\tilde{\tilde C}\tilde{\tilde A}}\eta_{\tilde\Psi\,\tilde{\tilde C}\tilde{\tilde K}} - \eta_{\tilde\Psi}\eta_{\tilde\Lambda\,\tilde{\tilde K}}{}^{\tilde{\tilde A}} + \eta_{\tilde\Lambda}\eta_{\tilde\Psi\,\tilde{\tilde K}}{}^{\tilde{\tilde A}}.\end{gathered} \tag{11.25}$$

Thus, the contraction of (11.22) with $\eta_{\tilde\Psi}\eta^{\tilde\Omega}$ determines the expression $\Gamma_{\tilde\Lambda\tilde{\tilde A}}{}^{\tilde{\tilde A}}$, after substituting of which in (11.22), the coefficients $\Gamma_{\tilde\Lambda\tilde{\tilde A}}{}^{\tilde{\tilde D}}$ are uniquely determined. For this purpose, it is enough to contract (11.22) with $\eta_{\tilde\Psi\,\tilde{\tilde K}\tilde{\tilde L}}\eta^{\tilde\Omega\tilde{\tilde M}\tilde{\tilde L}}$. It is caused by the identity $\eta_{\Psi\,\tilde{\tilde A}\tilde{\tilde B}}\eta^{\Psi}{}_{\tilde{\tilde C}\tilde{\tilde D}} = \varepsilon_{\tilde{\tilde A}\tilde{\tilde B}\tilde{\tilde C}\tilde{\tilde D}} = \varepsilon_{\tilde{\tilde A}\tilde{\tilde C}}\varepsilon_{\tilde{\tilde B}\tilde{\tilde D}}$, where $\varepsilon_{\tilde{\tilde B}\tilde{\tilde D}}$, as before, is the antisymmetric metric spin-tensor (see Example 8.1 which is easily adaptable for the inclusion $\mathbb{R}^4 \subset \mathbb{C}^4$). But in this case, the condition (11.21) is equivalent to (11.16).

If we perform the symmetrization of the expression (11.22) on Ω and omitted Ψ then from the analogue of (8.17), we obtain

$$\underbrace{\frac{N^2}{2}(\partial_{\tilde\Lambda}G_{\tilde\Psi\tilde\Omega} - 2\Gamma_{\tilde\Lambda(\tilde\Omega\tilde\Psi)})}_{=\frac{N^2}{2}\nabla_\Lambda G_{\tilde\Psi\tilde\Omega}} + \underbrace{(\frac{1}{2N^2}\eta_{\tilde\Psi\,\tilde{\tilde A}\tilde{\tilde B}}\eta_{\tilde\Omega\,\tilde{\tilde C}\tilde{\tilde D}}\partial_\Lambda\varepsilon^{\tilde{\tilde A}\tilde{\tilde B}\tilde{\tilde C}\tilde{\tilde D}} + \tilde\Gamma_{\tilde\Lambda\tilde{\tilde B}}{}^{\tilde{\tilde B}}G_{\tilde\Psi\tilde\Omega})}_{=\frac{1}{2N^2}\eta_{\tilde\Psi\,\tilde{\tilde A}\tilde{\tilde B}}\eta_{\tilde\Omega\,\tilde{\tilde C}\tilde{\tilde D}}\nabla_\Lambda\varepsilon^{\tilde{\tilde A}\tilde{\tilde B}\tilde{\tilde C}\tilde{\tilde D}}} = 0. \tag{11.26}$$

If the metric tensor is covariantly constant then

$$\eta_{\tilde\Psi\,\tilde{\tilde A}\tilde{\tilde B}}\eta_{\tilde\Omega\,\tilde{\tilde C}\tilde{\tilde D}}\nabla_{\tilde\Lambda}\varepsilon^{\tilde{\tilde A}\tilde{\tilde B}\tilde{\tilde C}\tilde{\tilde D}} = 0. \tag{11.27}$$

Accordingly, the complex connection is determined by the relations

$$\nabla_\Lambda := m_\Lambda{}^{\tilde A}\nabla_{\tilde A}, \quad \bar\nabla_{\Lambda'} := \bar m_{\Lambda'}{}^{\tilde A}\nabla_{\tilde A} \tag{11.28}$$

that in the absence of torsion, will lead to the connection

$$\nabla_\Lambda g_{\Psi\Phi} = 0, \quad \bar\nabla_{\Lambda'}\bar g_{\Psi'\Phi'} = 0 \tag{11.29}$$

which prolongs into the complex spinor bundle by the conditions (see (6.33))

$$\begin{gathered}\nabla_\Lambda(\tilde M_K)_A{}^{\tilde{\tilde B}} = 0, \quad \nabla_\Lambda(\tilde{\tilde M}_K)_A{}^{\tilde{\tilde B}} = 0, \quad \bar\nabla_{\Lambda'}(\tilde M_K)_A{}^{\tilde{\tilde B}} = 0, \quad \bar\nabla_{\Lambda'}(\tilde{\tilde M}_K)_A{}^{\tilde{\tilde B}} = 0, \\ \sum_{K=1}^{2N}(\eta_K)^\Psi{}_{AB}\bar\nabla_{\Lambda'}(\eta_K)_\Omega{}^{AB} = 0, \quad \sum_{K=1}^{2N}(\eta_K)^\Psi{}_{AB}\nabla_\Lambda(\eta_K)_\Omega{}^{AB} = 0.\end{gathered} \tag{11.30}$$

It should be noted that the upper relations

$$\begin{gathered}\partial_\Lambda(\tilde M_K)_A{}^{\tilde{\tilde B}} + \tilde\Gamma_{\Lambda\tilde{\tilde C}}{}^{\tilde{\tilde B}}(\tilde M_K)_A{}^{\tilde{\tilde C}} - (\tilde\Gamma_K)_{\Lambda A}{}^C(\tilde M_K)_C{}^{\tilde{\tilde B}} = 0, \\ \partial_\Lambda(\tilde{\tilde M}_K)_A{}^{\tilde{\tilde B}} + \tilde{\tilde\Gamma}_{\Lambda\tilde{\tilde C}}{}^{\tilde{\tilde B}}(\tilde{\tilde M}_K)_A{}^{\tilde{\tilde C}} - (\tilde{\tilde\Gamma}_K)_{\Lambda A}{}^C(\tilde{\tilde M}_K)_C{}^{\tilde{\tilde B}} = 0\end{gathered} \tag{11.31}$$

are define the different coefficients of the connection: $(\tilde\Gamma_K)_{\Lambda A}{}^C$ and $(\tilde{\tilde\Gamma}_K)_{\Lambda A}{}^C$; the bottom relations are rewritten as

$$\begin{gathered}\sum_{K=1}^{2N}((\eta_K)^\Psi{}_{AB}\partial_\Lambda(\eta_K)_\Omega{}^{AB} - \tfrac{N}{2}\cdot\Gamma_{\Lambda\Omega}{}^\Psi + \\ +(\tilde\Gamma_K)_{\Lambda D}{}^A(\eta_K)^\Psi{}_{AB}(\eta_K)_\Omega{}^{DB} + (\tilde{\tilde\Gamma}_K)_{\Lambda D}{}^B(\eta_K)^\Psi{}_{AB}(\eta_K)_\Omega{}^{AD}) = 0.\end{gathered} \tag{11.32}$$

Therefore, in order to prolong uniquely the connection into the spinor bundle, it is necessary to demand

$$(\tilde{\tilde{\Gamma}}_K)_{\Lambda L}{}^M := \tfrac{2}{N}(\tilde{\tilde{\Gamma}}_K)_{\Lambda D}{}^A((\eta_K)_{\Psi LB}(\eta_K)_\Omega{}^{MB}(\eta_K)^\Psi{}_{AC}(\eta_K)^{\Omega DC} - \tfrac{(n-2)}{4}\delta_L{}^M\delta_D{}^A),$$

$$(\tilde{\Gamma}_K)_{\Lambda A}{}^C := -\tfrac{2}{n-4}(\tilde{\tilde{\Gamma}}_K)_{\Lambda K}{}^L((\varepsilon_K)_{LA}{}^{KC} - \tfrac{n-2}{N}\delta_L{}^K\delta_A{}^C). \tag{11.33}$$

Therefore, from the covariant constancy of the metric tensor

$$\begin{aligned} \nabla_\Lambda g_{\Psi\Omega} = (\tfrac{2}{N})^2\nabla_\Lambda((\eta_K)_{\Psi AB}(\eta_K)_{\Omega CD}(\varepsilon_K)^{ABCD}) = \\ = \tfrac{2}{N}((\eta_K)_\Psi{}^{AB}\nabla_\Lambda(\eta_K)_{\Omega AB} + (\eta_K)_\Omega{}^{AB}\nabla_\Lambda(\eta_K)_{\Psi AB}) + \\ +(\tfrac{2}{N})^2(\eta_K)_{\Psi AB}(\eta_K)_{\Omega CD}\nabla_\Lambda(\varepsilon_K)^{ABCD} = 0, \end{aligned} \tag{11.34}$$

the relation

$$\sum_{K=1}^{2N}(\eta_K)_{\Psi AB}(\eta_K)_{\Omega CD}\nabla_\Lambda(\varepsilon_K)^{ABCD} = 0 \tag{11.35}$$

will follow. We now turn to a real inclusion. The equation

$$\nabla_\Lambda H_i{}^\Psi = 0 \tag{11.36}$$

can be rewritten as

$$\partial_\Lambda H_i{}^\Psi - \Gamma_{\Lambda i}{}^j H_j{}^\Psi + \Gamma_{\Lambda\Phi}{}^\Psi H_i{}^\Phi = 0. \tag{11.37}$$

Accordingly, in the spinor bundle, the real connection can be prolonged by the condition

$$\sum_{I=1}^{2N}(\eta_K)^i{}_{AB}\nabla_k(\eta_K)_j{}^{AB} = 0. \tag{11.38}$$

This means that on the real space, the connection compatible with the metric takes the form

$$\nabla_i g_{ik} = 0, \tag{11.39}$$

where

$$\nabla_i := H_i{}^\Lambda\nabla_\Lambda. \tag{11.40}$$

Example 11.1. *n=4. Metric index is equal to 1.*
Let

$$g^i{}_{AB'} := (\eta_1)^i{}_{AB}\bar{S}_{B'}{}^B, \quad g_i{}^{AB'} := (\eta_1)_i{}^{AB}S_B{}^{B'}, \quad \overline{g_j{}^{A'B}} = \bar{g}_j{}^{B'A} = g_j{}^{AB'}. \tag{11.41}$$

Then the special case of the coordination for the connections can look like

$$g^i{}_{AB'}\nabla_k g_j{}^{AB'} + \bar{g}^i{}_{B'A}\nabla_k\bar{g}_j{}^{B'A} = 0. \tag{11.42}$$

The condition can be rewritten as

$$\begin{aligned} g^i{}_{AB'}(\partial_k g_j{}^{AB'} - \Gamma_{kj}{}^l g_l{}^{AB'} + \tilde{\tilde{\Gamma}}_{kC}{}^A g_j{}^{CB'} + \tilde{\Gamma}_{kC'}{}^{B'} g_j{}^{AC'} + \partial_k g_j{}^{AB'} - \Gamma_{kj}{}^l g_l{}^{AB'} + \\ +\overline{\tilde{\tilde{\Gamma}}}_{kC'}{}^{B'} g_j{}^{AC'} + \overline{\tilde{\Gamma}}_{kC}{}^A g_j{}^{CB'}) = 0. \end{aligned} \tag{11.43}$$

Then due to Note 11.1, this relation can be rewritten as

$$\partial_k g_j{}^{AB'} - \Gamma_{kj}{}^l g_l{}^{AB'} + \frac{1}{2}(\tilde{\tilde{\Gamma}}_{kC}{}^A + \bar{\tilde{\Gamma}}_{kC}{}^A) g_j{}^{CB'} + \frac{1}{2}\overline{(\tilde{\tilde{\Gamma}}_{kC}{}^B + \bar{\tilde{\Gamma}}_{kC}{}^B)} g_j{}^{AC'} = 0. \tag{11.44}$$

Define $\Gamma_{kC}{}^A := \frac{1}{2}(\tilde{\tilde{\Gamma}}_{kC}{}^A + \bar{\tilde{\Gamma}}_{kC}{}^A)$ *which will lead to the connection from [22, v. 1, p. 230 (eng)]*

$$\nabla_k g_j{}^{AB'} = 0. \tag{11.45}$$

This proves the theorem.

Theorem 11.1. *Let the pseudo-Riemannian manifold* V_{2n} $(n \geq 4)$ *be given. Let in the tangent bundle with fibers isomorphic to* $R^{2n}_{(n,n)}$, *a torsion-free connection be introduced. Then the compatibility condition of the connections between the spinor and tangent bundles can have the form*

$$\eta^{\tilde{\varPsi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{B}}} \nabla_{\tilde{\varLambda}} \eta_{\tilde{\varOmega}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}} = 0. \tag{11.46}$$

In order to prolong uniquely the constructed connection into the spinor bundle, it is necessary to demand

$$\begin{aligned} \tilde{\tilde{\Gamma}}_{\tilde{\varLambda}\tilde{\tilde{L}}}{}^{\tilde{\tilde{M}}} &:= \tfrac{1}{N^2}\tilde{\tilde{\Gamma}}_{\tilde{\varLambda}\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}(\eta_{\tilde{\varPsi}\tilde{\tilde{L}}\tilde{\tilde{B}}}\eta_{\tilde{\varOmega}}{}^{\tilde{\tilde{M}}\tilde{\tilde{B}}}\eta^{\tilde{\varPsi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{C}}}\eta^{\tilde{\varOmega}\tilde{\tilde{K}}\tilde{\tilde{C}}} - \tfrac{(2n-2)}{4}\delta_{\tilde{\tilde{L}}}{}^{\tilde{\tilde{M}}}\delta_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}), \\ \tilde{\Gamma}_{\tilde{\varLambda}\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}} &:= -\tfrac{2}{2n-4}\tilde{\tilde{\Gamma}}_{\tilde{\varLambda}\tilde{\tilde{C}}}{}^{\tilde{\tilde{D}}}(\varepsilon_{\tilde{\tilde{D}}\tilde{\tilde{K}}}{}^{\tilde{\tilde{C}}\tilde{\tilde{A}}} - \tfrac{2n-2}{2N^2}\delta_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}\delta_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{C}}}). \end{aligned} \tag{11.47}$$

In order to prolong uniquely the connection onto the complex representation $\mathbb{C}V_n$ *with fibers isomorphic* $\mathbb{C}^n$, *it is necessary to demand*

$$\nabla_{\tilde{\varLambda}} m_{\tilde{\varPsi}}{}^{\Psi} = 0, \quad \nabla_{\tilde{\varLambda}} \bar{m}_{\tilde{\varPsi}}{}^{\Psi'} = 0, \quad \nabla_{\Lambda} := m^{\tilde{\varLambda}}{}_{\Lambda}\nabla_{\tilde{\varLambda}}, \quad \bar{\nabla}_{\Lambda'} := \bar{m}^{\tilde{\varLambda}}{}_{\Lambda'}\nabla_{\tilde{\varLambda}}. \tag{11.48}$$

Then the compatibility condition of the connections will have the form

$$\begin{gathered} \nabla_{\Lambda}(\tilde{M}_K)_A{}^{\tilde{\tilde{B}}} = 0, \quad \nabla_{\Lambda}(\tilde{\tilde{M}}_K)_A{}^{\tilde{\tilde{B}}} = 0, \quad \bar{\nabla}_{\Lambda'}(\tilde{M}_K)_A{}^{\tilde{\tilde{B}}} = 0, \quad \bar{\nabla}_{\Lambda'}(\tilde{\tilde{M}}_K)_A{}^{\tilde{\tilde{B}}} = 0, \\ \sum_{K=1}^{2N} (\eta_K)^{\Psi}{}_{AB}\bar{\nabla}_{\Lambda'}(\eta_K)_{\Omega}{}^{AB} = 0, \quad \sum_{K=1}^{2N} (\eta_K)^{\Psi}{}_{AB}\nabla_{\Lambda}(\eta_K)_{\Omega}{}^{AB} = 0. \end{gathered} \tag{11.49}$$

In order to prolong uniquely the connection, we will demand

$$\begin{gathered} (\tilde{\tilde{\Gamma}}_K)_{\Lambda L}{}^M := \tfrac{2}{N}(\tilde{\tilde{\Gamma}}_K)_{\Lambda D}{}^A((\eta_K)_{\Psi LB}(\eta_K)_{\Omega}{}^{MB}(\eta_K)^{\Psi}{}_{AC}(\eta_K)^{\Omega DC} - \tfrac{(n-2)}{4}\delta_L{}^M\delta_D{}^A), \\ (\tilde{\Gamma}_K)_{\Lambda A}{}^C := -\tfrac{2}{n-4}(\tilde{\tilde{\Gamma}}_K)_{\Lambda K}{}^L((\varepsilon_K)_{LA}{}^{KC} - \tfrac{n-2}{N}\delta_L{}^K\delta_A{}^C). \end{gathered} \tag{11.50}$$

In the case of an inclusion of a real space into the complex space, it is necessary to satisfy the relations

$$\nabla_{\Lambda} H_i{}^{\Psi} = 0, \quad \bar{\nabla}_{\Lambda'}\bar{H}_i{}^{\Psi'} = 0, \quad \nabla_i := H_i{}^{\Lambda}\nabla_{\Lambda}. \tag{11.51}$$

Then the compatibility condition will have the form

$$\sum_{I=1}^{2N} (\eta_K)^i{}_{AB}\nabla_k(\eta_K)_j{}^{AB} = 0. \tag{11.52}$$

Corollary 11.1. *If in the tangent bundle to the manifold V_{2n}, a torsion-free connection compatible with the metric*

$$\nabla_{\tilde{\Lambda}} G_{\tilde{\Psi}\tilde{\Omega}} = 0 \tag{11.53}$$

is given then in the tangent bundle of the complex representation $\mathbb{C}V_n$, the connection

$$\nabla_{\Lambda} g_{\Omega\Psi} = 0, \quad \bar{\nabla}_{\Lambda'} \bar{g}_{\Omega'\Psi'} = 0 \tag{11.54}$$

is induced. For an inclusion $V_n \subset \mathbb{C}V_n$, the connection

$$\nabla_i g_{ik} = 0 \tag{11.55}$$

is induced.

We construct the operators $P^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}} := \eta^{\tilde{\Psi}}{}_{\tilde{\tilde{B}}\tilde{\tilde{A}}} X^{\tilde{\tilde{B}}}$ and $P_{\tilde{\Psi}}{}^{\tilde{\tilde{A}}} := \eta_{\tilde{\Psi}}{}^{\tilde{\tilde{B}}\tilde{\tilde{A}}} Y_{\tilde{\tilde{B}}}$ such that $X^{\tilde{\tilde{A}}} Y_{\tilde{\tilde{A}}} = 2$ and $P_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}} P_{\tilde{\Psi}\tilde{\tilde{A}}} = G_{\tilde{\Lambda}\tilde{\Psi}}$. This is always possible for $n \geq 8$. Then the compatibility condition of the connections for the real representation has the form

$$\nabla_{\tilde{\Lambda}} P_{\tilde{\Psi}}{}^{\tilde{\tilde{A}}} = 0 \tag{11.56}$$

which can be rewritten as

$$\partial_{\tilde{\Lambda}} P_{\tilde{\Psi}}{}^{\tilde{\tilde{A}}} - \Gamma_{\tilde{\Lambda}\tilde{\Psi}}{}^{\tilde{\Phi}} P_{\tilde{\Phi}}{}^{\tilde{\tilde{A}}} + \Gamma_{\tilde{\Lambda}\tilde{\tilde{C}}}{}^{\tilde{\tilde{A}}} P_{\tilde{\Psi}}{}^{\tilde{\tilde{C}}} = 0. \tag{11.57}$$

Therefore, we can uniquely prolong the connection into the spinor bundle as

$$\Gamma_{\tilde{\Lambda}\tilde{\tilde{C}}}{}^{\tilde{\tilde{A}}} := -P^{\tilde{\Psi}}{}_{\tilde{\tilde{C}}} \partial_{\tilde{\Lambda}} P_{\tilde{\Psi}}{}^{\tilde{\tilde{A}}} + \Gamma_{\tilde{\Lambda}\tilde{\Psi}}{}^{\tilde{\Phi}} P^{\tilde{\Psi}}{}_{\tilde{\tilde{C}}} P_{\tilde{\Phi}}{}^{\tilde{\tilde{A}}}. \tag{11.58}$$

For the complex representation, the operators

$$\begin{gathered}
(P_K)^{\Psi}{}_A := m_{\tilde{\Psi}}{}^{\Psi} P^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}} (\tilde{M}_K)_A{}^{\tilde{\tilde{A}}}, \quad (P_K)^{\Psi'}{}_A := \bar{m}_{\tilde{\Psi}}{}^{\Psi'} P^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}} (\tilde{M}_K)_A{}^{\tilde{\tilde{A}}}, \\
(P^*_K)_{\Psi}{}^A := m^{\tilde{\Psi}}{}_{\Psi} P_{\tilde{\Psi}}{}^{\tilde{\tilde{A}}} (\tilde{M}^*_K)^A{}_{\tilde{\tilde{A}}}, \quad (P^*_K)_{\Psi'}{}^A := \bar{m}^{\tilde{\Psi}}{}_{\Psi'} P_{\tilde{\Psi}}{}^{\tilde{\tilde{A}}} (\tilde{M}^*_K)^A{}_{\tilde{\tilde{A}}}.
\end{gathered} \tag{11.59}$$

are defined.

Theorem 11.2. *Let the pseudo-Riemannian manifold V_{2n} $(n \geq 4)$ be given. Let in the tangent bundle with fibers isomorphic to $R^{2n}_{(n,n)}$, a torsion-free connection be introduced. Then the compatibility condition of the connections between the spinor and tangent bundles can have the form*

$$\nabla_{\tilde{\Lambda}} P_{\tilde{\Psi}}{}^{\tilde{\tilde{A}}} = 0. \tag{11.60}$$

In order to prolong uniquely the constructed connection into the spinor bundle, it is necessary to demand

$$\Gamma_{\tilde{\Lambda}\tilde{\tilde{C}}}{}^{\tilde{\tilde{A}}} := -P^{\tilde{\Psi}}{}_{\tilde{\tilde{C}}} \partial_{\tilde{\Lambda}} P_{\tilde{\Psi}}{}^{\tilde{\tilde{A}}} + \Gamma_{\tilde{\Lambda}\tilde{\Psi}}{}^{\tilde{\Phi}} P^{\tilde{\Psi}}{}_{\tilde{\tilde{C}}} P_{\tilde{\Phi}}{}^{\tilde{\tilde{A}}}. \tag{11.61}$$

In order to prolong uniquely the connection onto the complex representation $\mathbb{C}V_n$ with fibers isomorphic $\mathbb{C}^n$, it is necessary to demand

$$\nabla_{\tilde{\Lambda}} m_{\tilde{\Psi}}{}^{\Psi} = 0, \quad \nabla_{\tilde{\Lambda}} \bar{m}_{\tilde{\Psi}}{}^{\Psi'} = 0, \quad \nabla_{\Lambda} := m^{\tilde{\Lambda}}{}_{\Lambda} \nabla_{\tilde{\Lambda}}, \quad \bar{\nabla}_{\Lambda'} := \bar{m}^{\tilde{\Lambda}}{}_{\Lambda'} \nabla_{\tilde{\Lambda}}. \tag{11.62}$$

Then the compatibility condition of the connections has the form

$$\begin{gathered} \nabla_{\Lambda} (\tilde{M}_K)_A{}^{\tilde{\tilde{B}}} = 0, \quad \bar{\nabla}_{\Lambda'} (\tilde{M}_K)_A{}^{\tilde{\tilde{B}}} = 0, \\ \nabla_{\Lambda} (P_K)^{\Psi}{}_A = 0, \quad \bar{\nabla}_{\Lambda'} (P_K)^{\Psi}{}_A = 0, \quad \nabla_{\Lambda} (\bar{P}_K)^{\Psi'}{}_A = 0, \quad \bar{\nabla}_{\Lambda'} (\bar{P}_K)^{\Psi'}{}_A = 0. \end{gathered} \tag{11.63}$$

In the case of an inclusion of a real space in the complex space, it is necessary to satisfy the relations

$$\nabla_{\Lambda} H_i{}^{\Psi} = 0, \quad \bar{\nabla}_{\Lambda'} \bar{H}_i{}^{\Psi'} = 0, \quad \nabla_i := H_i{}^{\Lambda} \nabla_{\Lambda}. \tag{11.64}$$

Then the compatibility condition of the connections has the form

$$\nabla_i (P_K)^j{}_A = 0, \quad \nabla_i (\bar{P}_K)^j{}_{A'} = 0. \tag{11.65}$$

Now one can construct the operators

$$\begin{aligned} (Y_K)^A &:= y^{\Psi} (P^*_K)_{\Psi}{}^A, \quad y^{\Psi} = \sum_{J=1}^{2N} (Y_J)^B (P_J)^{\Psi}{}_B, \\ (\bar{Y}_K)^A &:= \bar{y}^{\Psi'} (\bar{P}^*_K)_{\Psi'}{}^A, \quad \bar{y}^{\Psi'} = \sum_{J=1}^{2N} (\bar{Y}_J)^B (\bar{P}_J)^{\Psi'}{}_B, \\ L_x (Y_K)^A &:= x^{\Omega} \nabla_{\Omega} (Y_K)^A - \sum_{J=1}^{2N} (Y_J)^B (P_J)^{\Psi}{}_B (P^*_K)_{\Omega}{}^A \nabla_{\Psi} x^{\Omega}, \\ \bar{L}_{\bar{x}} (\bar{Y}_K)^A &:= \bar{x}^{\Omega'} \bar{\nabla}_{\Omega'} (\bar{Y}_K)^A - \sum_{J=1}^{2N} (\bar{Y}_J)^B (\bar{P}_J)^{\Psi'}{}_B (\bar{P}^*_K)_{\Omega'}{}^A \bar{\nabla}_{\Psi'} \bar{x}^{\Omega'}. \end{aligned} \tag{11.66}$$

These operators are invariants of the connection only if

$$\begin{gathered} \partial_{\Lambda} (P_K)^{\Psi}{}_B = 0, \quad \bar{\partial}_{\Lambda'} (P_K)^{\Psi}{}_B = 0, \quad \partial_{\Lambda} (P^*_K)_{\Psi}{}^B = 0, \quad \bar{\partial}_{\Lambda'} (P^*_K)_{\Psi}{}^B = 0, \\ \partial_{\Lambda} (\bar{P}_K)^{\Psi'}{}_B = 0, \quad \bar{\partial}_{\Lambda'} (\bar{P}_K)^{\Psi'}{}_B = 0, \quad \partial_{\Lambda} (\bar{P}^*_K)_{\Psi'}{}^B = 0, \quad \bar{\partial}_{\Lambda'} (\bar{P}^*_K)_{\Psi'}{}^B = 0 \end{gathered} \tag{11.67}$$

under the condition

$$\partial_{\tilde{\Lambda}} P^{\tilde{\Psi}}{}_{\tilde{\tilde{B}}} = 0, \quad \partial_{\tilde{\Lambda}} P_{\tilde{\Psi}}{}^{\tilde{\tilde{B}}} = 0, \quad \partial_{\tilde{\Lambda}} m^{\tilde{\Psi}}{}_{\Psi} = 0, \quad \partial_{\tilde{\Lambda}} m_{\tilde{\Psi}}{}^{\Psi} = 0, \quad \partial_{\tilde{\Lambda}} \bar{m}^{\tilde{\Psi}}{}_{\Psi'} = 0, \quad \partial_{\tilde{\Lambda}} \bar{m}_{\tilde{\Psi}}{}^{\Psi'} = 0. \tag{11.68}$$

Then from (11.58) and (6.33), the parities

$$(\Gamma_K)_{\Lambda C}{}^A = \Gamma_{\Lambda \Psi}{}^{\Phi} \underbrace{(P_K)^{\Psi}{}_C (P^*_K)_{\Phi}{}^A}_{:=(P_K)_{\Phi}{}^{\Psi}{}_C{}^A}, \quad (\bar{\Gamma}_K)_{\Lambda' C}{}^A = \Gamma_{\Lambda' \Psi}{}^{\Phi} \underbrace{(P_K)^{\Psi}{}_C (P^*_K)_{\Phi}{}^A}_{:=(P_K)_{\Phi}{}^{\Psi}{}_C{}^A} = 0 \tag{11.69}$$

will follow that will ensure the invariance of the operators. Then the following identities will be executed (the proof is given in Appendix)

$$\begin{aligned} L_{[x,y]} (Y_K)^A &= L_x L_y (Y_K)^A - L_y L_x (Y_K)^A, \\ \bar{L}_{[\bar{x},\bar{y}]} (\bar{Y}_K)^A &= \bar{L}_{\bar{x}} \bar{L}_{\bar{y}} (\bar{Y}_K)^A - \bar{L}_{\bar{y}} \bar{L}_{\bar{x}} (\bar{Y}_K)^A. \end{aligned} \tag{11.70}$$

For n = 4, such the connections were considered in [9]. However, in [9], the expression of the form $(\eta_K)^{\Psi}{}_{DC} (\eta_K)_{\Phi}{}^{DA}$ unlike the expression $(P_K)_{\Phi}{}^{\Psi}{}_C{}^A$ of this article was considered, and this would cause the torsion of such the operators.

Theorem 11.3. *The operators*

$$
\begin{aligned}
L_x(Y_K)^A &:= x^\Omega \partial_\Omega (Y_K)^A - \sum_{J=1}^{2N} (Y_J)^B (P_J)^\Psi{}_B (P^*_K)_\Omega{}^A \partial_\Psi x^\Omega, \\
\bar{L}_{\bar{x}}(\bar{Y}_K)^A &:= \bar{x}^{\Omega'} \bar{\partial}_{\Omega'} (\bar{Y}_K)^A - \sum_{J=1}^{2N} (\bar{Y}_J)^B (\bar{P}_J)^{\Psi'}{}_B (\bar{P}^*_K)_{\Omega'}{}^A \bar{\partial}_{\Psi'} \bar{x}^{\Omega'}
\end{aligned}
\tag{11.71}
$$

will be invariants of the complex connection from Theorem 11.2 under the condition of the constancy of the operators

$$
\begin{gathered}
\partial_\Lambda (P_K)^\Psi{}_B = 0, \quad \bar{\partial}_{\Lambda'} (P_K)^\Psi{}_B = 0, \quad \partial_\Lambda (P^*_K)_\Psi{}^B = 0, \quad \bar{\partial}_{\Lambda'} (P^*_K)_\Psi{}^B = 0, \\
\partial_\Lambda (\bar{P}_K)^{\Psi'}{}_B = 0, \quad \bar{\partial}_{\Lambda'} (\bar{P}_K)^{\Psi'}{}_B = 0, \quad \partial_\Lambda (\bar{P}^*_K)_{\Psi'}{}^B = 0, \quad \bar{\partial}_{\Lambda'} (\bar{P}^*_K)_{\Psi'}{}^B = 0, \\
\partial_{\tilde{\Lambda}} P^{\tilde{\Psi}}{}_{\tilde{\tilde{B}}} = 0, \quad \partial_{\tilde{\Lambda}} P_{\tilde{\Psi}}{}^{\tilde{\tilde{B}}} = 0, \\
\partial_{\tilde{\Lambda}} m^{\tilde{\Psi}}{}_\Psi = 0, \quad \partial_{\tilde{\Lambda}} m_{\tilde{\Psi}}{}^\Psi = 0, \quad \partial_{\tilde{\Lambda}} \bar{m}^{\tilde{\Psi}}{}_{\Psi'} = 0, \quad \partial_{\tilde{\Lambda}} \bar{m}_{\tilde{\Psi}}{}^{\Psi'} = 0, \\
\sum_{K=1}^{2N} (P_K)^\Psi{}_B (P^*_K)_\Phi{}^B = \delta_\Psi{}^\Phi, \quad \sum_{K=1}^{2N} (\bar{P}_K)^{\Psi'}{}_B (\bar{P}^*_K)_{\Phi'}{}^B = \delta_{\Psi'}{}^{\Phi'}.
\end{gathered}
\tag{11.72}
$$

These operators satisfy the identities

$$
\begin{aligned}
L_{[x,y]}(Y_K)^A &= L_x L_y (Y_K)^A - L_y L_x (Y_K)^A, \\
\bar{L}_{[\bar{x},\bar{y}]}(\bar{Y}_K)^A &= \bar{L}_{\bar{x}} L_{\bar{y}} (\bar{Y}_K)^A - \bar{L}_{\bar{y}} L_{\bar{x}} (\bar{Y}_K)^A
\end{aligned}
\tag{11.73}
$$

and are the natural prolongation of the operators $L_x(y)^\Psi = x^\Omega \partial_\Omega y^\Psi - y^\Omega \partial_\Omega x^\Psi$, $\bar{L}_{\bar{x}}(\bar{y})^{\bar{\Psi}} = \bar{x}^{\Omega'} \bar{\partial}_{\Omega'} \bar{y}^{\Psi'} - \bar{y}^{\Omega'} \bar{\partial}_{\Omega'} \bar{x}^{\Psi'}$ *into the spinor bundle*

$$
\begin{gathered}
L_x(y)^\Psi = \sum_{K=1}^{2N} L_x(Y_K)^A (P_K)^\Psi{}_A, \quad \bar{L}_{\bar{x}}(\bar{y})^{\Psi'} = \sum_{K=1}^{2N} \bar{L}_{\bar{x}}(\bar{Y}_K)^A (\bar{P}_K)^{\Psi'}{}_A, \\
L_x(Y_K)^A = L_x(y)^\Psi (P^*_K)_\Psi{}^A, \quad \bar{L}_{\bar{x}}(\bar{Y}_K)^A = \bar{L}_{\bar{x}}(\bar{y})^{\Psi'} (\bar{P}^*_K)_{\Psi'}{}^A.
\end{gathered}
\tag{11.74}
$$

12 On the classification of tensors with the symmetries of the Riemann curvature tensor. Curvature spinors

In this section, it is told about how to associate tensors possessing the symmetries of the curvature tensor to their spinor analogues. This simplifies the classification of such the tensors for small dimensions. The conclusion of all results of this section is made on the basis [22, v. 1, p. 231-246 (eng)].

After the introduction of the connection, we can pass to the study of the curvature tensor properties with the restriction $(\Gamma_K)_{\Lambda A}{}^B = (\tilde{\Gamma}_K)_{\Lambda A}{}^B = (\tilde{\tilde{\Gamma}}_K)_{\Lambda A}{}^B$ in the corresponding bundles. The curvature tensors can be calculated by the formulas

$$
\nabla_{[\Lambda} \nabla_{\Psi]} r^\Omega = R_{\Lambda\Psi\Phi}{}^\Omega r^\Phi, \quad \nabla_{[\Lambda} \nabla_{\Psi]} X^A = \sum_{K=1}^{2N} (\mathcal{R}_K)_{\Lambda\Psi C}{}^A X^C \tag{12.1}
$$

Consider the coordination of the complex connections induced by the formulas (11.30), (11.33) with the covariant constancy of the metric tensor. We require that

$$
\nabla_\Lambda (\varepsilon_K)_{AB}{}^{CD} = \partial_\Lambda (\varepsilon_K)_{AB}{}^{CD} = 0. \tag{12.2}
$$

Then

$$\sum_{K=1}^{2N} (\nabla_{\Theta}(\eta_K)^{\Psi}{}_{AB})(\nabla_{\Lambda}(\eta_K)_{\Omega}{}^{AB}) = 0. \tag{12.3}$$

Therefore, the integrability condition of (11.49) from Theorem 11.1 takes the form

$$\begin{gathered}\sum_{K=1}^{2N} (\eta_K)^{\Theta}{}_{AB}\nabla_{[\Lambda}\nabla_{\Psi]}(\eta_K)_{\Omega}{}^{AB} = 0,\\ N^2 R_{\Lambda\Psi\Omega}{}^{\Theta} = \sum_{K=1}^{2N} ((\mathcal{R}_K)_{\Lambda\Psi C}{}^{A}(\eta_K)^{\Theta}{}_{AB}(\eta_K)_{\Omega}{}^{CB} + (\mathcal{R}_K)_{\Lambda\Psi C}{}^{B}(\eta_K)^{\Theta}{}_{AB}(\eta_K)_{\Omega}{}^{AC}).\end{gathered} \tag{12.4}$$

Define

$$(A_K)_{\Theta\Phi L}{}^{X} := \frac{1}{2}(\eta_K)_{[\Theta}{}^{MX}(\eta_K)_{\Phi]ML}. \tag{12.5}$$

Then from (8.27), the identity

$$(A_K)_{\Phi\Theta L}{}^{X}(A_K)_{\Omega\Gamma X}{}^{L} = \frac{N}{8} g_{\Phi[\Gamma} g_{\Omega]\Theta} \tag{12.6}$$

will follow. Then we can set

$$\begin{gathered}(R_K)_{\Lambda\Psi C}{}^{N} := -R_{\Lambda\Psi\Theta\Phi}(A_K)^{\Theta\Phi}{}_{C}{}^{N}, \quad R_{\Lambda\Psi\Theta\Phi} = \tfrac{8}{N}(A_K)_{\Theta\Phi C}{}^{N}(R_K)_{\Lambda\Psi N}{}^{C},\\ (R_K)_{\Lambda\Psi C}{}^{M} = -\tfrac{2}{n-4}(R_K)_{\Lambda\Psi L}{}^{D}\varepsilon_{CD}{}^{ML}\end{gathered} \tag{12.7}$$

(the proof of the last equation is given in Appendix). It is obvious that $(R_K)_{\Lambda\Psi C}{}^{N} = (\mathcal{R}_K)_{\Lambda\Psi C}{}^{N}$, for example, if the coordination condition of the connections has the form

$$\nabla_{\Lambda}(\eta_K)_{\Psi}{}^{AB} = 0. \tag{12.8}$$

For further calculations, we should use the identities (the number K will be omitted since all the other calculations do not depend on this number)

$$\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_2} = \frac{N}{2} g_{\Lambda_1[\Lambda_4} g_{\Lambda_3]\Lambda_2}. \tag{12.9}$$

$$\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} = N g_{[\Lambda_3|[\Lambda_2} g_{\Lambda_1]|[\Lambda_6} g_{\Lambda_5]|\Lambda_4]}. \tag{12.10}$$

$$\begin{gathered}\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2} =\\ = \tfrac{N}{4}(g_{\Lambda_1[\Lambda_8} g_{\Lambda_7]\Lambda_2} g_{\Lambda_3[\Lambda_6} g_{\Lambda_5]\Lambda_4} - g_{\Lambda_1[\Lambda_6} g_{\Lambda_5]\Lambda_2} g_{\Lambda_4[\Lambda_8} g_{\Lambda_7]\Lambda_3} +\\ + g_{\Lambda_1[\Lambda_4} g_{\Lambda_3]\Lambda_2} g_{\Lambda_5[\Lambda_8} g_{\Lambda_7]\Lambda_6}) + N(g_{[\Lambda_3|[\Lambda_2} g_{\Lambda_1][\Lambda_8} g_{\Lambda_7][\Lambda_6} g_{\Lambda_5]|\Lambda_4]} +\\ + g_{[\Lambda_5|[\Lambda_4} g_{\Lambda_3][\Lambda_7} g_{\Lambda_8][\Lambda_2} g_{\Lambda_1]|\Lambda_6]} + g_{[\Lambda_3|[\Lambda_7} g_{\Lambda_8][\Lambda_5} g_{\Lambda_6][\Lambda_1} g_{\Lambda_2]|\Lambda_4]})\end{gathered} \tag{12.11}$$

(the proof is given in Appendix).

Theorem 12.1. *The classification of a bitensor, possessing the properties* ($\Lambda, \Psi, ... = \overline{1, n}$, $A, B, ... = \overline{1, N}$)

$$R_{\Lambda\Psi\Phi\Theta} = R_{[\Lambda\Psi][\Phi\Theta]}, \quad R_{\Lambda\Psi\Phi\Theta} = R_{\Phi\Theta\Lambda\Psi} \tag{12.12}$$

and belonging to the tangent bundle $\tau(\mathbb{C}V_n)$ *over the analytic Riemannian space* $\mathbb{C}V_n$*, can be reduced to the classification of a tensor* $R_A{}^B{}_C{}^D$ *of the N-dimensional complex spinor space such that*

$$R_C{}^D{}_S{}^R := R_{\Lambda\Psi\Phi\Theta}A^{\Lambda\Psi}{}_C{}^D A^{\Phi\Theta}{}_S{}^R, \quad A_{\Lambda\Psi C}{}^D := \frac{1}{2}\eta_{[\Lambda}{}^{AD}\eta_{\Psi]AC}. \tag{12.13}$$

Besides, the relations

$$R_K{}^K{}_S{}^R = R_S{}^R{}_K{}^K = 0, \quad R_C{}^D{}_S{}^R = R_S{}^R{}_C{}^D \tag{12.14}$$

will be executed. The decomposition

$$\begin{gathered} R_C{}^K{}_M{}^A = C_C{}^K{}_M{}^A - \tfrac{4}{N(n-2)} R_G{}^D{}_N{}^P \varepsilon^{ABKL} \varepsilon^{GN}{}_{ML} \varepsilon_{DPCB} - \\ -\big(\tfrac{1}{2(n-1)(n-2)} - \tfrac{1}{4(n-2)}\big) R \varepsilon^{ABKL} \varepsilon_{CBML} - \big(\tfrac{1}{4(n-2)} - \tfrac{n}{8(n-1)(n-2)}\big) R \delta_C{}^A \delta_M{}^K \end{gathered} \tag{12.15}$$

corresponds to the decomposition of the tensor $R_{\Lambda\Psi}{}^{\Phi\Theta}$

$$R_{\Lambda\Psi}{}^{\Phi\Theta} = C_{\Lambda\Psi}{}^{\Phi\Theta} + \frac{4}{n-2} R_{[\Lambda}{}^{[\Phi} g_{\Psi]}{}^{\Theta]} - \frac{2}{(n-1)(n-2)} R g_{[\Lambda}{}^{[\Phi} g_{\Psi]}{}^{\Theta]} \tag{12.16}$$

into the irreducible components not resulted by orthogonal transformations. At the same time, the Bianchi identity

$$R_{\Lambda\Psi\Phi\Theta} + R_{\Lambda\Theta\Psi\Phi} + R_{\Lambda\Phi\Theta\Psi} = 0 \tag{12.17}$$

will have the form

$$R_L{}^D{}_S{}^L = -\frac{1}{8} \cdot R \delta_S{}^D. \tag{12.18}$$

Proof.

1. The first step. The Bianchi identity $R_{[\Lambda\Psi\Phi]\Theta} = 0$.

$$\begin{gathered} R_{[\Lambda\Psi\Phi]\Theta} \eta^\Lambda{}_{AB} \eta^{\Psi AC} \eta^\Phi{}_{DC} \eta^{\Theta DL} = \\ = R_{\Lambda\Psi\Phi\Theta} \tfrac{1}{3} (\eta^\Lambda{}_{AB} \eta^{\Psi AC} \eta^\Phi{}_{DC} + \eta^\Phi{}_{AB} \eta^{\Lambda AC} \eta^\Psi{}_{DC} + \eta^\Psi{}_{AB} \eta^{\Phi AC} \eta^\Lambda{}_{DC}) \eta^{\Theta DL} = \\ = R_{\Lambda\Psi\Phi\Theta} \tfrac{1}{3} (\eta^\Lambda{}_{AB} \eta^{\Psi AC} \eta^\Phi{}_{DC} + g^{\Phi\Lambda} \eta^\Psi{}_{DB} - \\ - \eta^\Lambda{}_{AB} \eta^{\Phi AC} \eta^\Psi{}_{DC} + \eta^\Lambda{}_{AB} \eta^{\Psi AC} \eta^\Phi{}_{DC} - g^{\Phi\Psi} \eta^\Lambda{}_{DB}) \eta^{\Theta DL} = \\ = R_{\Lambda\Psi\Phi\Theta} (\eta^\Lambda{}_{AB} \eta^{\Psi AC} \eta^\Phi{}_{DC} + g^{\Phi\Lambda} \eta^\Psi{}_{DB}) \eta^{\Theta DL} = 4 R_B{}^C{}_C{}^L + \tfrac{1}{2} R \delta_B{}^L = 0. \end{gathered} \tag{12.19}$$

   Inversely,

$$\begin{gathered} (4 R_{A_1}{}^C{}_C{}^{A_5} + \tfrac{1}{2} R \delta_{A_1}{}^{A_5}) \eta_{[\Lambda_5|A_5 A_6|} \eta_{\Lambda_6]}{}^{A_7 A_6} \eta_{[\Lambda_7|A_7 A_8|} \eta_{\Lambda_8]}{}^{A_1 A_8} = \\ = R_{\Lambda_1\Lambda_2\Lambda_3\Lambda_4} \eta^{\Lambda_1}{}_{A_1 A_2} \eta^{\Lambda_2 A_3 A_2} \eta^{\Lambda_3}{}_{A_3 A_4} \eta^{\Lambda_4 A_5 A_4} \eta_{[\Lambda_5|A_5 A_6|} \eta_{\Lambda_6]}{}^{A_7 A_6} \eta_{[\Lambda_7|A_7 A_8|} \eta_{\Lambda_8]}{}^{A_1 A_8} + \\ + \tfrac{1}{2} R \eta_{[\Lambda_5|A_1 A_6|} \eta_{\Lambda_6]}{}^{A_7 A_6} \eta_{[\Lambda_7|A_7 A_8|} \eta_{\Lambda_8]}{}^{A_1 A_8} = \tfrac{N}{4} (R_{\Lambda_8\Lambda_7\Lambda_6\Lambda_5} - R_{\Lambda_6\Lambda_5\Lambda_7\Lambda_8} - \\ - R g_{\Lambda_5[\Lambda_8} g_{\Lambda_7]\Lambda_6}) + N(-R_{[\Lambda_8|[\Lambda_5} g_{\Lambda_6]\Lambda_7]} + R_{[\Lambda_6|[\Lambda_8\Lambda_7]|\Lambda_5]} + \\ + R_{[\Lambda_6|[\Lambda_7} g_{\Lambda_8]|\Lambda_5]}) + \tfrac{1}{2} \cdot \tfrac{N}{2} R g_{\Lambda_5[\Lambda_8} g_{\Lambda_7]\Lambda_6} = \tfrac{N}{2} (R_{\Lambda_8\Lambda_7\Lambda_6\Lambda_5} + 2 R_{\Lambda_6[\Lambda_8\Lambda_7]\Lambda_5}) = 0. \end{gathered} \tag{12.20}$$

2. Second step. The Ricci tensor $R_{\Lambda\Psi}$.
   We use the two auxiliary identities.

$$\begin{gathered} R_{\Lambda\Psi C}{}^M = -\tfrac{2}{n-4} R_{\Lambda\Psi L}{}^D \varepsilon_{CD}{}^{ML}, \\ \eta_{[\Gamma}{}^{CL} \eta_{\Omega]CD} = -\tfrac{2}{n-4} \eta_{[\Gamma}{}^{CN} \eta_{\Omega]MN} \varepsilon_{CD}{}^{ML}. \end{gathered} \tag{12.21}$$

   This can determine the identity

$$\eta_{[\Gamma|CD|} \eta_{\Omega]}{}^{CL} R_{\Lambda\Psi L}{}^D = -\eta_{[\Gamma}{}^{CN} \eta_{\Omega]MN} R_{\Lambda\Psi C}{}^M. \tag{12.22}$$

Therefor,

$$\begin{gathered}
\tfrac{4}{N}\eta_{\Lambda}{}^{AB}\eta_{\Psi CD}R_{A}{}^{C}{}_{B}{}^{D}-\tfrac{1}{4}Rg_{\Lambda\Psi}= \\
=\tfrac{1}{N}\eta_{\Lambda}{}^{AB}\eta_{\Psi CD}\eta_{\Gamma AL}\eta_{\Phi}{}^{CL}\eta_{\Omega NB}\eta_{\Theta}{}^{ND}R^{\Gamma\Phi\Omega\Theta}-\tfrac{1}{4}Rg_{\Lambda\Psi}= \\
=\tfrac{1}{N}(g_{\Phi\Psi}\eta_{\Lambda}{}^{AB}\eta_{\Gamma AL}\eta_{[\Theta}{}^{NL}\eta_{\Omega]NB}-\eta_{\Lambda}{}^{AB}\eta_{\Gamma AL}\eta_{\Psi}{}^{CL}\eta_{\Phi CD}\eta_{\Theta}{}^{ND}\eta_{\Omega NB})R^{\Gamma\Phi\Omega\Theta}- \\
-\tfrac{1}{4}Rg_{\Lambda\Psi}=\tfrac{1}{2}g_{\Phi\Psi}g_{\Lambda\Omega}g_{\Gamma\Theta}R^{\Gamma\Phi\Omega\Theta}-\tfrac{1}{N}(g_{\Gamma\Psi}\eta_{\Lambda}{}^{AB}\eta_{\Phi AD}\eta_{[\Theta}{}^{ND}\eta_{\Omega]NB}- \\
-\eta_{(\Lambda}{}^{AB}\eta_{\Psi)AL}\eta_{\Gamma}{}^{CL}\eta_{\Phi CD}\eta_{[\Theta}{}^{ND}\eta_{\Omega]NB})R^{\Gamma\Phi\Omega\Theta}-\tfrac{1}{4}Rg_{\Lambda\Psi}= \\
=-\tfrac{1}{2}R_{\Lambda\Psi}-\tfrac{1}{2}g_{\Phi\Theta}g_{\Lambda\Omega}g_{\Gamma\Psi}R^{\Gamma\Phi\Omega\Theta}+\tfrac{1}{4}g_{\Lambda\Psi}g_{\Gamma\Omega}g_{\Theta\Phi}R^{\Gamma\Phi\Omega\Theta}-\tfrac{1}{4}Rg_{\Lambda\Psi}=-R_{\Lambda\Psi}.
\end{gathered} \tag{12.23}$$

Whence,

$$P_{KL}{}^{MN}:=R_{\Lambda\Psi}\eta^{\Lambda}{}_{KL}\eta^{\Psi MN}=\frac{4}{N}\varepsilon_{KL}{}^{AB}\varepsilon^{MN}{}_{CD}R_{A}{}^{C}{}_{B}{}^{D}-\frac{1}{4}R\varepsilon_{KL}{}^{MN}. \tag{12.24}$$

Moreover,

$$\begin{gathered}
g^{[\Delta|[\Psi}R^{\Phi]|\Lambda]}\eta_{[\Delta}{}^{AB}\eta_{\Lambda]CB}\eta_{[\Psi}{}^{KL}\eta_{\Phi]ML}= \\
=g^{\Delta\Psi}R^{\Phi\Lambda}(\eta_{\Delta}{}^{AB}\eta_{\Lambda CB}-\tfrac{1}{2}g_{\Delta\Lambda}\delta_{C}{}^{A})\eta_{[\Psi}{}^{KL}\eta_{\Phi]ML}= \\
=g^{\Delta\Psi}R^{\Phi\Lambda}\eta_{\Delta}{}^{AB}\eta_{\Lambda CB}(\eta_{\Psi}{}^{KL}\eta_{\Phi ML}-\tfrac{1}{2}g_{\Psi\Phi}\delta_{M}{}^{K})= \\
=\varepsilon^{ABKL}R^{\Phi\Lambda}\eta_{\Phi ML}\eta_{\Lambda CB}-\tfrac{1}{2}R^{\Delta\Lambda}\eta_{\Delta}{}^{AB}\eta_{\Lambda CB}\delta_{M}{}^{K}= \\
=\varepsilon^{ABKL}R^{\Phi\Lambda}\eta_{\Phi ML}\eta_{\Lambda CB}-\tfrac{1}{4}R\delta_{C}{}^{A}\delta_{M}{}^{K}= \\
=-\varepsilon^{ABKL}(\tfrac{4}{N}\eta^{\Phi GN}\eta^{\Lambda}{}_{DP}R_{G}{}^{D}{}_{N}{}^{P}-\tfrac{1}{4}Rg^{\Phi\Lambda})\eta_{\Lambda CB}\eta_{\Phi ML}-\tfrac{1}{4}R\delta_{C}{}^{A}\delta_{M}{}^{K}= \\
=-\tfrac{4}{N}R_{G}{}^{D}{}_{N}{}^{P}\varepsilon^{ABKL}\varepsilon^{GN}{}_{ML}\varepsilon_{DPCB}+\tfrac{1}{4}R(\varepsilon^{ABKL}\varepsilon_{CBML}-\delta_{C}{}^{A}\delta_{M}{}^{K}).
\end{gathered} \tag{12.25}$$

3. Third step. The scalar curvature R.

$$\begin{gathered}
g^{[\Delta|[\Psi}g^{\Phi]|\Lambda]}\eta_{[\Delta}{}^{AB}\eta_{\Lambda]CB}\eta_{[\Psi}{}^{KL}\eta_{\Phi]ML}= \\
=g^{\Delta\Psi}g^{\Phi\Lambda}\eta_{\Delta}{}^{AB}\eta_{\Lambda CB}(\eta_{\Psi}{}^{KL}\eta_{\Phi ML}-\tfrac{1}{2}g_{\Psi\Phi}\delta_{M}{}^{K})=\varepsilon^{ABKL}\varepsilon_{CBML}-\tfrac{n}{4}\delta_{C}{}^{A}\delta_{M}{}^{K}.
\end{gathered} \tag{12.26}$$

Now it is necessary to gather all results and to receive the analogue of the decomposition on irreducible components not resulted by orthogonal transformations. □

Corollary 12.1.

1. *The simplicity condition of an bivector of the n-dimensional space $\mathbb{C}^n$ can be written down as*
$$p^{[\Lambda\Psi}p^{\Phi\Omega]}=0. \tag{12.27}$$
The coordinates of the bivector can be associated to the traceless complex matrix
$$p_{L}{}^{D}p_{S}{}^{L}-\frac{1}{N}(p_{L}{}^{K}p_{K}{}^{L})\delta_{S}{}^{D}=0. \tag{12.28}$$

2. *A simple bivector of the space $\mathbb{C}^n$ with the condition $p^{\Lambda\Psi}p_{\Lambda\Psi}=0$ can be associated to a nilpotent operator with index equal to 2: $p_{L}{}^{D}p_{S}{}^{L}=0$.*

Proof.
1). A bivector is simple if and only if when there is the decomposition

$$p^{\Lambda\Psi}=x^{\Lambda}y^{\Psi}-y^{\Lambda}x^{\Psi}. \tag{12.29}$$

Therefore, the conditions (12.27) are satisfied automatically. Conversely, if the conditions (12.27) are executed then they can be written down as

$$p^{\Lambda\Psi}p^{\Phi\Omega} - p^{\Lambda\Phi}p^{\Psi\Omega} + p^{\Psi\Phi}p^{\Lambda\Omega} = 0. \tag{12.30}$$

Contract this equation with the nonzero covectors t_Ω and z_Φ, that $p^{\Phi\Omega}z_\Phi t_\Omega \neq 0$

$$p^{\Lambda\Psi} = \frac{1}{p^{\Theta\Xi}z_\Theta t_\Xi}(p^{\Lambda\Phi}z_\Phi p^{\Psi\Omega}t_\Omega - p^{\Psi\Phi}z_\Phi p^{\Lambda\Omega}t_\Omega). \tag{12.31}$$

Set

$$x^\Lambda := \frac{1}{p^{\Theta\Xi}z_\Theta t_\Xi}p^{\Lambda\Phi}z_\Phi, \quad y^\Psi := \frac{1}{p^{\Theta\Xi}z_\Theta t_\Xi}p^{\Psi\Omega}t_\Omega \tag{12.32}$$

from which the simplicity condition will imply. Since the tensor $R_{\Lambda\Psi\Phi\Omega} = p_{\Lambda\Psi}p_{\Phi\Omega}$ satisfies the conditions of the classification theorem, the formula (12.28) is a direct consequence of the Bianchi identity.

2). In the condition of the first paragraph, we add the condition $p^{\Lambda\Psi}p_{\Lambda\Psi} = 0$ which takes the form $p_L{}^K p_K{}^L = 0$. From here, the existence of the nilpotent operator with index equal to 2 will follow. □

We construct the analogue of the differential Bianchi identity

$$\nabla_{[\Lambda}R_{\Psi\Phi]\Theta\Omega} = 0 \tag{12.33}$$

under the covariant constancy of the connecting operators. For this purpose, we will contract it with the expression $\eta^\Lambda{}_{AB}A^{\Psi\Phi}{}_C{}^D A^{\Theta\Omega}{}_K{}^L$

$$\begin{gathered}\eta^\Lambda{}_{AB}A^{\Psi\Phi}{}_C{}^D(\eta_\Lambda{}^{MN}A_{\Psi\Phi P}{}^Q - \eta_\Psi{}^{MN}\eta_\Lambda{}^{XQ}\eta_{\Phi XP})\nabla_{MN}R_Q{}^P{}_K{}^L = \\ = -\tfrac{N^2}{16}\nabla_{AB}R_C{}^D{}_K{}^L - \tfrac{N}{8}(\varepsilon_{YCXP}\varepsilon_{AB}{}^{XQ}\nabla^{YD} - \varepsilon^{YD}{}_{XP}\varepsilon_{AB}{}^{XQ}\nabla_{YC})R_Q{}^P{}_K{}^L = 0.\end{gathered} \tag{12.34}$$

This will lead to the differential spinor Bianchi identity

$$\nabla_{AB}R_C{}^D{}_K{}^L = \frac{4}{N}(\varepsilon^{YD}{}_{XP}\varepsilon_{AB}{}^{XQ}\nabla_{YC} - \frac{1}{2}\delta_C{}^D\varepsilon_{AB}{}^{XQ}\nabla_{XP})R_Q{}^P{}_K{}^L. \tag{12.35}$$

13 Twistor equation

In this section, it is told about how to construct and to solve the n-dimensional twistor equation, and then to investigate its properties. The conclusion of all results of this section is made on the basis [22, v. 1, pp. 352-357, v. 2, pp. 43-46, p. 463 (eng)].

We define the twistor equation as

$$\eta_{\Lambda AB}\nabla_\Psi X^A + \eta_{\Psi AB}\nabla_\Lambda X^A = \frac{2}{n}g_{\Lambda\Psi}\eta^\Phi{}_{AB}\nabla_\Phi X^A. \tag{13.1}$$

We contract it with $\eta^{\Psi CB}$

$$\begin{gathered}\eta_{\Lambda AB}\nabla^{CB}X^A + \tfrac{n}{2}\nabla_\Lambda X^C = \tfrac{2}{n}\eta_\Lambda{}^{CB}\nabla_{AB}X^A, \\ \nabla_\Lambda X^C - \eta_\Lambda{}^{CB}\nabla_{AB}X^A + \tfrac{n}{2}\nabla_\Lambda X^C = \tfrac{2}{n}\eta_\Lambda{}^{CB}\nabla_{AB}X^A,\end{gathered} \tag{13.2}$$

$$\nabla_\Lambda X^C = \frac{2}{n}\eta_\Lambda{}^{CB}\nabla_{AB}X^A. \tag{13.3}$$

The integrability condition of the equation for an arbitrary X^A have the form (the proof is given in Appendix)

$$C_{\Phi\Psi\Lambda\Delta} = 0 \tag{13.4}$$

which corresponds to a conformally flat space. Let Ω be an arbitrary scalar field. Consider the conformal rescaling of the metric

$$g_{\Lambda\Psi} \to \hat{g}_{\Lambda\Psi} = \Omega g_{\Lambda\Psi}. \tag{13.5}$$

Therefore, we can set

$$\hat{\eta}_\Lambda{}^{AB} := \eta_\Lambda{}^{AB}, \quad \hat{\eta}_{\Lambda AB} := \Omega\eta_{\Lambda AB}. \tag{13.6}$$

Let's demand the performance of

$$\nabla_\Lambda \eta_\Psi{}^{AB} = 0, \quad \nabla_\Lambda \eta_{\Psi AB} = 0, \quad \hat{\nabla}_\Lambda \hat{\eta}_\Psi{}^{AB} = 0, \quad \hat{\nabla}_\Lambda \hat{\eta}_{\Psi AB} = 0. \tag{13.7}$$

This will lead to the system

$$\begin{gathered}
\begin{cases} (\hat{\nabla}_\Lambda - \nabla_\Lambda)\eta_\Psi{}^{AB} = 0, \\ (\hat{\nabla}_\Lambda - \nabla_\Lambda)\eta_{\Psi AB} = -(\Omega^{-1}\nabla_\Lambda\Omega)\eta_{\Psi AB}, \end{cases} \\
\begin{cases} Q_{\Lambda\Psi}{}^\Theta \eta_\Theta{}^{AB} = Q_{\Lambda K}{}^A \eta_\Psi{}^{KB} + Q_{\Lambda K}{}^B \eta_\Psi{}^{AK}, \\ Q_{\Lambda\Psi}{}^\Theta \eta_{\Theta AB} = -Q_{\Lambda A}{}^K \eta_{\Psi KB} - Q_{\Lambda B}{}^K \eta_{\Psi AK} + (\Omega^{-1}\nabla_\Lambda\Omega)\eta_{\Psi AB}, \end{cases}
\end{gathered} \tag{13.8}$$

where $Q_{\Lambda\Psi}{}^\Theta$ is *strain tensor*. Then

$$\begin{gathered}
\begin{cases} Q_{\Lambda(\Psi\Omega)} = \frac{2}{N} g_{\Psi\Omega} Q_{\Lambda K}{}^K, \\ Q_{\Lambda(\Psi\Omega)} = -\frac{2}{N} g_{\Psi\Omega} Q_{\Lambda K}{}^K + (\Omega^{-1}\nabla_\Lambda\Omega) g_{\Psi\Omega}, \end{cases} \\
\Upsilon_\Lambda := \tfrac{1}{2}\Omega^{-1}\nabla_\Lambda\Omega, \quad Q_{\Lambda(\Psi\Omega)} = g_{\Psi\Omega}\Upsilon_\Lambda, \quad Q_{\Lambda K}{}^K = \tfrac{N}{2}\Upsilon_\Lambda, \\
\tilde{Q}_{\Lambda\Psi\Omega} := Q_{\Lambda[\Psi\Omega]}, \quad \tilde{Q}_{\Lambda K}{}^A := Q_{\Lambda K}{}^A - \tfrac{1}{N} Q_{\Lambda L}{}^L \delta_K{}^A.
\end{gathered} \tag{13.9}$$

Therefore,

$$\tilde{Q}_{\Lambda\Psi\Omega} = \frac{8}{N} A_{\Psi\Omega A}{}^K \tilde{Q}_{\Lambda K}{}^A. \tag{13.10}$$

If we require the preservation of the twistor equation form

$$\begin{cases} \nabla_\Lambda X^C = \frac{2}{n}\eta_\Lambda{}^{CB}\nabla_{AB}X^A, \\ \hat{\nabla}_\Lambda \hat{X}^C = \frac{2}{n}\hat{\eta}_\Lambda{}^{CB}\hat{\nabla}_{AB}\hat{X}^A, \end{cases} \quad \hat{X}^C := X^C, \tag{13.11}$$

then it would impose the condition

$$Q_{\Lambda K}{}^N = \frac{2}{n}\eta_\Lambda{}^{CN}\eta_{\Psi CB} Q^\Psi{}_K{}^B, \tag{13.12}$$

on the strain tensor. Then

$$\begin{gathered}
(\tilde{Q}_{\Lambda K}{}^{N} + \tfrac{1}{2}\Upsilon_\Lambda \delta_K{}^N)\eta_\Phi{}^{LK}\eta_{\Theta LN} = \tfrac{2}{n}\eta_\Lambda{}^{CN}\eta_{\Psi CB}\eta_\Phi{}^{LK}\eta_{\Theta LN}(\tilde{Q}^\Psi{}_K{}^B + \tfrac{1}{2}\Upsilon^\Psi\delta_K{}^B),\\
\underbrace{\tilde{Q}_{\Lambda\Phi\Theta} + \Upsilon_\Lambda g_{\Phi\Theta}}_{=Q_{\Lambda\Phi\Theta}} = \tfrac{8}{nN}\eta_\Lambda{}^{CN}\eta_{\Psi CB}\eta_\Phi{}^{LK}\eta_{\Theta LN}\tilde{Q}^\Psi{}_K{}^B + \tfrac{1}{n}(g_{\Lambda\Psi}g_{\Phi\Theta} + g_{\Lambda\Theta}g_{\Phi\Psi} - g_{\Lambda\Phi}g_{\Theta\Psi})\Upsilon^\Psi,\\
Q_{\Lambda\Phi\Theta} - \tfrac{1}{n}(\Upsilon_\Lambda g_{\Phi\Theta} + \Upsilon_\Phi g_{\Lambda\Theta} - \Upsilon_\Theta g_{\Lambda\Phi}) = \tfrac{8}{nN}\eta_\Lambda{}^{CN}\eta_{\Psi CB}\eta_\Phi{}^{LK}\eta_{\Theta LN}\tilde{Q}^\Psi{}_K{}^B,\\
Q_{\Lambda(\Phi\Theta)} - \tfrac{1}{n}\Upsilon_\Lambda g_{\Phi\Theta} = \tfrac{1}{2n}(Q^\Psi{}_{\Lambda\Psi} - Q^\Psi{}_{\Psi\Lambda})g_{\Phi\Theta},\\
Q^\Psi{}_{\Lambda\Psi} - Q^\Psi{}_{\Psi\Lambda} = 2(n-1)\Upsilon_\Lambda,\\
Q_{\Lambda[\Phi\Theta]} + \tfrac{1}{n}(-\Upsilon_\Phi g_{\Lambda\Theta} + \Upsilon_\Theta g_{\Lambda\Phi}) = \tfrac{4}{nN}\eta_\Lambda{}^{CN}\eta_{\Psi CB}\eta_{[\Phi}{}^{LK}\eta_{\Theta]LN}\eta_{[\Gamma|SK|}\eta_{\Omega]}{}^{SB}\tilde{Q}^{\Psi[\Gamma\Omega]} =\\
= \tfrac{1}{2n}(g_{\Lambda\Psi}g_{\Omega\Theta}g_{\Phi\Gamma} - g_{\Lambda\Psi}g_{\Omega\Phi}g_{\Theta\Gamma})\tilde{Q}^{\Psi[\Gamma\Omega]} + \tfrac{4}{n}g_{[\Theta|[\Lambda}g_{\Psi][\Omega}g_{\Gamma]|\Phi]}Q^{\Psi\Gamma\Omega},\\
(1-\tfrac{1}{n})Q_{\Lambda[\Phi\Theta]} + \tfrac{1}{n}(-\Upsilon_\Phi g_{\Lambda\Theta} + \Upsilon_\Theta g_{\Lambda\Phi}) =\\
= \tfrac{1}{n}(g_{\Lambda\Theta}g_{\Omega\Psi}g_{\Phi\Gamma} - g_{\Psi\Theta}g_{\Omega\Lambda}g_{\Phi\Gamma} + g_{\Phi\Psi}g_{\Omega\Lambda}g_{\Theta\Gamma} - g_{\Lambda\Phi}g_{\Omega\Psi}g_{\Theta\Gamma})Q^{\Psi[\Gamma\Omega]} =\\
= \tfrac{1}{n}((n-1)\Upsilon_\Phi g_{\Theta\Lambda} - Q_{\Theta[\Phi\Lambda]} + Q_{\Phi[\Theta\Lambda]} - (n-1)\Upsilon_\Theta g_{\Phi\Lambda}).
\end{gathered}\tag{13.13}$$

Whence, we definitively obtain

$$Q_{[\Lambda\Phi\Theta]} = 0,\ \ Q_{\Lambda[\Phi\Theta]} = \Upsilon_\Phi g_{\Lambda\Theta} - \Upsilon_\Theta g_{\Lambda\Phi},\ \ Q_{\Lambda\Phi\Theta} = \Upsilon_\Lambda g_{\Phi\Theta} + \Upsilon_\Phi g_{\Lambda\Theta} - \Upsilon_\Theta g_{\Lambda\Phi}. \tag{13.14}$$

Try to solve the twistor equation in the flat space ($R_{\Lambda\Psi K}{}^A X^K = \nabla_{[\Lambda}\nabla_{\Psi]}X^A = 0$) under the covariant constancy condition of the connecting operators ($\nabla_\Lambda \eta_\Psi{}^{CB} = 0$)

$$\begin{gathered}
\nabla_\Lambda X^C = \tfrac{2}{n}\eta_\Lambda{}^{CB}\nabla_{AB}X^A,\\
\nabla_\Psi\nabla_\Lambda X^C = \tfrac{2}{n}\eta_\Lambda{}^{CB}\nabla_{AB}(\nabla_\Psi X^A),\\
\nabla_\Psi\nabla_\Lambda X^C = \tfrac{2}{n}\eta_\Lambda{}^{CB}\nabla_{AB}(\tfrac{2}{n}\eta_\Psi{}^{AK}\nabla_{LK}X^L),\\
\nabla_{(\Psi}\nabla_{\Lambda)}X^C = \tfrac{4}{n^2}(\eta_{(\Lambda}{}^{CK}\nabla_{\Psi)} - \eta_{(\Lambda}{}^{CB}\eta_{\Psi)AB}\nabla^{AK})\nabla_{LK}X^L,\\
(1-\tfrac{2}{n})\nabla_{(\Psi}\nabla_{\Lambda)}X^C = -\tfrac{2}{n^2}g_{\Lambda\Psi}\nabla^{CK}\nabla_{LK}X^L = -\tfrac{1}{n}g_{\Lambda\Psi}\nabla^\Omega\nabla_\Omega X^C,\\
\nabla_\Psi\nabla_\Lambda X^C = 0.
\end{gathered}\tag{13.15}$$

Thus, $\nabla_\Lambda X^C$ is a constant and can be represented as

$$\nabla_\Lambda X^C := i\eta_\Lambda{}^{CA}\dot{Y}_A, \tag{13.16}$$

where as $\dot{Y}_A$, $\dot{X}^C$ denote the constant spinor fields. Integrating this equation, we obtain

$$X^C := \dot{X}^C + iR^{CA}\dot{Y}_A. \tag{13.17}$$

We are interested in the case, where $X^C = 0$. Omitting the terms above the spinors, we obtain the following relation

$$X^C = -iR^{CA}Y_A. \tag{13.18}$$

From the geometric point of view, the equation (13.16), rewritten in terms of the operators γ_Λ, is anything other as the derivational equation [16, eq. (1.2)] of the normalized Grassmannian, and (13.17) is the equation [16, eq. (2.6)]. Such the normalization is called *spinor normalization*. For n = 6, it was constructed in [7].

We return to the twistor equation (13.1), but use the covariant constancy operators (11.59) constructed in the previous section. Then for the real representation, the twistor equation (13.16) takes the form

$$\nabla_{\tilde{\Lambda}} X^{\tilde{\tilde{C}}} := i\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{C}}\tilde{\tilde{A}}}\dot{Y}_{\tilde{\tilde{A}}} =: P_{\tilde{\Lambda}}{}^{\tilde{\tilde{C}}}, \tag{13.19}$$

and for the complex representation,

$$\begin{aligned} \nabla_\Lambda (X_K)^C = i(\eta_K)_\Lambda{}^{C\tilde{\tilde{A}}} \dot{Y}_{\tilde{\tilde{A}}} =: (P^*_K)_\Lambda{}^C, \\ (P_K)_{\Lambda A} \nabla_\Psi (X_K)^A + (P_K)_{\Psi A} \nabla_\Lambda (X_K)^A = \tfrac{2}{n} g_{\Lambda\Psi} (P_K)^\Phi{}_A \nabla_\Phi (X_K)^A. \end{aligned} \tag{13.20}$$

Carrying out the summation on K and defining $x_\Lambda := \sum\limits_{K=1}^{2N} (P_K)_{\Lambda A}(X_K)^A$, we obtain the conformal Killing equation

$$\nabla_\Lambda x_\Psi + \nabla_\Psi x_\Lambda = \frac{2}{n} g_{\Lambda\Psi} \nabla_\Phi x^\Phi.$$

14 Spinor formalism for n=6 and n=8

In this section, it is told about how to construct the spinor formalism for the small dimensions. For clarity, we construct various geometric interpretations of some algebraic relations. Presented, how to pass to the structure constants of the octonion algebra with the help of the Cartan triality principle. The conclusion of all results of this section is made on the basis [1] and the literature to it, [11], [12], [13], [24], [25], [26], [28].

14.1 General isomorphisms

In this subsection, it is told about how to construct the basic isomorphisms of the spinor formalism for n = 6. The conclusion of all results of this subsection is made on the basis [1], [3].

1. $\mathbb{C}^6 \cong \Lambda^2 \mathbb{C}^4$.
   Let $\alpha, \beta, ... = \overline{1,6}$, $a, b, a_1, b_1, ... = \overline{1,4}$. Then from (8.17), the identities

   $$\frac{1}{2} \eta^\alpha{}_{aa_1} \eta_\beta{}^{aa_1} = \delta_\alpha{}^\beta, \quad \eta^\alpha{}_{aa_1} \eta_\alpha{}^{bb_1} = \delta_{aa_1}{}^{bb_1} := 2\delta_{[a}{}^{[b} \delta_{a_1]}{}^{b_1]}, \tag{14.1}$$

   $$r^\alpha = 1/2 \cdot \eta^\alpha{}_{aa_1} R^{aa_1}, \quad R^{aa_1} = \eta_\alpha{}^{aa_1} r^\alpha, \tag{14.2}$$

   $$\begin{aligned} g^{\alpha\beta} = 1/4 \cdot \eta^\alpha{}_{aa_1} \eta^\beta{}_{bb_1} \varepsilon^{aa_1bb_1}, \quad \varepsilon^{aa_1bb_1} = \eta_\alpha{}^{aa_1} \eta_\beta{}^{bb_1} g^{\alpha\beta}, \\ g_{\alpha\beta} = 1/4 \cdot \eta_\alpha{}^{aa_1} \eta_\beta{}^{bb_1} \varepsilon_{aa_1bb_1}, \quad \varepsilon_{aa_1bb_1} = \eta^\alpha{}_{aa_1} \eta^\beta{}_{bb_1} g_{\alpha\beta} \end{aligned} \tag{14.3}$$

   will follow. This defines the isomorphism between the space $\mathbb{C}^6$ and the bivector space $\Lambda^2\mathbb{C}^4$, where $g_{\alpha\beta}$ is the metric tensor of the space $\mathbb{C}^6$.

2. $SO(6, \mathbb{C}) \cong SL(4, \mathbb{C})/\pm 1$.
   According to (6.42), the special orthogonal transformation can be represented as

   $$S_\alpha{}^\beta \eta_\beta{}^{aa_1} = \eta_\alpha{}^{bb_1} S_b{}^a S_{b_1}{}^{a_1}. \tag{14.4}$$

   In this case, $\tilde{E}_a{}^b$ from (8.19′) is the identity transformation multiplied by the imaginary unit. For non-special transformations, we have the identity

   $$S_\alpha{}^\beta \eta_\beta{}^{aa_1} = \eta_{\alpha bb_1} S^{ba} S^{b_1a_1}. \tag{14.5}$$

3. $so(6,\mathbb{C}) \cong sl(4,\mathbb{C})$.
   Define

$$A_{\alpha\beta d}{}^{c} = \eta_{[\alpha}{}^{ca}\eta_{\beta]da}. \tag{14.6}$$

Then by analogy with (10.6) and (12.5), we obtain

$$T_{\alpha\beta} = A_{\alpha\beta d}{}^{c}T_{c}{}^{d}, \quad T_{k}{}^{k} = 0, \quad T_{\alpha\beta} = -T_{\alpha\beta}, \tag{14.7}$$

$$\begin{gathered}
A_{\alpha\beta d}{}^{c}A^{\alpha\beta}{}_{r}{}^{s} = \tfrac{1}{2}\delta_{r}{}^{s}\delta_{d}{}^{c} - 2\delta_{r}{}^{c}\delta_{d}{}^{s}, \quad A_{\alpha\beta d}{}^{c}A^{\lambda\mu}{}_{c}{}^{d} = 2\delta_{[\alpha}{}^{\mu}\delta_{\beta]}{}^{\lambda}, \\
A_{\alpha\beta d}{}^{c}A_{\gamma}{}^{\beta}{}_{r}{}^{s} = \\
= (\eta_{\alpha}{}^{cs}\eta_{\gamma rd} + \eta_{\alpha}{}^{ck}\eta_{\gamma kr}\delta_{d}{}^{s}) + 1/2(\eta_{\alpha}{}^{sn}\eta_{\gamma rn}\delta_{d}{}^{c} + \eta_{\alpha}{}^{ck}\eta_{\gamma dk}\delta_{r}{}^{s}) - 1/4g_{\alpha\gamma}\delta_{r}{}^{s}\delta_{d}{}^{c}, \\
T_{m}{}^{n} = \tfrac{1}{2}A^{\alpha\beta}{}_{m}{}^{n}T_{\beta\alpha}, \quad T_{\beta\alpha} = -T_{\alpha\beta}.
\end{gathered} \tag{14.8}$$

In addition, the main identity for the 4-vector

$$\begin{gathered}
e_{\alpha\beta\gamma\delta} = e_{[\alpha\beta\gamma\delta]}, \quad e_{\alpha\beta\gamma\delta} = A_{\alpha\beta b}{}^{a}A_{\gamma\delta d}{}^{c}e_{a}{}^{b}{}_{c}{}^{d}, \\
e_{m}{}^{n}{}_{r}{}^{s} = (e_{m}{}^{s}\delta_{r}{}^{n} + e_{r}{}^{n}\delta_{m}{}^{s}) - \tfrac{1}{2}(e_{r}{}^{s}\delta_{m}{}^{n} + e_{m}{}^{n}\delta_{r}{}^{s}), \\
e_{a}{}^{b} := \tfrac{1}{3}e_{a}{}^{k}{}_{k}{}^{b}, \quad B_{\alpha\beta\gamma\delta r}{}^{k} := A_{\alpha\beta r}{}^{d}A_{\gamma\delta d}{}^{k} + A_{\alpha\beta c}{}^{k}A_{\gamma\delta r}{}^{c}, \quad e_{\alpha\beta\gamma\delta} := B_{\alpha\beta\gamma\delta r}{}^{k}e_{k}{}^{r}, \quad e_{k}{}^{k} = 0
\end{gathered} \tag{14.9}$$

and the main identity for the 6-vector

$$\begin{gathered}
e_{\alpha\beta\gamma\delta\lambda\mu} = A_{\alpha\beta b}{}^{a}A_{\gamma\delta d}{}^{c}A_{\lambda\mu l}{}^{k}e_{a}{}^{b}{}_{c}{}^{d}{}_{k}{}^{l}, \\
e_{a}{}^{b}{}_{c}{}^{d}{}_{k}{}^{l} = \tfrac{i}{8}(2((4\delta_{k}{}^{b}\delta_{c}{}^{l} - \delta_{k}{}^{l}\delta_{c}{}^{b})\delta_{a}{}^{d} + (4\delta_{k}{}^{d}\delta_{a}{}^{l} - \delta_{k}{}^{l}\delta_{a}{}^{d})\delta_{c}{}^{b}) - \\
-(4\delta_{k}{}^{b}\delta_{a}{}^{l} - \delta_{k}{}^{l}\delta_{a}{}^{b})\delta_{c}{}^{d} - (4\delta_{k}{}^{d}\delta_{c}{}^{l} - \delta_{k}{}^{l}\delta_{c}{}^{d})\delta_{a}{}^{b}), \\
e_{\alpha\gamma\lambda\nu\pi\sigma} = 2\eta_{[\alpha}{}^{bb_1}\eta_{\gamma]dd_1}\eta_{[\lambda}{}^{mm_1}\eta_{\nu]xx_1}\eta_{[\pi}{}^{rr_1}\eta_{\sigma]}{}^{ss_1} \cdot i\ \delta_{r}{}^{d}\delta_{m}{}^{s}\delta_{b}{}^{x}\delta_{b_1}{}^{d_1}\delta_{m_1}{}^{x_1}\delta_{r_1}{}^{s_1} = \\
= \tfrac{1}{4}\eta_{[\alpha}{}^{bb_1}\eta_{\gamma]}{}^{dd_1}\eta_{[\lambda}{}^{mm_1}\eta_{\nu]}{}^{xx_1}\eta_{[\pi}{}^{rr_1}\eta_{\sigma]}{}^{ss_1} \cdot i\ \varepsilon_{rb_1dd_1}\varepsilon_{mr_1ss_1}\varepsilon_{bm_1xx_1}.
\end{gathered} \tag{14.10}$$

are executed. The proof of these identities is given in Appendix.

14.2 On the classification of tensors with the symmetries of the Riemann curvature tensor for n=6. Curvature spinors

In this subsection, it is told about how to associate tensors possessing the symmetries of the curvature tensor to their spinor analogues. This simplifies the classification of such the tensors for n=6. The conclusion of all results of this subsection is made on the basis [2].

Theorem 14.1. *The classification of a bitensor, possessing the properties*

$$R_{\alpha\beta\gamma\delta} = R_{[\alpha\beta][\gamma\delta]}, \quad R_{\alpha\beta\gamma\delta} = R_{\gamma\delta\alpha\beta}, \quad R_{\alpha\beta\gamma\delta} + R_{\alpha\delta\beta\gamma} + R_{\alpha\gamma\delta\beta} = 0 \tag{14.11}$$

and belonging to the tangent bundle $\tau(\mathbb{C}V_6)$ over the six-dimensional analytic Riemannian space $\mathbb{C}V_6$, can be reduced to the classification of a tensor $R_{a}{}^{b}{}_{c}{}^{d}$ of the 4-dimensional complex vector space

$$R_{\alpha\beta\gamma\delta} = A_{\alpha\beta d}{}^{c}A_{\gamma\delta r}{}^{s}R_{c}{}^{d}{}_{s}{}^{r}. \tag{14.12}$$

In addition, the following relations

$$R_{k}{}^{k}{}_{s}{}^{r} = R_{s}{}^{r}{}_{k}{}^{k} = 0, \quad R_{c}{}^{d}{}_{s}{}^{r} = R_{s}{}^{r}{}_{c}{}^{d} \tag{14.13}$$

are executed. The decomposition

$$R_c{}^d{}_s{}^r = C_c{}^d{}_s{}^r - P_{cs}{}^{dr} - \frac{1}{40} \cdot R(3\delta_s{}^d\delta_c{}^r - 2\delta_s{}^r\delta_c{}^d) \tag{14.14}$$

corresponds to the decomposition

$$R_{\alpha\beta}{}^{\gamma\delta} = C_{\alpha\beta}{}^{\gamma\delta} + R_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]} - 1/10 R g_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]} \tag{14.15}$$

into irreducible components not resulted by an orthogonal transformations. These components will satisfy the following relations

$$P_{cs}{}^{rd} = -4(R_{[c}{}^{[r}{}_{s]}{}^{d]} + R_k{}^{[r}{}_{[c}{}^{|k|}\delta_{s]}{}^{d]}), \tag{14.16}$$

$$C_c{}^d{}_s{}^r = R_{(c}{}^{(d}{}_{s)}{}^{r)} + \frac{1}{40} \cdot R\delta_{(s}{}^d\delta_{c)}{}^r, \quad C_c{}^d{}_s{}^r = C_{(c}{}^{(d}{}_{s)}{}^{r)}, \tag{14.17}$$

$$R = R_\beta{}^\beta = -2 \cdot R_k{}^r{}_r{}^k, \quad P_{kc}{}^{kd} = 1/2 \cdot R\delta_c{}^d, \tag{14.18}$$

$$R_l{}^d{}_s{}^l = -\frac{1}{8} \cdot R\delta_s{}^d \tag{14.19}$$

the last of which is equivalent of the Bianchi identity.

Proof. We have the following equality

$$R_{\alpha\beta\gamma\delta} = 1/16 \cdot \eta_\alpha{}^{aa_1}\eta_\beta{}^{bb_1}\eta_\gamma{}^{cc_1}\eta_\delta{}^{dd_1}R_{aa_1bb_1cc_1dd_1}. \tag{14.20}$$

Set

$$R_c{}^d{}_s{}^r := \frac{1}{4}R_{ck}{}^{dk}{}_{st}{}^{rt}, \quad R_{\beta\gamma} = \frac{1}{4}\eta_\beta{}^{cs}\eta_{\gamma rd} \cdot P_{cs}{}^{rd}. \tag{14.21}$$

From this, the parties

$$R_{\alpha\beta\gamma\delta} = A_{\alpha\beta d}{}^c A_{\gamma\delta r}{}^s R_c{}^d{}_s{}^r, \tag{14.22}$$

$$\begin{aligned} R_{\beta\delta} = R_{\alpha\beta}{}^\alpha{}_\delta = A_{\alpha\beta d}{}^c A^\alpha{}_{\delta r}{}^s R_c{}^d{}_s{}^r = (\eta_\beta{}^{cs}\eta_{\delta rd} + \eta_\beta{}^{ck}\eta_{\delta kr}\delta_d{}^s)R_c{}^d{}_s{}^r = \\ = \tfrac{1}{4}\eta_\beta{}^{cs}\eta_{\delta rd} \cdot 4(R_{[c}{}^{[d}{}_{s]}{}^{r]} - R_{[c}{}^k{}_{|k|}{}^{[r}\delta_{s]}{}^{d]}) \end{aligned} \tag{14.23}$$

will follow. Set

$$P_{cs}{}^{rd} := -4(R_{[c}{}^{[r}{}_{s]}{}^{d]} + R_{[c}{}^k{}_{|k|}{}^{[r}\delta_{s]}{}^{d]}) \tag{14.24}$$

then

$$R_{\beta\delta} = \frac{1}{4}\eta_\beta{}^{cs}\eta_{\delta rd}P_{cs}{}^{rd}. \tag{14.25}$$

Therefore, the scalar curvature has the form

$$R = R_\beta{}^\beta = \tfrac{1}{4}\eta_\beta{}^{cc_1}\eta^\beta{}_{aa_1}P_{cc_1}{}^{aa_1} = \tfrac{1}{4}\varepsilon_{aa_1}{}^{cc_1}P_{cc_1}{}^{aa_1} = \tfrac{1}{2}P_{aa_1}{}^{aa_1} = -2R_k{}^r{}_r{}^k, \tag{14.26}$$

and the condition

$$\begin{aligned} P_{ks}{}^{kd} = -4(R_{[k}{}^{[k}{}_{s]}{}^{r]} + R_{[r}{}^k{}_{|k|}{}^{[r}\delta_{s]}{}^{d]}) = \\ = -4(-\tfrac{1}{2}R_k{}^d{}_s{}^k + \tfrac{1}{4}(R_k{}^r{}_r{}^k\delta_s{}^d + 4R_k{}^d{}_s{}^k - 2R_k{}^d{}_s{}^k)) = -R_k{}^r{}_r{}^k\delta_s{}^d = \tfrac{1}{2}R\delta_s{}^d \end{aligned} \tag{14.27}$$

is satisfied. The Bianchi identity can be rewritten as

$$(A_{\alpha\beta d}{}^c A_{\gamma\delta r}{}^s + A_{\alpha\gamma d}{}^c A_{\delta\beta r}{}^s + A_{\alpha\delta d}{}^c A_{\beta\gamma r}{}^s) \cdot R_c{}^d{}_s{}^r = 0. \tag{14.28}$$

Contracting this equation with $A^{\alpha\beta}{}_{t}{}^{l}A^{\gamma\delta}{}_{m}{}^{n}$, we find that

$$4R_{k}{}^{l}{}_{m}{}^{k}\delta_{t}{}^{n}+4R_{r}{}^{n}{}_{t}{}^{r}\delta_{m}{}^{l}-2R_{k}{}^{l}{}_{t}{}^{k}\delta_{m}{}^{n}-2R_{k}{}^{n}{}_{m}{}^{k}\delta_{t}{}^{l}-2R_{k}{}^{r}{}_{r}{}^{k}\delta_{t}{}^{n}\delta_{m}{}^{l}+R_{r}{}^{k}{}_{k}{}^{r}\delta_{m}{}^{n}\delta_{t}{}^{l}=0. \tag{14.29}$$

The contraction of this equation with $\delta_{n}{}^{t}$ leads us to the spinor analog of the Bianchi identity. In this case, all 15 significant equations reserved. Let

$$C_{\alpha\beta}{}^{\gamma\delta}:=A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}C_{c}{}^{d}{}_{s}{}^{r}, \tag{14.30}$$

then

$$C_{\alpha\beta}{}^{\gamma\delta}:=R_{\alpha\beta}{}^{\gamma\delta}-R_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]}+1/10Rg_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]}, \tag{14.31}$$

$$R_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]}=A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\cdot\frac{1}{4}(P_{sc}{}^{dr}-1/2R\delta_{s}{}^{d}\delta_{c}{}^{r}+\frac{1}{4}R\delta_{s}{}^{r}\delta_{c}{}^{d}), \tag{14.32}$$

$$g_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]}=A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\cdot\frac{1}{4}(1/2\delta_{s}{}^{r}\delta_{c}{}^{d}-2\delta_{s}{}^{d}\delta_{c}{}^{r}) \tag{14.33}$$

from which we obtain the decomposition into irreducible components not resulted by orthogonal transformations. All calculations are given in Appendix. □

Note 14.1. *At n=6, there are the two classification schemes of the Weyl spinor:*

$$1.\quad C_{k}{}^{l}{}_{m}{}^{n}\phi_{l}{}^{k}=\lambda\phi_{t}{}^{n},\qquad 2.\quad C_{k}{}^{l}{}_{m}{}^{n}\phi^{km}=\lambda\phi^{ln}. \tag{14.34}$$

Corollary 14.1. *1. The simplicity conditions of a bivector of the 6-dimensional space $\mathbb{C}^6$ can be written down as*

$$p_{[\alpha\beta}p_{\gamma\beta]}=0. \tag{14.35}$$

The coordinates of the bivector can be associated to the traceless complex matrix 4×4

$$p_{l}{}^{d}p_{s}{}^{l}-1/4(p_{l}{}^{k}p_{k}{}^{l})\delta_{s}{}^{d}=0. \tag{14.36}$$

2. A simple bivector of the space $\mathbb{C}^6$, constructed on isotropic vectors ($p^{\alpha\beta}p_{\alpha\beta}=0$), can be associated to a degenerate Rosenfeld null-pair: a covector and a vector of the space $\mathbb{C}^4$, the contraction of which is zero. In this case, the vector and the covector are determined to within a complex factor.

Proof.
The proof is similar to the one for Corollary 12.1. The only difference lies in the fact that isotropic vectors r^α of the space $\mathbb{C}^6$ are represented as $r^\alpha\eta_\alpha{}^{ab}=R^{ab}=X^aY^b-X^bY^a$. Such the representation is possible due to the perform of the identities for the isotropic vector r^α ($r^\alpha r_\alpha=0$)

$$24R^{[ab}R^{cd]}=\varepsilon_{klmn}R^{kl}R^{mn}\varepsilon^{abcd}=4(r^\alpha r_\alpha)\varepsilon^{abcd}=0, \tag{14.37}$$

where ε_{abcd} is a spin-tensor of the form (8.17). Moreover, the spin-tensor is antisymmetric on all 4 indices and satisfies to the parities

$$\begin{gathered}\varepsilon_{abcd}\varepsilon^{klmn}=24\delta_{[a}{}^{k}\delta_{b}{}^{l}\delta_{c}{}^{m}\delta_{n]}{}^{d},\quad \varepsilon_{abcd}\varepsilon^{almn}=6\delta_{[b}{}^{l}\delta_{c}{}^{m}\delta_{n]}{}^{d},\quad \varepsilon_{abcd}\varepsilon^{abmn}=4\delta_{[c}{}^{m}\delta_{n]}{}^{d},\\ \varepsilon_{abcd}\varepsilon^{abcn}=6\delta_{d}{}^{n},\quad \varepsilon_{abcd}\varepsilon^{abcd}=24.\end{gathered} \tag{14.38}$$

Set $x^{\alpha}\eta_{\alpha}{}^{ab} := X^aY^b - Y^aX^b$, $y^{\alpha}\eta_{\alpha}{}^{ab} := Z^aT^b - T^aZ^b$. From the condition $p^{\alpha\beta}p_{\alpha\beta} = 0$, the relation $x^{\alpha}y_{\alpha} = 0$ will follow. This means that $\varepsilon_{abcd}X^aY^bZ^cT^d = 0$. Therefore, the vectors X^a, Y^b, Z^c, T^d are linearly dependent. Set $T^a := \alpha X^a + \beta Y^a + \gamma Z^a$ and obtain

$$\begin{gathered}
p_a{}^b := \tfrac{1}{2}A_{\alpha\beta a}{}^b p_{\alpha\beta} = \tfrac{1}{4}\eta_{\alpha}{}^{kb}\eta_{\beta}{}^{cd}\varepsilon_{cdka}(x^{\alpha}y^{\beta} - y^{\alpha}x^{\beta}) = \\
= \tfrac{1}{4}((X^kY^b - X^bY^k)Z^cT^d - X^cY^d(Z^kT^b - T^kZ^b))\varepsilon_{cdka} = \\
= Z^c(\beta Y^d)X^k\varepsilon_{cdka}Y^b - Z^c(\alpha X^d)Y^k\varepsilon_{cdka}X^b - X^cY^dZ^k\varepsilon_{cdka}(\alpha X^b + \beta Y^b + \gamma Z^b) + \\
+ X^cY^d(\gamma Z^k)\varepsilon_{cdka}Z^b = \underbrace{X^cY^dZ^k\varepsilon_{cdka}}_{:=M_a}\underbrace{(-2\alpha X^b - 2\beta Y^b)}_{:=N^b} = M_aN^b.
\end{gathered} \tag{14.39}$$

In this case, N^b and M_a are defined up to transformations

$$N^b \longmapsto e^{\phi}N^b, \quad M_a \longmapsto e^{-\phi}M_a. \tag{14.40}$$

□

Note that the pair (N^b, M_a) is *Rosenfeld null-pair* [19]. In the space $\mathbb{CP}^4 = {}'\mathbb{C}^4/{}'\mathbb{C}$ (where ${}'\mathbb{C}^s = \mathbb{C}^s/0$), N^b defines the point and M_a defines the plane with *incidence condition* $M^bN_a = 0$. Therefore, we can construct the space $\mathbb{C}\Pi^4 =' \mathbb{C}^{*4}/{}'\mathbb{C}$ which is *dual space* to $\mathbb{CP}^4$. Then the space $\mathbb{CP}^4 \times \mathbb{C}\Pi^4$ is *Rosenfeld null-pair space.* It should be noted that such the spaces studied for the first time in [23] and [14].

Now we substitute $\varepsilon_{ab}{}^{cd} = 2\delta_{[a}{}^{c}\delta_{b]}{}^{d}$ into (12.35). Then the differential Bianchi identity will become simpler to

$$\nabla_{[cm}R_{t]}{}^{k}{}_{r}{}^{s} = \delta_{[m}{}^{k}\nabla_{c|n|}R_{t]}{}^{n}{}_{r}{}^{s}. \tag{14.41}$$

Contract this equation with $\delta_k{}^c$

$$\nabla_{c(m}R_{t)}{}^{c}{}_{r}{}^{s} = 0 \tag{14.42}$$

then contract with $\delta_s{}^m$

$$\nabla_{cm}R_t{}^{c}{}_{r}{}^{m} = 1/8\nabla_{rt}R \tag{14.43}$$

which is the spinor analogue of the equation

$$\nabla^{\alpha}(R_{\alpha\beta} - 1/2Rg_{\alpha\beta}) = 0. \tag{14.44}$$

14.3 Geometric representation of a twistor in $\mathbb{R}^6_{(2,4)}$

In this subsection, it is told about how to construct a geometric interpretation of isotropic spinors (twistors) on the isotropic cone $\mathbb{R}^6_{(2,4)}$. This interpretation is similar to the spinor representation in $\mathbb{R}^4_{(1,3)}$ with that only a difference that the dimension of the flagpole and the flag-plane is incremented. The conclusion of all results of this subsection is made on the basis [5], [32].

Let the metric of the space $\mathbb{R}^6_{(2,4)}$ have the form

$$dS^2 = dT^2 + dV^2 - dW^2 - dX^2 - dY^2 - dZ^2 \tag{14.45}$$

and let the cross section of the light-cone K_6

$$T^2 + V^2 - W^2 - X^2 - Y^2 - Z^2 = 0 \tag{14.46}$$

be set by the plane V+W=1. Let's consider the stereographic projection of this section on the plane (V=0,W=1) with the pole $N(0, \frac{1}{2}, \frac{1}{2}, 0, 0, 0)$ so that the point P(T,V,W,X,Y,Z) corresponds to $p(t, 0, 1, x, y, z)$ in the plane (V=0,W=1). Then

$$T/t = X/x = Y/y = Z/z = -\frac{(V - \frac{1}{2})}{\frac{1}{2}}. \tag{14.47}$$

We make the substitution

$$\varsigma = ix - y, \quad \omega = i(t + z), \quad \eta = i(t - z) \tag{14.48}$$

and obtain

$$\varsigma = \frac{-iX + Y}{2V - 1}, \quad \eta = \frac{-i(T + Z)}{2V - 1}, \quad \omega = \frac{i(Z - T)}{2V - 1}. \tag{14.49}$$

Therefore, the metric, induced in the cross-section, has the form

$$ds^2 := dT^2 - dX^2 - dY^2 - dZ^2 = -\frac{d\varsigma d\bar{\varsigma} + d\omega d\eta}{(\varsigma\bar{\varsigma} + \eta\omega)^2}. \tag{14.50}$$

The proof of this fact is given in Appendix. Set

$$X := \begin{pmatrix} \omega & \varsigma \\ -\bar{\varsigma} & \eta \end{pmatrix}, \quad dX := \begin{pmatrix} d\omega & d\varsigma \\ -d\bar{\varsigma} & d\eta \end{pmatrix}, \quad \frac{\partial}{\partial X} := \begin{pmatrix} \frac{\partial}{\partial\omega} & \frac{\partial}{\partial\varsigma} \\ -\frac{\partial}{\partial\bar{\varsigma}} & \frac{\partial}{\partial\eta} \end{pmatrix}. \tag{14.51}$$

Then

$$ds^2 = -\frac{det(dX)}{(det(X))^2}, \quad \bar{X}^T + X = 0. \tag{14.52}$$

Consider the linear-fractional group L

$$\tilde{X} = (AX + B)(CX + D)^{-1}, \quad S := \begin{pmatrix} A & B \\ C & D \end{pmatrix}, \quad detS = 1. \tag{14.53}$$

The reality condition, imposed on X ($X^* + X = 0$), gives the linear-fractional unitary subgroup $LU(2,2)$ so that the matrix S satisfies (see Appendix) the relation

$$S^*\hat{E}S = \hat{E}, \quad \hat{E} := \begin{pmatrix} 0 & E \\ E & 0 \end{pmatrix}. \tag{14.54}$$

Further, we define the special basis [22, v. 2, p. 65, p. 305 (eng)]

$$\begin{array}{rcccl} R^{12} & = & 1/\sqrt{2}(V + W) & = & \omega^0\xi^1 - \omega^1\xi^0, \\ R^{14} & = & i/\sqrt{2}(T + Z) & = & \omega^0\bar{\eta}^{0'} - \xi^0\bar{\pi}^{0'}, \\ R^{34} & = & 1/\sqrt{2}(V - W) & = & \bar{\pi}^{0'}\bar{\eta}^{1'} - \bar{\pi}^{1'}\bar{\eta}^{0'}, \\ R^{23} & = & i/\sqrt{2}(Z - T) & = & \xi^1\bar{\pi}^{1'} - \omega^1\bar{\eta}^{1'}, \\ R^{24} & = & 1/\sqrt{2}(Y + iX) & = & \omega^1\bar{\eta}^{0'} - \xi^1\bar{\pi}^{0'}, \\ R^{13} & = & 1/\sqrt{2}(Y - iX) & = & \xi^0\bar{\pi}^{1'} - \omega^0\bar{\eta}^{1'}. \end{array} \tag{14.55}$$

This relations are remarkable because this shows as the bivector R^{ab} is expressed in terms of its spinor components. Set

$$X := YZ^{-1},\quad \tilde{Y} = AY + BZ,\quad \tilde{Z} = CY + DZ,\quad Y = \begin{pmatrix} \omega^0 & \xi^0 \\ \omega^1 & \xi^1 \end{pmatrix},\quad Z = \begin{pmatrix} \bar{\pi}_0 & \bar{\eta}_0 \\ \bar{\pi}_1 & \bar{\eta}_1 \end{pmatrix}, \tag{14.56}$$

then

$$R := \parallel R^{ab} \parallel = \begin{pmatrix} (detY)J & YZ^{-1}(detZ)J \\ -(YZ^{-1}(detZ)J)^T & (detZ)J \end{pmatrix} = \begin{pmatrix} Y \\ Z \end{pmatrix} J \begin{pmatrix} Y^T & Z^T \end{pmatrix}, \\ J := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},\quad \tilde{R} = SRS^T. \tag{14.57}$$

Determine

$$\tilde{S} := \tilde{I}S\tilde{I}^{-1},\quad \tilde{I} := \frac{1}{\sqrt{2}}\begin{pmatrix} E & E \\ -E & E \end{pmatrix}, \tag{14.58}$$

then

$$\tilde{S}^*\tilde{E}\tilde{S} = \tilde{E}. \tag{14.59}$$

The matrixes S form the group isomorphic $SU(2,2)$ so the matrixes $\tilde{S}$ form the group $SU(2,2)$. A transformation from the group $LU(2,2)$ is called *twistor transformation.* Due to the double covering of the connected identity component of the group $SO(2,4)$ (which is denoted as $SO^+(2,4)$) by the group $SU(2,2)$ and due to the double covering of the conformal group $C_+^{\uparrow 4}(1,3)$ [22, v. 2, p. 304 (eng)] by the group $SO^+(2,4)$, the existence of the isomorphisms

$$SU(2,2)/\{\pm 1; \pm i\} \cong LU(2,2) \cong C_+^{\uparrow 4}(1,3) \cong SO^+(2,4)/\pm 1 \tag{14.60}$$

will imply. This means that the group $LU(2,2)$ exhausts all conformal transformations of the group $\mathbb{C}_+^{\uparrow 4}(1,3)$. The matrix S is restored up to a factor λ such that $\lambda^4 = 1$ (det(S)=1) from which we obtain the ambiguity. Since we have the equalities

$$Y = AX + B \quad\Rightarrow\quad dX = AdY,\quad Y = X^{-1} \quad\Rightarrow\quad dX = -X^{-1}dXX^{-1}, \tag{14.61}$$

where A and B are some constant matrixes, then

$$\tilde{Z}^*\ d\tilde{X}\ \tilde{Z} = Z^*\ dX\ Z. \tag{14.62}$$

This equation is an invariant under the group LU(2,2). The proof of this fact is considered in Appendix. Other invariant can be obtained with the help of the identity

$$Y = AX + B \quad\Rightarrow\quad \frac{\partial}{\partial X} = A^T\frac{\partial}{\partial Y},\quad Y = X^{-1} \quad\Rightarrow\quad \frac{\partial}{\partial X} = -Y^T\frac{\partial}{\partial Y}Y^T \tag{14.63}$$

where A and B are also some constant matrixes. This invariant will have the form

$$\tilde{Z}^{-1}\frac{\partial}{\partial \tilde{X}^T}\tilde{Z}^{*\ -1} = Z^{-1}\frac{\partial}{\partial X^T}Z^{*\ -1}. \tag{14.64}$$

The proof can be found in Appendix. This means that there is a real vector $\tilde{L}$ tangent to the hyperboloid obtained in the cross-section of the cone K_6 by the plane V + W = 1. This vector is an invariant under transformations of a basis from the group LU(2,2) (i.e. coordinate-independent in the tangent space to this hyperboloid). The vector is uniquely determined by the matrix

$$\hat{L} := \tfrac{1}{\sqrt{2}}(Z^{-1}\tfrac{\partial}{\partial X^T}Z^{*\ -1} - \bar{Z}^{-1}\tfrac{\partial}{\partial X}Z^{T\ -1}) =$$

$$= \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}(\tfrac{\partial}{\partial\omega}(-\bar{\eta}^0\pi^0 + \eta^0\bar{\pi}^0) + \tfrac{\partial}{\partial\eta}(-\bar{\eta}^1\pi^1 + \eta^1\bar{\pi}^1) + \tag{14.65}$$

$$+\tfrac{\partial}{\partial\xi}(-\bar{\eta}^1\pi^0 + \eta^0\bar{\pi}^1) + \tfrac{\partial}{\partial\bar{\xi}}(\bar{\eta}^0\pi^1 - \eta^1\bar{\pi}^0)) \cdot \tfrac{1}{(det(Z))^2\sqrt{2}} := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}\tilde{L}$$

The norm of this vector in our metric will be such that

$$\parallel \tilde{L} \parallel = -\frac{1}{2(det(Y))^2} = -\frac{1}{(V+W)^2} \tag{14.66}$$

An isotropic vector k is called a vector of *first type unit extension* [22, v. 1, p. 36, eq. (1.4.16) (eng)] in the case when k sets the point belonging to the cross-section of the isotropic cone by the plane V + W = 1. Then $\parallel \tilde{L} \parallel = -1$ and any isotropic vector K collinear with k is defined as

$$K = (- \parallel \tilde{L} \parallel)^{\frac{1}{2}} k. \tag{14.67}$$

However, when V =- W we will obtain a vector with the infinite first type extension. To learn to distinguish between them, it is necessary to set the cross-section of the cone K_6 by the plane T+Z=1 and to enter a vector $\tilde{\tilde{L}}$ with the norm

$$\parallel \tilde{\tilde{L}} \parallel = -\frac{1}{(T+Z)^2} \tag{14.68}$$

in the same way. An isotropic vector k we call a vector of *second type unit extension* in the case when k is sets the point belonging to the cross-section of the isotropic cone by the plane T + Z = 1 and the first type extension will not be finite. We define *extension* of the vector K as

1. *first type extension* if such the extension is finite;
2. *second type extension* if the first type extension is infinite.

Note that the vector $\tilde{L}$ is not coordinate-independent in the space $\mathbb{R}^6_{(2,4)}$ although it is an invariant of the tangent space to the hyperboloid obtained in the cross-section of the cone K_6 by the plane V+W=1. Now there is a possibility to represent *isotropic twistor* in the space ${\mathbb{R}^6}_{(2,4)}$ visually. Consider a pair of vectors of equal extension in ${\mathbb{R}^6}_{(2,4)}$

$$K^\alpha = {\eta^\alpha}_{ab}\ iT^{[a}X^{b]},\quad N^\alpha = {\eta^\alpha}_{ab}\ T^{[a}Z^{b]}\ \Rightarrow\ K^\alpha K_\alpha = 0,\quad N^\alpha K_\alpha = 0,\quad N^\alpha N_\alpha = 0. \tag{14.69}$$

We choose a vector Y^a in such a way as to satisfy the conditions

$$\begin{gathered} Y^aY_a = 0,\quad Y^aX_a = 0,\quad Y^aZ_a = 0, \\ \varepsilon^{abcd}X_aY_bZ_cT_d = X^cZ_cY^dT_d = 1,\quad \varepsilon^{abcd} = 24X^{[a}Y^bZ^cT^{d]}. \end{gathered} \tag{14.70}$$

Thus, we obtain the vector basis X^a, Y^a, Z^a, T^a (recall that $Y_a = s_{aa'}Y^{a'}$)

$$\begin{gathered} Y^aY_a = 0, \quad Y^aX_a = 0, \quad Y^aZ_a = 0, \quad X^aX_a = 0, \\ X^aT_a = 0, \quad Z^aZ_a = 0, \quad Z^aT_a = 0, \quad T^aT_a = 0. \end{gathered} \tag{14.71}$$

Whence,

$$\varepsilon_{abcd}T^{[c}Y^{d]} = -2X_{[a}Z_{b]}. \tag{14.72}$$

Therefore, the vectors

$$L^\alpha = \eta^\alpha{}_{ab}(-T^{[a}Y^{b]} + X^{[a}Z^{b]}), \quad M^\alpha = \eta^\alpha{}_{ab}(-i)(T^{[a}Y^{b]} + X^{[a}Z^{b]}) \tag{14.73}$$

satisfy the parities

$$\begin{gathered} L^\alpha = \bar{L}^\alpha, \quad M^\alpha = \bar{M}^\alpha, \quad L^\alpha K_\alpha = 0, \quad L^\alpha M_\alpha = 0, \quad M^\alpha K_\alpha = 0, \\ L^\alpha N_\alpha = 0, \quad M^\alpha N_\alpha = 0, \quad L^\alpha L_\alpha = -2, \quad M^\alpha M_\alpha = -2. \end{gathered} \tag{14.74}$$

Now we can construct the threevector

$$P^{\alpha\beta\gamma} = 6K^{[\alpha}N^\beta L^{\gamma]}. \tag{14.75}$$

Knowing K^α, we know T^a and X^a up to

$$X^a \longmapsto \lambda_1 X^a + \mu_1 T^a, \quad T^a \longmapsto \nu_1 X^a + \xi_1 T^a, \quad det\begin{pmatrix} \lambda_1 & \mu_1 \\ \nu_1 & \xi_1 \end{pmatrix} = 1. \tag{14.76}$$

And if we know N^α then an arbitrariness in a choice of T^a and Z^a will be such that

$$Z^a \longmapsto \lambda_2 Z^a + \mu_2 T^a, \quad T^a \longmapsto \nu_2 Z^a + \xi_2 T^a, \quad det\begin{pmatrix} \lambda_2 & \mu_2 \\ \nu_2 & \xi_2 \end{pmatrix} = 1. \tag{14.77}$$

Therefor, $\nu_1 = \nu_2 = 0$ and $\xi_2 = \xi_1$. For Y^a, we obtain

$$Y^a \longmapsto \alpha X^a + \beta Y^a + \gamma Z^a + \delta T^a. \tag{14.78}$$

If we now demand that two such bases are related by a transformation from the group LU(2,2) then we obtain

$$\begin{array}{lllllll} X^a & \longmapsto & \tau^{-1}X^a + \mu T^a, & & T^a & \longmapsto & \tau T^a, \\ Z^a & \longmapsto & \tau^{-1}Z^a + \chi T^a, & & Y^a & \longmapsto & -\bar{\chi}X^a + \tau Y^a - \bar{\mu}Z^a + \delta T^a, \end{array} \tag{14.79}$$

$$\bar{\chi}\mu + \bar{\mu}\chi + \tau\bar{\delta} + \bar{\tau}\delta = 0, \quad \tau\bar{\tau} = 1. \tag{14.80}$$

Whence,

$$\begin{array}{lllllll} X^{[a}T^{b]} & \longmapsto & X^{[a}T^{b]} & \Leftrightarrow & K^\alpha & \longmapsto & K^\alpha \\ Z^{[a}T^{b]} & \longmapsto & Z^{[a}T^{b]} & \Leftrightarrow & N^\alpha & \longmapsto & N^\alpha, \end{array} \tag{14.81}$$

$$\begin{array}{lll} T^{[a}Y^{b]} & \longmapsto & -\tau\bar{\chi}T^{[a}X^{b]} + \tau^2 T^{[a}Y^{b]} - \bar{\mu}\tau T^{[a}Z^{b]} \\ X^{[a}Z^{b]} & \longmapsto & \tau^{-2}X^{[a}Z^{b]} + \tau^{-1}\chi X^{[a}T^{b]} + \tau^{-1}\mu T^{[a}Z^{b]}. \end{array} \tag{14.82}$$

Set $\tau =: e^{i\Theta}$ then

$$\begin{gathered} L^\alpha \longmapsto L^\alpha \cos(2\Theta) + M^\alpha \sin(2\Theta) - i(\bar{\chi}\tau - \bar{\tau}\chi)K^\alpha + (\mu\bar{\tau} + \tau\bar{\mu})N^\alpha, \\ M^\alpha \longmapsto M^\alpha \cos(2\Theta) - L^\alpha \sin(2\Theta) + (\bar{\chi}\tau + \bar{\tau}\chi)K^\alpha - i(\mu\bar{\tau} - \tau\bar{\mu})N^\alpha. \end{gathered} \tag{14.83}$$

Thus, the 3-half-plane, spanned by $K^\alpha, N^\alpha, L^\alpha$ is a coordinate-independent in the space $\mathbb{R}^6_{(2,4)}$. Thus, our design can be presented as follows. The first type extension of the vectors K^α and N^α should be the same. K^α and N^α determine *flagpole*: the set of vectors with

1. the first type extension is equal to the first type extension of the vector K^{α};
2. the start coinciding with the beginning of the vector K^{α}.

K^{α}, N^{α}, L^{α} determine the 3-half-plane which we call *flag-plane*. Thus, knowing K^{α} and N^{α}, we know the twistor T^{a} up to the phase Θ. In turn, in the 2-plane (L^{α}, M^{α}), 2Θ is an angle of the rotation of the flag (3-half-plane $P^{\alpha\beta\gamma}$) around the flagpole (N^{α}, K^{α}). Therefore, a rotation of the flag on 2π will lead to the twistor $-T^{a}$, and only a rotation on 4π will return our design to the original state. In addition, collinear twistors can be distinguished from each other using the concept *extension of the vector* for the K^{α} so that under the transformation $T^{a} \longmapsto rT^{a}$, $Y^{a} \longmapsto r^{-1}Y^{a}$ $(r \in R\backslash\{0\})$, the flagpole is multiplied by r and the flag-plane remains unchanged. Finally, it should be noted that the mentioned geometrical structure is uniquely determined by the twistor T^{a}. In the case of the infinite first type extension of the vector K^{α}, we consider the cone $K_4 \subset K_6$ on which the vector N^{α} lays. But non-zero vectors K^{α} and N^{α} have the finite second type extension giving the geometric interpretation of a spinor on the isotropic cone K_4.

14.4 Structure constants of the octonion algebra. Spinor formalism for n = 8

In this subsection, it is told about how on the connecting operators, satisfying the Clifford equation, to construct the structural constants of the octonion algebra. Such the algebra is the special case of the normalized division hypercomplex algebra for n = 8. We consider the axioms of such the algebra. The conclusion of all results of this subsection is made on the basis [4].

Let the operators ${\eta_{\alpha}}^{ab}$ $(\alpha\ ,\beta\ ,... = \overline{1,6}; a\ ,b\ ,... = \overline{1,4})$ be the antisymmetric operators constructed above for the spinor formalism for n = 6. And let the operators ${\eta_{\Lambda}}^{AB}$ $(\Lambda\ ,\Psi\ ,... = \overline{1,8}; A\ ,B\ ,... = \overline{1,8})$ for the spinor formalism for n = 8 be constructed according to the scheme (8.20). Then the tensor (8.16), (8.17) will have the symmetries

$$\varepsilon_{AB(CD)} = \frac{1}{2}\varepsilon_{AB}\varepsilon_{CD}, \quad \varepsilon_{A(B|C|D)} = \frac{1}{2}\varepsilon_{AC}\varepsilon_{BD}, \tag{14.84}$$

where according to (8.19), $\varepsilon_{AC} = \tilde{\varepsilon}_{AC}$ is the metric spin-tensor in the spinor space $\mathbb{C}^8$

$$\eta_{\Lambda AB} = \frac{1}{4}{\eta_{\Lambda}}^{CD}\varepsilon_{ABCD} = {\eta_{\Lambda}}^{CD}\varepsilon_{AD}\varepsilon_{BC}, \quad \eta_{\Lambda} := \frac{1}{4}{\eta_{\Lambda}}^{CD}\varepsilon_{CD}. \tag{14.85}$$

The second identity (14.84) is very important. It leads to the non-degeneracy operator ${\varepsilon_{XA}}^{YB}X^{X}X_{Y}$ and, as a result, to the normability of the octonion algebra. In this case, the identity (9.11) splits into three parts: the alternative identity (left and right) and the flexible identity. As it happens.

Theorem 14.2. *For every $r^{\Lambda} \in \mathbb{C}^8$, there is the decomposition*

$$r^{\Lambda} = {\eta^{\Lambda}}_{AB}X^{A}Y^{B} \tag{14.86}$$

for some X^{A}, $Y^{B} \in \mathbb{C}^8$.

Proof. From the Clifford equation (8.1), the parities

$$R^{AB}R_{CB} = \frac{1}{2}r^{\Lambda}r_{\Lambda}\delta_{C}{}^{A}, \quad R_{KL}\delta_{A}{}^{B} = R_{AR}\varepsilon_{KL}{}^{BR} + R^{BR}\varepsilon_{KLAR} \tag{14.87}$$

will follow. Contract this equation with $P^K P^A$ choosing P^K so that $p := \frac{1}{2}\varepsilon_{AK}P^K P^A \neq 0$ and assuming that $Q_L := R_{KL}P^K$

$$Q_L P^B = Q_R P^K \varepsilon_{KL}{}^{BR} + pR^B{}_L. \tag{14.88}$$

Then from (14.84), the identities

$$R^{BL} = \frac{1}{p}Q^R P^K(\delta_R{}^L\delta_K{}^B - \varepsilon_K{}^{LB}{}_R) = \frac{1}{p}Q^R P^K \varepsilon^{BL}{}_{KR}. \tag{14.89}$$

will follow. Set

$$X^A := \frac{1}{\sqrt{p}}P^A, \quad Y^B := \frac{1}{\sqrt{p}}Q^B \tag{14.90}$$

that proves this theorem. □

Define

$$P^{\Lambda}{}_A := \eta^{\Lambda}{}_{AB}X^A, \quad X^A X_A := 2, \tag{14.91}$$

then

$$P^{\Lambda}{}_A P^{\Psi}{}_B \varepsilon^{AB} = g^{\Lambda\Psi}, \quad P^{\Lambda}{}_A P^{\Psi}{}_B g_{\Lambda\Psi} = \varepsilon_{AB}. \tag{14.92}$$

Thus, the operator $P^{\Lambda}{}_A$ defines the isomorphism between the spaces $\mathbb{C}^n \cong \mathbb{C}^N$ ($n = N = 8$)

$$r^{\Lambda} = P^{\Lambda}{}_B Y^B. \tag{14.93}$$

Set

$$\eta_{\Lambda}{}^{\Psi\Theta} := \sqrt{2}\eta_{\Lambda}{}^{AB}P^{\Psi}{}_A P^{\Theta}{}_B, \tag{14.94}$$

then for these structural constants the alternative octonion identities are executed

$$\eta_{\Lambda\Theta}{}^{\Phi}\eta_{(\Psi\Upsilon)}{}^{\Theta} = \eta_{\Lambda(\,\Psi}{}^{\Theta}\eta_{|\Theta|\Upsilon\,)}{}^{\Phi}, \quad \eta_{(\Lambda\Psi)}{}^{\Theta}\eta_{\Theta\Phi}{}^{\Upsilon} = \eta_{(\,\Lambda|\Theta|}{}^{\Upsilon}\eta_{\Psi\,)\Phi}{}^{\Theta} \tag{14.95}$$

and the central Moufang identity

$$r^{\Phi}\eta_{\Phi\Theta}{}^{\Omega}\eta_{\Lambda\Psi}{}^{\Theta}\eta_{\Omega\Upsilon}{}^{\Gamma}r^{\Upsilon} = r^{\Phi}\eta_{\Phi\Lambda}{}^{\Theta}\eta_{\Theta\Omega}{}^{\Gamma}\eta_{\Psi\Upsilon}{}^{\Omega}r^{\Upsilon} \tag{14.96}$$

is executed too. The proof is given in Appendix. Now, in order to enter the octonion structure constants, we must use the inclusion operator $H_i{}^{\Lambda} : \mathbb{R}^6 \subset \mathbb{C}^6$. Normalizability of the octonion algebra will follow from the Clifford equation (8.1) written out for the structural constants

$$(\eta_{\Phi\Theta}{}^{\Omega}\eta_{\Psi\Xi}{}^{\Delta} + \eta_{\Psi\Theta}{}^{\Omega}\eta_{\Phi\Xi}{}^{\Delta})g_{\Omega\Delta} = 2g_{\Phi\Psi}g_{\Theta\Xi}. \tag{14.97}$$

The operators of the same type (12.5)

$$A_{\Theta\Phi L}{}^{X} := \tfrac{1}{2}\eta_{[\Theta}{}^{MX}\eta_{\Phi]ML}, \quad \tilde{A}_{\Theta\Phi L}{}^{X} := \tfrac{1}{4}(\eta_{[\Theta}{}^{MX}\eta_{\Phi]ML} + \eta_{[\Theta}{}^{XM}\eta_{\Phi]LM}) \tag{14.98}$$

will satisfy the identities

$$\begin{aligned} A_{\Lambda\Psi}{}^{AB}A^{\Lambda\Psi}{}_{CD} &= \delta_{[C}{}^{A}\delta_{D]}{}^{B}, & \tilde{A}_{\Lambda\Psi}{}^{AB}\tilde{A}^{\Lambda\Psi}{}_{CD} &= \tfrac{1}{2}\delta_{[C}{}^{A}\delta_{D]}{}^{B} + \tfrac{1}{4}\varepsilon_{[C}{}^{A}{}_{D]}{}^{B}, \\ A_{\Lambda\Psi}{}^{AB}A^{\Upsilon\Omega}{}_{AB} &= \delta_{[\Lambda}{}^{\Upsilon}\delta_{\Psi]}{}^{\Omega}, & \tilde{A}_{\Lambda\Psi}{}^{AB}\tilde{A}^{\Upsilon\Omega}{}_{AB} &= \delta_{[\Lambda}{}^{\Upsilon}\delta_{\Psi]}{}^{\Omega} + \eta^{[\Upsilon}\delta_{[\Lambda}{}^{\Omega]}\eta_{\Psi]}. \end{aligned} \tag{14.99}$$

The proof is given in Appendix.

14.5 Geometry of the inductive step from n=6 to n=8

In this subsection, it is told about how to construct two quadrics associated to the Cartan triality principle [29, p. 175 (rus)]. The construction is carried out in the explicit form justifying the standard scheme of the induction transition from n = 6 to n = 8. The conclusion of all results of this subsection is made on the basis [6], [29], [30, v. 2].

14.5.1 Rosenfeld null-pair

In this subsubsection, it is told about how to construct an explicit solution of the twistor equation for n=6. The conclusion of all results of this subsubsection is made on the basis [6], [19].

Denote by $\mathbb{A}^{\mathbb{C}*}$ the spinor 4-dimensional complex vector space. Such the space is dual to the space $\mathbb{A}^{\mathbb{C}}$. Then the 8-dimensional complex space $\mathbb{T}^2$ is formed as the direct sum $\mathbb{A}^{\mathbb{C}} \oplus \mathbb{A}^{\mathbb{C}*}$. That is, if X^a $(a, b, ... = \overline{1, 4})$ are the coordinates of a vector in $A^{\mathbb{C}}$, and Y_b are the coordinates of a covector in $A^{\mathbb{C}*}$ then $X^A := (X^a, Y_b),\ (A, B, ... = \overline{1, 8})$ are the coordinates of a vector of $\mathbb{T}^2$. We will consider the bivector coordinates r^{ab} (13.18)

$$X^a = ir^{ab}Y_b \tag{14.100}$$

as the ones in the complex affine space $\mathbb{C}\mathbb{A}_6$. This is a system of 4 linear equations with 6 unknowns. To determine its rank, we consider *homogeneous equation* $r^{ab}Y_b = 0$ which has nontrivial solutions if and only if the bivector is simple: $r_{kc}r^{ac} = \frac{1}{4}r^{ab}r_{ab}\delta_k{}^d = 0$, and therefore can be represented as

$$r^{ab}_{\text{homogeneous}} = P^aQ^b - P^bQ^a, \tag{14.101}$$

where P^a and Q^a are defined up to linear combinations of them. From this, it follows that $P^aY_a = 0,\ \ Q^aY_a = 0$. For $X^aY_a = 0$, by X^a, S^a, Z^a, we denote all solutions, that form a basis. Then our solution takes the form

$$r^{ab}_{\text{homogeneous}} = \lambda_1 S^{[a}X^{b]} + \lambda_2 X^{[a}Z^{b]} + \lambda_3 S^{[a}Z^{b]} \tag{14.102}$$

and hence determines a 3-dimensional subspace in the bivector space. From here, *common solution*

$$r^{ab} = r^{ab}_{\text{particular}} + \lambda_1 S^{[a}X^{b]} + \lambda_2 X^{[a}Z^{b]} + \lambda_3 S^{[a}Z^{b]} \tag{14.103}$$

is obtained, where $r^{ab}_{\text{particular}}$ is an arbitrary bivector being *particular solution* of (14.100).

14.5.2 Construction of the quadrics $\mathbb{C}Q_6$ and $\mathbb{C}\tilde{Q}_6$

In this subsubsection, it is told about how to construct a quadric satisfying the Cartan triality principle. The conclusion of all results of this subsubsection is made on the basis [6], [29], [30, v. 2].

The space $\mathbb{T}^2$ will be a complex space in which the scalar square of a vector is determined by the quadratic form

$$\varepsilon_{AB}X^AX^B = 2X^aY_a \tag{14.104}$$

so that the matrix of the spin-tensor ε_{AB} has the form

$$\parallel \varepsilon_{AB} \parallel = \begin{pmatrix} 0 & \delta_a{}^c \\ \delta^b{}_d & 0 \end{pmatrix}. \tag{14.105}$$

For the fixed r^{ab}, the equation (14.100) will define the 4-dimensional subspace in $\mathbb{T}^2$ which will be the 4-dimensional planar generators of the cone $\varepsilon_{AB}X^AX^B = 0$. Thus, in the projective space $\mathbb{CP}_7$, we can consider the quadric $\mathbb{C}Q_6$ defined by the equation (14.104). 4 basis points of the generator satisfy the condition $\varepsilon_{AB}X^A_IX^B_J = 0, \quad (I, J, ... = \overline{1, 4})$. Set

$$X^A_1 := (X^a, Y_b), \quad X^A_2 := (Z^a, T_b), \quad X^A_3 := (L^a, N_b), \quad X^A_4 := (K^a, M_b). \tag{14.106}$$

On the basis of the common solution of the equation (14.100), each point of the quadric $\mathbb{C}Q_6$ can be associated to the 3-dimensional isotropic plane of the space $\mathbb{CA}_6$. The point (t,v,w,x,y,z) of the space $\mathbb{CA}_6$ can be represented by the line $(\lambda T, \lambda V, \lambda U, \lambda S, \lambda W, \lambda X, \lambda Y, \lambda Z)$ of the space $\mathbb{CR}^8$ having the metric

$$dL^2 = dT^2 + dV^2 + dU^2 + dS^2 + dW^2 + dX^2 + dY^2 + dZ^2. \tag{14.107}$$

This line will be a generator of the isotropic cone $\mathbb{C}K_8$. The intersection of the 7-plane $U - iS = 1$ with the cone $\mathbb{C}K_8$ has the induced metric

$$d\tilde{L}^2 = dT^2 + dV^2 + dW^2 + dX^2 + dY^2 + dZ^2. \tag{14.108}$$

This space has the form of a paraboloid in $\mathbb{C}K_8$ and is identical to the space $\mathbb{CR}^6$: $U = 1+iS = \frac{1}{2}(1-T^2-V^2-W^2-X^2-Y^2-Z^2)$. Every generator of the cone (a set of points belonging to $\mathbb{C}K_8$ with the constant ratio T:V:U:S:W:X:Y:Z), not lying on the hyperplane $U = iS$, intersects the paraboloid in the single point. Every generator of the cone, lying on the hyperplane $U = iS$, corresponds to the point belonging to the infinity of the space $\mathbb{CR}^6$. Thus, straight lines of $\mathbb{CR}^8$, passing through the origin of $\mathbb{CR}^8$, correspond to points of the projective space $\mathbb{CP}_7$. The stereographic projection of this section on the plane (S=0, U=1) with the pole $N(0, 0, \frac{1}{2}, \frac{i}{2}, 0, 0, 0, 0)$ maps the point P(T,V,U,S,W,X,Y,Z) of the hyperboloid to the point p(t,v,1,0,w,x,y,z) of the plane (S=0, U=1)

$$\begin{gathered} \lambda T = t, \quad \lambda V = v, \quad \lambda W = w, \quad \lambda X = x, \quad \lambda Y = y, \quad \lambda Z = z, \quad \lambda = \tfrac{1}{U+iS}, \\ r^\alpha r_\alpha = -\tfrac{U-iS}{U+iS}, \quad \lambda U = \tfrac{1}{2}(1 - t^2 - v^2 - w^2 - x^2 - y^2 - z^2) = -\lambda iS + 1. \end{gathered} \tag{14.109}$$

All generators of the same cone $\mathbb{C}K_8$ form the quadric $\mathbb{C}\tilde{Q}_6$ in the projective space $\mathbb{CP}_7$

$$g_{\Lambda\Psi}r^\Lambda r^\Psi = 0. \tag{14.110}$$

14.6 Correspondence $\mathbb{C}Q_6 \longmapsto \mathbb{C}\tilde{Q}_6$

In this subsubsection, it is told about how to construct a correspondence between generators of the quadric of different ranks according to the Cartan triality principle. The conclusion of all results of this subsubsection is made on the basis [6], [29], [30, v. 2].

1. The common solution of the equation (14.100)

   $$r^{ab} = r^{ab}_{\text{particular}} + r^{ab}_{\text{homogeneous}} = r^{ab}_{\text{particular}} + \lambda_1 S^{[a}X^{b]} + \lambda_2 X^{[a}Z^{b]} + \lambda_3 S^{[a}Z^{b]} \tag{14.111}$$

   determines the 4-dimensional planar generator of the cone $\mathbb{C}K_8$. Then for such the generator, the system

   $$\begin{cases} ir^{ab}Y_b &= X^a, \\ ir^{ab}T_b &= Z^a, \\ ir^{ab}N_b &= L^a, \\ ir^{ab}M_b &= K^a \end{cases} \tag{14.112}$$

   will be determined with the conditions

   $$\begin{gathered} X^aY_a = 0, \quad Z^aT_a = 0, \quad L^aN_a = 0, \quad K^aM_a = 0, \\ X^aT_a = -Z^aY_a, \quad X^aN_a = -L^aY_a, \quad X^aM_a = -K^aY_a, \\ Z^aN_a = -L^aT_a, \quad Z^aM_a = -K^aT_a, \quad K^aN_a = -L^aM_a. \end{gathered} \tag{14.113}$$

   Thus, from the 16 equations with the 6 unknowns r^{ab} only 6 from them will be significant equations (the 10 communication conditions) that defines the point of $\mathbb{CA}_6$ and hence the point of the quadric $\mathbb{C}\tilde{Q}_6$.

2. If we know only one equation

   $$ir^{ab}Y_b = X^a \tag{14.114}$$

   with the condition

   $$X^aY_a = 0 \tag{14.115}$$

   then from the 4 equations, only 3 from them will be significant equations (the 1 communication condition). This means that the point of the quadric $\mathbb{C}Q_6$ will uniquely define the 3-dimensional planar generator $\mathbb{CP}_3$ belonging to the quadric $\mathbb{C}\tilde{Q}_6$.

3. If we know the two equations

   $$\begin{cases} ir^{ab}Y_b &= X^a, \\ ir^{ab}T_b &= Z^a \end{cases} \tag{14.116}$$

   with the conditions

   $$X^aY_a = 0, \quad Z^aT_a = 0, \quad X^aT_a = -Z^aY_a \tag{14.117}$$

   then from 8 equations, only 5 from them will be significant (the 6 unknowns and the 3 communication conditions). This means that the rectilinear generator $\mathbb{CP}_1$ of the quadric $\mathbb{C}Q_6$ will uniquely define the rectilinear generator $\mathbb{CP}_1$ belonging to the quadric $\mathbb{C}\tilde{Q}_6$. In this case, the manifold of generators $\mathbb{CP}_1(\mathbb{C}Q_6)$, belonging to the same generator $\mathbb{CP}_3(\mathbb{C}Q_6)$, defines the beam of generators $\mathbb{CP}_1(\mathbb{C}\tilde{Q}_6)$ belonging to the quadric $\mathbb{C}\tilde{Q}_6$ (this beam is a cone). The center of the beam is determined by the system (14.112).



Fig. 1: Correspondence $\forall CP_2 \subset CP_3 \leftrightarrow R$



Fig. 2: Correspondence $CP_3 \supset CP_1 \leftrightarrow CP_1 \subset K_6$

4. If we know the three equations

$$\left\{\begin{array}{lcl} ir^{ab}Y_b & = & X^a, \\ ir^{ab}T_b & = & Z^a, \\ ir^{ab}N_b & = & L^a \end{array}\right. \tag{14.118}$$

with the conditions

$$\begin{gathered} X^aY_a = 0, \quad Z^aT_a = 0, \quad L^aN_a = 0, \\ X^aT_a = -Z^aY_a, \quad X^aN_a = -L^aY_a, \quad Z^aN_a = -L^aT_a \end{gathered} \tag{14.119}$$

then from the 12 equations only 6 from them will be significant (the 6 unknowns and the 6 communication conditions). This means, that the 2-dimensional generator $\mathbb{CP}_2$ of the quadric $\mathbb{C}Q_6$ will uniquely define the point of the quadric $\mathbb{C}\tilde{Q}_6$. At the same time, the manifold of generators $\mathbb{CP}_2(\mathbb{C}Q_6)$, belonging to the same generator $\mathbb{CP}_3(\mathbb{C}Q_6)$, uniquely determines the same point of the quadric $\mathbb{C}\tilde{Q}_6$. This point is determined by the system (14.112).

14.6.1 Geometry of the transition to the connecting operators ${\eta_\Lambda}^{KL}$ for n = 8

In this subsubsection, it is told about how to construct the connecting operators of the spinor formalism for n = 8. The conclusion of all results of this subsubsection is made on the basis [6], [29], [30, v. 2].

Further we consider the bivector $\hat{R}^{AB}$ such that its vector components ${X_1}^A, {X_2}^A$ are defined as

$$X_1^A := (X^a, Y_b), \quad X_2^A := (Z^a, T_b), \quad \left\{\begin{array}{lcl} ir^{ab}Y_b & = & X^a, \\ ir^{ab}T_b & = & Z^a. \end{array}\right. \tag{14.120}$$

This defines the identity chain

$$\begin{gathered} ir^{ab}Y_b = X^a, \quad ir_{ab}Z^b = pf(r)T_a, \\ \tfrac{1}{2}ir_{cd}\varepsilon^{abcd}Y_b\varepsilon_{aklm} = X^a\varepsilon_{aklm}, \\ 3ir_{[kl}Y_{m]} = X^a\varepsilon_{aklm}, \\ ir_{kl}Y_m + 2ir_{m[k}Y_{l]} = X^a\varepsilon_{aklm}. \end{gathered} \tag{14.121}$$

Contract the last identity with Z^m that will give

$$\begin{gathered} ir_{kl}Y_mZ^m = X^mZ^n\varepsilon_{klmn} + pf(r)\cdot 2T_{[k}Y_{l]}, \\ X^mZ^n - X^nZ^m = -ir^{mn}T_kX^k + \varepsilon^{klmn}pf(r)Y_{[k}T_{l]}. \end{gathered} \tag{14.122}$$

Based on the foregoing, for the quadric $\mathbb{C}Q_6$, we consider the rectilinear generator defined by the bivector $(T_kX^k \neq 0)$

$$\hat{R}^{AB} = 2X_1^{[A}X_2^{B]} = \begin{pmatrix} X^aZ^b - X^bZ^a & X^aT_d - Y_dZ^a \\ Y_cZ^b - T_cX^b & Y_cT_d - Y_dT_c \end{pmatrix} =$$

$$= \begin{pmatrix} -ir^{ab}T_kX^k + \varepsilon^{klab}pf(r)Y_kT_l & 2ir^{ar}Y_{[r}T_{b]} \\ i\varepsilon^{kbmn}r_{ck}Y_{[m}T_{n]} - \delta_c{}^bX^kT_k & 2Y_{[c}T_{d]} \end{pmatrix} =$$

$$= \underbrace{T_lX^l \begin{pmatrix} -\frac{1}{2}i\delta^a{}_kr^\gamma r_\gamma & r^{ar} \\ r_{ck} & -i\delta_c{}^r \end{pmatrix}}_{:=R^A{}_K} \cdot \underbrace{\frac{1}{T_lX^l} \begin{pmatrix} \varepsilon^{kbmn}iY_mT_n & 0 \\ -i\delta_r{}^bX^nT_n & 2iY_{[r}T_{d]} \end{pmatrix}}_{:=P^{KB}} := R^A{}_KP^{KB}, \tag{14.123}$$

$$R^{AB} := \varepsilon^{BC}R^A{}_C = T_lX^l \begin{pmatrix} r^{an} & -\frac{1}{2}i\delta^a{}_kr^\gamma r_\gamma \\ -i\delta_c{}^n & r_{ck} \end{pmatrix}, \tag{14.124}$$

$$R_{AB} := \varepsilon_{AC}R^C{}_B = T_lX^l \begin{pmatrix} r_{ck} & -i\delta_c{}^n \\ -\frac{1}{2}i\delta^a{}_kr^\gamma r_\gamma & r^{an} \end{pmatrix}$$

where ε_{AC} is the same metric symmetric spin-tensor. At the same time, the equation

$$R_A{}^C\hat{R}^{AB} = 0 \tag{14.125}$$

will be true that means that any spin-tensor $\hat{R}^{AB}$, representing the generator $\mathbb{CP}_1(\mathbb{C}Q_6)$, will contain the same tensor $R^A{}_K$ in its expansion, wherein, the second spin-tensor P^{KB} of the decomposition will be responsible for the position of $\mathbb{CP}_1$ in $\mathbb{CP}_3$. Therefore, there is a reason to assign the bispinor R^{AB} to the point of the quadric $\mathbb{C}\tilde{Q}_6$. This point is uniquely determined. In the transition to the space $\mathbb{CR}^8$ on the basis of

$$\begin{array}{lllllllll} r^{12} & = & \frac{1}{\sqrt{2}}(v+iw), & r^{13} & = & \frac{1}{\sqrt{2}}(x+iy), & r^{14} & = & \frac{i}{\sqrt{2}}(t+iz), \\ r^{23} & = & \frac{i}{\sqrt{2}}(iz-t), & r^{24} & = & \frac{1}{\sqrt{2}}(-x+iy), & r^{34} & = & \frac{1}{\sqrt{2}}(v-iw) \end{array} \tag{14.126}$$

we define *homogeneous coordinates* of $\mathbb{CR}^8$ as follows

$$\lambda = \left\{ \begin{array}{ccccccccccccccc} R^{12} & : & R^{13} & : & R^{14} & : & R^{23} & : & R^{24} & : & R^{34} & : & R^{15} & : & R^{51}, \\ \\ r^{12} & : & r^{13} & : & r^{14} & : & r^{23} & : & r^{24} & : & r^{34} & : & -\frac{1}{2}ir^\gamma r_\gamma & : & -i, \end{array} \right. \tag{14.127}$$

$$\begin{array}{lllllllll} R^{12} & = & \frac{1}{\sqrt{2}}(V+iW), & R^{13} & = & \frac{1}{\sqrt{2}}(iY+X), & R^{14} & = & \frac{1}{\sqrt{2}}(-Z+iT), \\ R^{23} & = & \frac{1}{\sqrt{2}}(-Z-iT), & R^{24} & = & \frac{1}{\sqrt{2}}(-X+iY), & R^{34} & = & \frac{1}{\sqrt{2}}(V-iW), \\ R^{15} & = & \frac{1}{2}(iU+S), & R^{51} & = & -iU+S, & & & \end{array} \tag{14.128}$$

$$\begin{array}{lllllll} R^{12} & = & -R^{21} & = & R^{78} & = & -R^{87}, \\ R^{13} & = & -R^{31} & = & R^{86} & = & -R^{68}, \\ R^{14} & = & -R^{41} & = & R^{67} & = & -R^{76}, \\ R^{23} & = & -R^{32} & = & R^{58} & = & -R^{85}, \\ R^{24} & = & -R^{42} & = & R^{75} & = & -R^{57}, \\ R^{34} & = & -R^{43} & = & R^{56} & = & -R^{65}, \\ R^{15} & = & R^{26} & = & R^{37} & = & R^{48}, \\ R^{51} & = & R^{62} & = & R^{73} & = & R^{84}. \end{array} \tag{14.129}$$

$$R^{AB}R_{AB} = 8(R^{12}R^{34} - R^{13}R^{24} + R^{14}R^{23} + R^{15}R^{51}) =$$

$$= 4(T^2 + V^2 + U^2 + S^2 + W^2 + X^2 + Y^2 + Z^2) := 4nf(R), \tag{14.130}$$

$$R^{AB}R_{CB} = \tfrac{1}{2}nf(R){\delta_C}^A \quad \Rightarrow \quad R^{AB}R_{AB} = 4nf(R).$$

For $(r_i)^\Lambda$, to define a generator of the quadric $\mathbb{C}\tilde{Q}_6$, it is necessary and sufficient to have the condition

$$g_{\Lambda\Psi}(r_I)^\Lambda(r_J)^\Psi = 0. \tag{14.131}$$

We define some connecting operators ${\eta_\Lambda}^{BC}$ so that to satisfy the conditions

$$r_\Lambda = \frac{1}{4}{\eta_\Lambda}^{BC}R_{BC}, \quad r^\Lambda = \frac{1}{4}{\eta^\Lambda}_{BC}R^{BC}. \tag{14.132}$$

Then these operators satisfy the Clifford equation

$$g_{\Lambda\Psi}{\delta_K}^L = {\eta_{\Lambda K}}^R{{\eta_\Psi}^L}_R + {\eta_{\Lambda K}}^R{{\eta_\Psi}^L}_R. \tag{14.133}$$

Define

$$\varepsilon_{PQRT} := {\eta^\Lambda}_{PQ}{\eta^\Psi}_{RT}g_{\Lambda\Psi}, \quad \varepsilon_{PQRT} = \varepsilon_{RTPQ} \tag{14.134}$$

that will give another metric spin-tensor ε_{PQRT} with which the help we can raise and lower a pair of indices

$$\begin{gathered}
g_{\Lambda\Psi} = \tfrac{1}{4}{\eta_\Lambda}^{PQ}\eta_{\Psi PQ}, \quad \eta_{\Lambda RT} = \tfrac{1}{4}{\eta_\Lambda}^{PQ}\varepsilon_{PQRT}, \quad \varepsilon_{PQ} = \tfrac{1}{4}\varepsilon_{PQRT}\varepsilon^{RT}, \\
\varepsilon_{STPQ}{\delta_K}^L = {\varepsilon_{STK}}^R{{\varepsilon_{PQ}}^L}_R + {\varepsilon_{PQK}}^R{{\varepsilon_{ST}}^L}_R, \quad \varepsilon_{STPQ} = \tfrac{1}{4}{\varepsilon_{ST}}^{KR}\varepsilon_{PQKR}, \\
\varepsilon_{PQ(KL)} = \tfrac{1}{2}\varepsilon_{PQ}\varepsilon_{KL}, \quad \varepsilon_{[PQ](KL)} = 0, \quad {\eta_\Lambda}^{(MN)} = \tfrac{1}{8}{\eta_\Lambda}^{KL}\varepsilon_{KL}\varepsilon^{MN}.
\end{gathered} \tag{14.135}$$

The result of the applying of the two metric spin-tensors should be the same. If the matrixes of the tensors $g_{\Lambda\Psi}$ and ε_{KL} have the form

$$\parallel g_{\Lambda\Psi} \parallel = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad \parallel \varepsilon_{KL} \parallel = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix} \tag{14.136}$$

then the significant coordinates of the operators ${\eta^\Lambda}_{KL}$ in some basis would be such that

$$\begin{array}{llll}
{\eta^2}_{12} = \frac{1}{\sqrt{2}}, & {\eta^2}_{34} = \frac{1}{\sqrt{2}}, & {\eta^5}_{12} = -\frac{i}{\sqrt{2}}, & {\eta^5}_{34} = \frac{i}{\sqrt{2}}, \\
{\eta^2}_{78} = \frac{1}{\sqrt{2}}, & {\eta^2}_{56} = \frac{1}{\sqrt{2}}, & {\eta^5}_{78} = -\frac{i}{\sqrt{2}}, & {\eta^5}_{56} = \frac{i}{\sqrt{2}}, \\
{\eta^1}_{14} = -\frac{i}{\sqrt{2}}, & {\eta^1}_{23} = \frac{i}{\sqrt{2}}, & {\eta^8}_{14} = -\frac{1}{\sqrt{2}}, & {\eta^8}_{23} = -\frac{1}{\sqrt{2}}, \\
{\eta^1}_{67} = -\frac{i}{\sqrt{2}}, & {\eta^1}_{58} = \frac{i}{\sqrt{2}}, & {\eta^8}_{67} = -\frac{1}{\sqrt{2}}, & {\eta^8}_{58} = -\frac{1}{\sqrt{2}}, \\
{\eta^7}_{13} = -\frac{i}{\sqrt{2}}, & {\eta^7}_{24} = -\frac{i}{\sqrt{2}}, & {\eta^6}_{13} = \frac{1}{\sqrt{2}}, & {\eta^6}_{24} = -\frac{1}{\sqrt{2}}, \\
{\eta^7}_{68} = \frac{i}{\sqrt{2}}, & {\eta^7}_{57} = \frac{i}{\sqrt{2}}, & {\eta^6}_{68} = -\frac{1}{\sqrt{2}}, & {\eta^6}_{57} = \frac{1}{\sqrt{2}}, \\
{\eta^4}_{15} = \frac{1}{\sqrt{2}}, & {\eta^4}_{51} = \frac{1}{\sqrt{2}}, & {\eta^3}_{15} = -\frac{i}{\sqrt{2}}, & {\eta^3}_{51} = \frac{i}{\sqrt{2}}, \\
{\eta^4}_{26} = \frac{1}{\sqrt{2}}, & {\eta^4}_{62} = \frac{1}{\sqrt{2}}, & {\eta^3}_{26} = -\frac{i}{\sqrt{2}}, & {\eta^3}_{62} = \frac{i}{\sqrt{2}}, \\
{\eta^4}_{37} = \frac{1}{\sqrt{2}}, & {\eta^4}_{73} = \frac{1}{\sqrt{2}}, & {\eta^3}_{37} = -\frac{i}{\sqrt{2}}, & {\eta^3}_{73} = \frac{i}{\sqrt{2}}, \\
{\eta^4}_{48} = \frac{1}{\sqrt{2}}, & {\eta^4}_{84} = \frac{1}{\sqrt{2}}, & {\eta^3}_{48} = -\frac{i}{\sqrt{2}}, & {\eta^3}_{84} = \frac{i}{\sqrt{2}},
\end{array} \tag{14.137}$$

and the significant coordinates of the operators ${\eta_\Lambda}^{KL}$ would be such that

$$\begin{array}{llll}
{\eta_2}^{12} = \frac{1}{\sqrt{2}}, & {\eta_2}^{34} = \frac{1}{\sqrt{2}}, & {\eta_5}^{12} = \frac{i}{\sqrt{2}}, & {\eta_5}^{34} = -\frac{i}{\sqrt{2}}, \\
{\eta_2}^{78} = \frac{1}{\sqrt{2}}, & {\eta_2}^{56} = \frac{1}{\sqrt{2}}, & {\eta_5}^{78} = \frac{i}{\sqrt{2}}, & {\eta_5}^{56} = -\frac{i}{\sqrt{2}}, \\
{\eta_1}^{14} = \frac{i}{\sqrt{2}}, & {\eta_1}^{23} = -\frac{i}{\sqrt{2}}, & {\eta_8}^{14} = -\frac{1}{\sqrt{2}}, & {\eta_8}^{23} = -\frac{1}{\sqrt{2}}, \\
{\eta_1}^{67} = \frac{i}{\sqrt{2}}, & {\eta_1}^{58} = -\frac{i}{\sqrt{2}}, & {\eta_8}^{67} = -\frac{1}{\sqrt{2}}, & {\eta_8}^{58} = -\frac{1}{\sqrt{2}}, \\
{\eta_7}^{13} = \frac{i}{\sqrt{2}}, & {\eta_7}^{24} = \frac{i}{\sqrt{2}}, & {\eta_6}^{13} = \frac{1}{\sqrt{2}}, & {\eta_6}^{24} = -\frac{1}{\sqrt{2}}, \\
{\eta_7}^{68} = -\frac{i}{\sqrt{2}}, & {\eta_7}^{57} = -\frac{i}{\sqrt{2}}, & {\eta_6}^{68} = -\frac{1}{\sqrt{2}}, & {\eta_6}^{57} = \frac{1}{\sqrt{2}}, \\
{\eta_4}^{15} = \frac{1}{\sqrt{2}}, & {\eta_4}^{51} = \frac{1}{\sqrt{2}}, & {\eta_3}^{15} = \frac{i}{\sqrt{2}}, & {\eta_3}^{51} = -\frac{i}{\sqrt{2}}, \\
{\eta_4}^{26} = \frac{1}{\sqrt{2}}, & {\eta_4}^{62} = \frac{1}{\sqrt{2}}, & {\eta_3}^{26} = \frac{i}{\sqrt{2}}, & {\eta_3}^{62} = -\frac{i}{\sqrt{2}}, \\
{\eta_4}^{37} = \frac{1}{\sqrt{2}}, & {\eta_4}^{73} = \frac{1}{\sqrt{2}}, & {\eta_3}^{37} = \frac{i}{\sqrt{2}}, & {\eta_3}^{73} = -\frac{i}{\sqrt{2}}, \\
{\eta_4}^{48} = \frac{1}{\sqrt{2}}, & {\eta_4}^{84} = \frac{1}{\sqrt{2}}, & {\eta_3}^{48} = \frac{i}{\sqrt{2}}, & {\eta_3}^{84} = -\frac{i}{\sqrt{2}}.
\end{array} \tag{14.138}$$

The basis is slightly different from one in which the homogeneous coordinates of $\mathbb{CR}^8$ were defined above. Thus, the bispinor $\hat{R}^{AB}$ determines the rectilinear generator $\mathbb{CP}_1(\mathbb{C}Q_6)$ belonging to the planar generator $\mathbb{CP}_3(\mathbb{C}Q_6)$ which determines the point with the coordinates R^{AB} for the quadric $\mathbb{C}\tilde{Q}_6$. We define the spin-tensor $\hat{\hat{R}}^{AB}$

$$\hat{\hat{R}}^{AB} := \hat{R}^{AK}\hat{P}_K{}^B, \quad \hat{P}_K{}^B := \begin{pmatrix} 2i\delta_m{}^k & 0 \\ 0 & -2pf(r)i\delta^n{}_r \end{pmatrix}. \tag{14.139}$$

The tensor $\hat{\hat{R}}^{AB}$ will continue to represent the rectilinear generator $\mathbb{CP}_1(\mathbb{C}Q_6)$ belonging to the planar generator $\mathbb{CP}_3(\mathbb{C}Q_6)$. We apply the operators $\eta^\Lambda{}_{KL}$ to the spin-tensor $\hat{\hat{R}}^{AB}$ and obtain

$$\eta^\Lambda{}_{KL}\hat{\hat{R}}^{KL} = \eta^\Lambda{}_{KL}T_mX^m \begin{pmatrix} r^{an} & -\frac{1}{2}i\delta^a{}_k r^\gamma r_\gamma \\ -i\delta_c{}^n & r_{ck} \end{pmatrix} = \eta^\Lambda{}_{KL}R^{KL}. \tag{14.140}$$

Thus, the operators $\eta^\Lambda{}_{KL}$ take each rectilinear generator $\mathbb{CP}_1(\mathbb{C}Q_6)$, belonging to the planar generator $\mathbb{CP}_3(\mathbb{C}Q_6)$, to the same generator $\mathbb{CP}_3(\mathbb{C}Q_6)$, and this determines the point of the quadric $\mathbb{C}\tilde{Q}_6$. In the homogeneous coordinates, the same tensor R^{KL} determines the coordinates of the point R of the space $\mathbb{CR}^8$.

Let us find out what is the number of a family which comprises this generator $\mathbb{CP}_3(\mathbb{C}Q_6)$. For this purpose, we consider the conditions

$$\begin{gathered} \varepsilon_{AB}(X_I)^A(X_J)^B = 0, \\ X^{ABCD} := \varepsilon^{IJKL}(X_I)^A(X_J)^B(X_K)^C(X_L)^D, \end{gathered} \tag{14.141}$$

where ε^{IJKL} is a 4-vector antisymmetrical in all indices. Also we consider the 8-vector $e_{ABCDKLMN}$ antisymmetrical in all indices too. If in the condition

$$\frac{1}{24}e_{ABCDKLMN}X^{ABCD} = \rho\varepsilon_{KR}\varepsilon_{LT}\varepsilon_{MU}\varepsilon_{NV}X^{RTUV}, \quad \rho^2 = 1, \tag{14.142}$$

$\rho = 1$ then we say that the planar generator $\mathbb{CP}_3(\mathbb{C}Q_6)$ belongs to family I, and if $\rho = -1$ then the planar generator belongs to family II. In our case

$$(X_I)^A = ((X_I)^a, (Y_I)_b), \tag{14.143}$$

and then

$$\varepsilon^{IJKL}(X_I)^1(X_J)^2(X_K)^3(X_L)^4 = \rho\varepsilon^{IJKL}(X_I)^1(X_J)^2(X_K)^3(X_L)^4. \tag{14.144}$$

Whence, $\rho = 1$. This means that our generators should belong to family I. In addition, there is the tensor ${P_\Lambda}^B := {\eta_\Lambda}^{AB}X_A$ for $X_A = (1,0,0,0,1,0,0,0)$

$$\parallel {P_\Lambda}^B \parallel = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 & i & 0 & 0 & 0 & -i \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ -i & 0 & 0 & 0 & i & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & i & 0 & 0 & 0 & -i & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & i & 0 & 0 & 0 & -i & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 \end{pmatrix}, \quad det \parallel {P_\Lambda}^B \parallel = 1, \tag{14.145}$$

$$\varepsilon_{AB}{P_\Lambda}^A{P_\Psi}^B = g_{\Lambda\Psi}, \quad (X_I)^A = {P_\Lambda}^A(r_I)^\Lambda.$$

Then from

$$\begin{gathered} g_{\Lambda\Psi}(r_I)^\Lambda(r_J)^\Psi = 0, \\ X^{ABCD} = {P_\Lambda}^A{P_\Psi}^B{P_\Phi}^C{P_\Theta}^D r^{\Lambda\Psi\Phi\Theta} \end{gathered} \tag{14.146}$$

it follows that $\rho = 1$ and

$$\begin{gathered} \tfrac{1}{24}e_{ABCDKLMN}X^{ABCD} = \varepsilon_{KR}\varepsilon_{LT}\varepsilon_{MU}\varepsilon_{NV}X^{RTUV}, \\ \tfrac{1}{24}e_{ABCDKLMN}{P_\Lambda}^A{P_\Psi}^B{P_\Phi}^C{P_\Theta}^D r^{\Lambda\Psi\Phi\Theta} = \\ = \varepsilon_{KA}\varepsilon_{LB}\varepsilon_{MC}\varepsilon_{ND}{P_{\Lambda_1}}^A{P_{\Psi_1}}^B{P_{\Phi_1}}^C{P_{\Theta_1}}^D r^{\Lambda_1\Psi_1\Phi_1\Theta_1}, \\ \tfrac{1}{24}e_{ABCDKLMN}{P_\Lambda}^A{P_\Psi}^B{P_\Phi}^C{P_\Theta}^D{P_{\Lambda_1}}^K{P_{\Psi_1}}^L{P_{\Phi_1}}^M{P_{\Theta_1}}^N r^{\Lambda\Psi\Phi\Theta} = \\ = g_{\Lambda\Lambda_1}g_{\Psi\Psi_1}g_{\Phi\Phi_1}g_{\Theta\Theta_1}r^{\Lambda_1\Psi_1\Phi_1\Theta_1}, \\ \tfrac{1}{24}det \parallel {P_\Omega}^A \parallel e_{\Lambda\Psi\Phi\Theta\Lambda_1\Psi_1\Phi_1\Theta_1}r^{\Lambda_1\Psi_1\Phi_1\Theta_1} = g_{\Lambda\Lambda_1}g_{\Psi\Psi_1}g_{\Phi\Phi_1}g_{\Theta\Theta_1}r^{\Lambda_1\Psi_1\Phi_1\Theta_1}, \\ \tfrac{1}{24}e_{\Lambda\Psi\Phi\Theta\Lambda_1\Psi_1\Phi_1\Theta_1}r^{\Lambda_1\Psi_1\Phi_1\Theta_1} = g_{\Lambda\Lambda_1}g_{\Psi\Psi_1}g_{\Phi\Phi_1}g_{\Theta\Theta_1}r^{\Lambda_1\Psi_1\Phi_1\Theta_1}, \end{gathered} \tag{14.147}$$

This shows that the generators $\mathbb{CP}_3(\mathbb{C}\tilde{Q}_6)$ must belong to family I. To obtain family II, we must act by the elementary orthogonal reflection (with determinant equal to -1) on the operator ${P_\Lambda}^K$.

14.7 Correspondence $\mathbb{C}\tilde{Q}_6 \longmapsto \mathbb{C}Q_6$

In this subsubsection it is told about how to construct a reverse mapping for the quadrics between generators of different ranks according to the Cartan triality principle. The conclusion of all results of this subsubsection is made on the basis [6], [29], [30, v. 2].

Applying the operators ${\eta_\Lambda}^{KL}$ to $g_{\Lambda\Psi}(r_I)^\Lambda(r_J)^\Psi = 0$, we obtain

$$\begin{gathered} (R_I)^{AB}(R_J)_{AB} = 0 \Leftrightarrow ((R_I)^{AB} - (R_J)^{AB})((R_K)_{AB} - (R_L)_{AB}) = 0 \Leftrightarrow \\ ((R_I)^{ab} - (R_J)^{ab})((R_K)_{ab} - (R_L)_{ab}) = 0. \end{gathered} \tag{14.148}$$

Here as usual, by I, J, we denote the number of basis points. This defines the system

$$\left\{\begin{array}{lcl} i(R_1)^{ab}Y_b &=& X^a, \\ i(R_2)^{ab}Y_b &=& X^a, \\ i(R_3)^{ab}Y_b &=& X^a, \\ i(R_4)^{ab}Y_b &=& X^a, \end{array}\right. \Leftrightarrow \left\{\begin{array}{lcl} i((R_1)^{ab}-(R_2)^{ab})Y_b &=& 0, \\ i((R_1)^{ab}-(R_3)^{ab})Y_b &=& 0, \\ i((R_3)^{ab}-(R_4)^{ab})Y_b &=& 0, \\ i(R_1)^{ab}Y_b &=& X^a. \end{array}\right. \tag{14.149}$$

Next we consider only the right system. It is constructed as follows. A cospinor Y_a which resets 3 different simple bivectors always exists. This statement is reduced to the existence of a covector orthogonal to three linearly independent vectors since any simple bivector is decomposed by the formula $R^{ab}=2P^{[a}Q^{b]}$. From the fourth equation, the spinor X^a is defined. On the other hand, on the basis that all $(R_I)^{ab}$ have the form $R^{ab}_{\text{particular}}+\lambda_1 S^{[a}X^{b]}+\lambda_2 X^{[a}Z^{b]}+\lambda_3 S^{[a}Z^{b]}$ for fixed $\lambda_1,\lambda_2,\lambda_3$ the equation

$$((R_I)^{ab}-(R_J)^{ab})((R_K)_{ab}-(R_L)_{ab})=0 \tag{14.150}$$

is satisfied.

1. So, let us know the last equation (14.149)

   $$iR_1{}^{ab}Y_b=X^a \tag{14.151}$$

   then we have the 4 equations, all of which will be significant. Since we have the eight unknowns (X^a,Y_b) for fixed $(R_I)^{ab}$ then the point of the quadric $\mathbb{C}\tilde{Q}_6$ uniquely defines the 3-dimensional planar generator $\mathbb{CP}_3(\mathbb{C}Q_6)$.

2. If we know all the equations of (14.149) with the conditions

   $$((R_1)^{ab}-(R_2)^{ab})((R_1)_{ab}-(R_2)_{ab})=0,\quad ((R_1)^{ab}-(R_3)^{ab})((R_1)_{ab}-(R_3)_{ab})=0,$$

   $$((R_3)^{ab}-(R_4)^{ab})((R_3)_{ab}-(R_4)_{ab})=0,$$

   $$(R_1)^{ab}(R_2)_{ab}=0,\quad (R_1)^{ab}(R_3)_{ab}=0,\quad (R_1)^{ab}(R_4)_{ab}=0,$$

   $$(R_2)^{ab}(R_3)_{ab}=0,\quad (R_2)^{ab}(R_4)_{ab}=0,\quad (R_3)^{ab}(R_4)_{ab}=0, \tag{14.152}$$

   that from the 16 equations, only 7 from them will be significant (the 8 unknowns and the 9 communication conditions). Therefore, the generator $\mathbb{CP}_3(\mathbb{C}\tilde{Q}_6)$ will uniquely define the point of the quadric $\mathbb{C}Q_6$.

3. If we know the three equations of the system (14.149)

   $$\left\{\begin{array}{lcl} i((R_1)^{ab}-(R_2)^{ab})Y_b &=& 0, \\ i((R_1)^{ab}-(R_3)^{ab})Y_b &=& 0, \\ i(R_1)^{ab}Y_b &=& X^a \end{array}\right. \tag{14.153}$$

   with the conditions

   $$((R_1)^{ab}-(R_2)^{ab})((R_1)_{ab}-(R_2)_{ab})=0,\quad ((R_1)^{ab}-(R_3)^{ab})((R_1)_{ab}-(R_3)_{ab})=0,$$

   $$(R_1)^{ab}(R_2)_{ab}=0,\quad (R_1)^{ab}(R_3)_{ab}=0,\quad (R_2)^{ab}(R_3)_{ab}=0, \tag{14.154}$$

that from the 12 equations, only 7 from them will be significant (the 8 unknowns and the 5 communication conditions). This means that the generator $\mathbb{CP}_2(\mathbb{C}\tilde{Q}_6)$ will uniquely define the point of the quadric $\mathbb{C}Q_6$. In this case, the manifold of the generators $\mathbb{CP}_2(\mathbb{C}\tilde{Q}_6)$, belonging to a single generator $\mathbb{CP}_3(\mathbb{C}\tilde{Q}_6)$, uniquely defines the point of the quadric $\mathbb{C}Q_6$.

4. If we know the two equations of the system (14.149)

$$\left\{ \begin{array}{lcl} i((R_1)^{ab} - (R_2)^{ab})Y_b & = & 0, \\ i(R_1)^{ab}Y_b & = & X^a \end{array} \right. \tag{14.155}$$

with the conditions

$$((R_1)^{ab} - (R^{ab}_2)((R_1)_{ab} - (R_2)_{ab}) = 0, \quad (R_1)^{ab}(R_2)_{ab} = 0, \tag{14.156}$$

then from the 8 equations, only 6 from them will be significant (the 8 unknowns and the 2 communication conditions). This means that the rectilinear generator $\mathbb{CP}_1$ of the quadric $\mathbb{C}\tilde{Q}_6$ will uniquely define the rectilinear generator $\mathbb{CP}_1$ belonging to the quadric $\mathbb{C}Q_6$. In this case, the manifold of generators $\mathbb{CP}_1(\mathbb{C}\tilde{Q}_6)$, belonging to the same generator $\mathbb{CP}_3(\mathbb{C}\tilde{Q}_6)$, defines the beam of generators $\mathbb{CP}_1(\mathbb{C}Q_6)$ belonging to the quadric $\mathbb{C}Q_6$ (this beam is a cone). The center of the beam is determined by the system (14.149).

14.7.1 On the Majorana spinors.

In this subsubsection, it is told about the Majorana condition in terms of an involution in the spinor space. The conclusion of all results of this subsubsection is made on the basis [21], [22].

As an illustration, we consider the Infeld-van der Waerden symbols [22, v. 1, p. 123 (eng)]. Let the Clifford equation, given on the vector space $\mathbb{C}^4$ in a special basis, have the form

$$\eta_\Lambda \sigma_\Psi + \eta_\Psi \sigma_\Lambda = 2\delta_{\Lambda\Psi}, \quad (\Lambda, \Psi, ... = \overline{0,3}). \tag{14.157}$$

If we specify the real inclusion $H_i{}^\Lambda : \mathbb{R}^4_{(1,3)} \subset \mathbb{C}^4$ then the involution $S_A{}^{A'} := \varepsilon^{A'B'}S_{AB'}$

$$S_\Lambda{}^{\Psi'}\bar{\eta}_{\Psi'}{}^{A'B'} = \eta_\Lambda{}^{BA}S_A{}^{A'}S_B{}^{B'}, \quad (A, A', ... = \overline{1,2}). \tag{14.158}$$

is defined according to Corollary 8.4, Case II). The Infeld-van der Waerden symbols are determined as follows

$$\begin{gathered} g_i{}^{AA'} := H_i{}^\Lambda \eta_\Lambda{}^{AB} S_B{}^{A'}, \\ \overline{g_i{}^{AD'}} = \bar{g}_i{}^{A'D} := \bar{H}_i{}^{\Lambda'}\bar{\eta}_{\Lambda'}{}^{A'B'}\bar{S}_{B'}{}^D = \bar{H}_i{}^{\Lambda'}\bar{S}_{\Lambda'}{}^\Lambda \eta_\Lambda{}^{DC}S_C{}^{A'} = H_i{}^\Lambda \eta_\Lambda{}^{DC}S_C{}^{A'} = g_i{}^{DA'}, \\ \bar{g}_i{}^{A'D} = g_i{}^{DA'}, \quad (i, j, ... = \overline{0,3}). \end{gathered} \tag{14.159}$$

By this equation, the involution $S_B{}^{A'}$ is introduced in the definition of the symbols $g_i{}^{AA'}$. The presence of the metric antisymmetric spin-tensor ε_{AB} allows to carry out such the operation with which we can raise and lower the single index only. We now construct the Dirac operators

$$\gamma_i = \begin{pmatrix} 0 & (g^T)_{iAA'} \\ g_i{}^{BB'} & 0 \end{pmatrix}. \tag{14.160}$$

In a special basis, they will have the form

$$\gamma_0 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad \gamma_1 = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix},$$

$$\gamma_2 = \begin{pmatrix} 0 & 0 & 0 & -i \\ 0 & 0 & i & 0 \\ 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \end{pmatrix}, \quad \gamma_3 = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}, \tag{14.161}$$

$$\begin{gathered} \gamma_i\bar{\gamma}_j + \gamma_j\bar{\gamma}_i = 2g_{ij}, \\ \gamma_i\bar{\gamma}_2 + \gamma_2\bar{\gamma}_i = 0, \quad \gamma_2\bar{\gamma}_2 = -1, \\ \gamma_i\bar{\gamma}_2 + \gamma_2\bar{\gamma}_i = 0, \quad i \neq 2, \\ \bar{\gamma}_2\gamma_i\bar{\gamma}_2 + \bar{\gamma}_2\gamma_2\bar{\gamma}_i = 0, \\ \bar{\gamma}_2\gamma_i\bar{\gamma}_2 = \bar{\gamma}_i, \end{gathered}$$

Accordingly, the involution ${S_A}^{A'}$ will generate the involution S for the complex Dirac operators $\bar{\gamma}_i = \bar{S}\gamma_i\bar{S}$, and in this special basis, it will be executed $||S|| = ||\gamma^2||$. Thus, we can define the Majorana spinor as

$$\psi^i = \gamma^2\bar{\psi}^i \tag{14.162}$$

which corresponds to the presentation [21, p. 98, Appendix E (eng)].

In the transition to the space $\mathbb{R}^6_{(2,4)}$, the Majorana condition will have the form $(A, A', ... = \overline{1,4},\ a, a', ... = \overline{1,2})$ $({\eta_\Lambda}^{AB} = -{\eta_\Lambda}^{BA})$

$$\begin{gathered} {S_\Lambda}^{\Lambda'} {\bar{\eta}_{\Lambda'}}{}^{A'B'} = \eta_{\Lambda AB} \underbrace{(i\tilde{S}^{AA'})}_{:=S^{AA'}} \underbrace{(i\tilde{S}^{BB'})}_{:=S^{BB'}}, \quad S^{AA'} = -\bar{S}^{A'A}, \\ \bar{X}^{A'} = (\bar{X}^{a'}, \bar{Y}_{b'}) = ({\bar{S}^{a'}}{}_a X^a, {\bar{S}_{b'}}{}^b Y_b) = \begin{pmatrix} 0 & {\bar{S}^{a'}}{}_a \\ {\bar{S}_{b'}}{}^b & 0 \end{pmatrix} \begin{pmatrix} Y_b \\ X^a \end{pmatrix} = \bar{S}^{A'A} X_A. \end{gathered} \tag{14.163}$$

However, in such the spinor space according to (8.39), there is no metric tensors, except $S_{AA'}$, capable to raise and to lower single indexes. The Majorana condition for the space $\mathbb{R}^6$ will be the same. In this case, we use the systems $(a, a', ... = \overline{1,4})$

$$\left\{ \begin{array}{l} iR^{ab}Y_b = X^a, \\ iR^{ab}N_b = L^a, \end{array} \right. \quad \left\{ \begin{array}{l} -iR_{ab}Z^b = T_a, \\ -iR_{ab}K^b = M_a. \end{array} \right. \tag{14.164}$$

If our space would be a complex $\mathbb{CR}^6$ then this system would establish a one-to-one correspondence between the respective rectilinear generators of the quadrics. The representation freedom would be dictated by the homogeneous solution of the system which would be isotropic. But, because of the real inclusion $\mathbb{R}^6 \subset \mathbb{CR}^6$, the isotropic vectors are represented only by the zero-vector.

$$\begin{array}{cc} R^{ab} = \frac{-i}{Y^kY_k}(X_dY_c\varepsilon^{abcd} + 2X^{[a}Y^{b]}), & R^{ab} = \frac{-i}{N^kN_k}(L_dN_c\varepsilon^{abcd} + 2L^{[a}N^{b]}), \\ Z^b := Y^b = S^{bb'}\bar{Y}_{b'}, \quad T_a := X_a = S_{aa'}\bar{X}^{a'}, & K^b := N^b = S^{bb'}\bar{N}_{b'}, \quad M_a := L_a = S_{aa'}\bar{L}^{a'}, \\ X^aY_a = 0, & L^aN_a = 0. \end{array} \tag{14.165}$$

Therefore, one can establish the one-to-one correspondence between a point of the quadric $\mathbb{C}\tilde{Q}_6$ and a rectilinear generator $\mathbb{CP}_1(\mathbb{C}Q_6)$. Since on a rectilinear generator of the quadrics $\mathbb{C}Q_6$ (the left system (14.164)), we construct the conjugate rectilinear generator $\mathbb{CP}_1(\mathbb{C}Q_6)$ (the right system (14.164)) then this will define the 3-dimensional planar generator $\mathbb{CP}_3(\mathbb{C}Q_6)$ which will uniquely identify the point to the coordinates R^{ab} on the quadric $\mathbb{C}\tilde{Q}_6$. This is the geometric meaning of the Majorana condition for the inclusion $\mathbb{R}_6 \subset \mathbb{CR}_6$.

$$X^aX_a + Y^aY_a(-\frac{1}{4}R^{ab}R_{ab}) = 0, \quad L^aL_a + N^aN_a(-\frac{1}{4}R^{ab}R_{ab}) = 0. \tag{14.166}$$

(Obviously, $R^{ab}R_{ab} > 0, \quad X^aX_a = S_{aa'}X^a\bar{X}^{a'} > 0, \quad Y^aY_a = S_{aa'}Y^a\bar{Y}^{a'} > 0$ for nonzero vectors.) The condition can be rewritten as follows

$$(X^a, Y_b)(X_a, \frac{-X^dX_d}{Y^cY_c}Y^b) = 0, \quad (L^a, N_b)(L_a, \frac{-L^dL_d}{N^cN_c}N^b) = 0. \tag{14.167}$$

Incidentally, R^{ab} is a null vector of the isotropic cone of the space $R^8_{(1,7)}$. The corresponding quadric cut out by a section of the cone.

Example 14.1. *Thus, let* $Y^kY_k = 2, \quad X^aY_a = 0$ *then in the special basis*

$$\begin{gathered} iR^{12} = -i\ \overline{R^{34}} = X^{[1}Y^{2]} + \overline{X^{[3}Y^{4]}}, \quad iR^{13} = i\ \overline{R^{24}} = X^{[1}Y^{3]} - \overline{X^{[2}Y^{4]}}, \\ iR^{14} = -i\ \overline{R^{23}} = X^{[1}Y^{4]} + \overline{X^{[2}Y^{3]}}. \end{gathered} \tag{14.168}$$

14.7.2 Theorem on two quadrics.

In this subsubsection, it is told about how to interpret the Cartan triality principle on the basis of previous results and to bring them together. The conclusion of all results of this subsubsection is made on the basis [6], [29], [30, v. 2, p. 258 (rus)].

Thus, the theorem is proved.

Theorem 14.3. *(the triality principle for the two B-cylinders).*
In the projective space $\mathbb{CP}_7$*, there are two quadrics (two B - cylinders) with the following main properties*

1. *The planar generator* $\mathbb{CP}_3$ *of a one quadric will define one-to-one the point* R *on the other quadric.*

2. *The planar generator* $\mathbb{CP}_2$ *of a one quadric will uniquely define the point* R *on the other quadric. But the point* R *of the second quadric can be associated to the manifold of planar generators* $\mathbb{CP}_2$ *belonging to the same planar generator* $\mathbb{CP}_3$ *of the first quadric.*

3. *The rectilinear generator* $\mathbb{CP}_1$ *of a one quadric will define one-to-one the rectilinear generator* $\mathbb{CP}_1$ *of the other quadric. And all the rectilinear generators belonging to the same planar generator* $\mathbb{CP}_3$ *of the first quadric define the beam centered at* R *belonging to the second quadric.*

This theorem is actually the generalization of the Klein correspondence. Prove this.

Proof.
On the quadric $\mathbb{C}Q_6$, we consider only those generators which have the form $X^A = (0, Y_b)$. The manifold of such generators is diffeomorphic to $\mathbb{CP}_3$. In this case, each generator can be associated to the point of the quadric $\mathbb{C}Q_4 \subset \mathbb{C}\tilde{Q}_6$. Then $R^{ab}Y_b = 0$. Until the end of the proof, we set $\mathbf{A}, \mathbf{B}, \mathbf{A}', \mathbf{B}', ... = \overline{1, 2}$. In addition, we consider the spinor representation of a twistor according to [22, eq. (6.1.24) and (6.2.18) (eng)].

$$Y_b = (\pi_{\mathbf{B}}, \bar{\omega}^{\mathbf{B}'}),$$

$$R^{ab} = const \begin{pmatrix} -\frac{1}{2}\varepsilon^{\mathbf{AB}} r_c r^c & iR^{\mathbf{A}}{}_{\mathbf{B}'} \\ -i\bar{R}_{\mathbf{A}'}{}^{\mathbf{B}} & \bar{\varepsilon}_{\mathbf{A}'\mathbf{B}'} \end{pmatrix}. \tag{14.169}$$

Therefore, the equation $R^{ab}Y_b = 0$ can be rewritten as a system of two equations

$$\begin{cases} -\frac{1}{2}\varepsilon^{\mathbf{AB}} r_c r^c \pi_{\mathbf{B}} + iR^{\mathbf{A}}{}_{\mathbf{B}'}\bar{\omega}^{\mathbf{B}'} & = & 0, \\ -i\bar{R}_{\mathbf{A}'}{}^{\mathbf{B}}\pi_{\mathbf{B}} + \bar{\varepsilon}_{\mathbf{A}'\mathbf{B}'}\bar{\omega}^{\mathbf{B}'} & = & 0, \end{cases} \tag{14.170}$$

and only one of them will be significant

$$ir^{\mathbf{AA}'}\bar{\pi}_{\mathbf{A}'} = \omega^{\mathbf{A}}. \tag{14.171}$$

Here we use the operation of conjugation and the metric spin-tensor $\bar{\varepsilon}_{\mathbf{A}'\mathbf{B}'}, \varepsilon_{\mathbf{AB}}$ with the help of which spinor indices rise and lover. This spinors pass each other by the conjugation. This defines the system

$$\begin{cases} ir^{\mathbf{AA}'}\bar{\pi}_{\mathbf{A}'} & = & \bar{\omega}^{\mathbf{A}'}, \\ ir^{\mathbf{AA}'}\bar{\eta}_{\mathbf{A}'} & = & \xi^{\mathbf{A}}, \end{cases} \quad Y_b = (\pi_{\mathbf{B}}, \bar{\omega}^{\mathbf{B}'}), \quad T_b = (\eta_{\mathbf{B}}, \bar{\xi}^{\mathbf{B}'}). \tag{14.172}$$

This system coincides with the system [22, eq. (6.2.14)] which, in turn, leads to the Klein correspondence. □

It should be noted in conclusion that from this theorem, the Cartan triality principle implies: There are the 3 diffeomorphic manifolds:

1. the manifold of all points of the quadric;
2. the manifold of I-family maximal planar generators;
3. the manifold of II-family maximal planar generators.

This is true, because the two constructed quadrics can be identified, for example, by means of the spin-tensor $P_\Lambda{}^L$. The manifold of points of the quadric is diffeomorphic to the maximal planar generator manifold. In addition, since the Cartan triality principle is performed then the operators $\eta^i{}_{KL}$ for the inclusion $\mathbb{R}^8 \subset \mathbb{C}^8$ define the octave algebra. This assertion is based on the results given in the monography [22, v. 2, pp. 460-462] which deals with the structure constants of this algebra.

14.8 Explicit construction of the Lie spin-operators for n=4.

In this subsection, it is told about how possible in a certain basis to get a representation of the operators P with which the help we can construct the spinor analog of the Lie operators. The conclusion of all results of this subsection is made on the basis [9], [22, v. 2, pp. 101-103 (eng)].

Let on the space $\mathbb{R}^8_{(4,4)}$, the metric

$$dS^2 = -dU^2 + dS^2 + dV^2 - dW^2 + dX^2 - dY^2 - dT^2 + dZ^2 \tag{14.173}$$

be given then in some basis, the connecting operators ${\eta_\Lambda}^{KL}$ have the form

$$\begin{array}{cccc}
{\eta_3}^{12} = \frac{1}{\sqrt{2}}, & {\eta_3}^{34} = \frac{1}{\sqrt{2}}, & {\eta_4}^{12} = \frac{1}{\sqrt{2}}, & {\eta_4}^{34} = -\frac{1}{\sqrt{2}}, \\
{\eta_3}^{78} = \frac{1}{\sqrt{2}}, & {\eta_3}^{56} = \frac{1}{\sqrt{2}}, & {\eta_4}^{78} = \frac{1}{\sqrt{2}}, & {\eta_4}^{56} = -\frac{1}{\sqrt{2}}, \\
{\eta_7}^{14} = \frac{1}{\sqrt{2}}, & {\eta_7}^{23} = -\frac{1}{\sqrt{2}}, & {\eta_8}^{14} = \frac{1}{\sqrt{2}}, & {\eta_8}^{23} = \frac{1}{\sqrt{2}}, \\
{\eta_7}^{67} = \frac{1}{\sqrt{2}}, & {\eta_7}^{58} = -\frac{1}{\sqrt{2}}, & {\eta_8}^{67} = \frac{1}{\sqrt{2}}, & {\eta_8}^{58} = \frac{1}{\sqrt{2}}, \\
{\eta_6}^{13} = \frac{1}{\sqrt{2}}, & {\eta_6}^{24} = \frac{1}{\sqrt{2}}, & {\eta_5}^{13} = \frac{1}{\sqrt{2}}, & {\eta_5}^{24} = -\frac{1}{\sqrt{2}}, \\
{\eta_6}^{68} = -\frac{1}{\sqrt{2}}, & {\eta_6}^{57} = -\frac{1}{\sqrt{2}}, & {\eta_5}^{68} = -\frac{1}{\sqrt{2}}, & {\eta_5}^{57} = \frac{1}{\sqrt{2}}, \\
{\eta_2}^{15} = \frac{1}{\sqrt{2}}, & {\eta_2}^{51} = \frac{1}{\sqrt{2}}, & {\eta_1}^{15} = -\frac{1}{\sqrt{2}}, & {\eta_1}^{51} = \frac{1}{\sqrt{2}}, \\
{\eta_2}^{26} = \frac{1}{\sqrt{2}}, & {\eta_2}^{62} = \frac{1}{\sqrt{2}}, & {\eta_1}^{26} = -\frac{1}{\sqrt{2}}, & {\eta_1}^{62} = \frac{1}{\sqrt{2}}, \\
{\eta_2}^{37} = \frac{1}{\sqrt{2}}, & {\eta_2}^{73} = \frac{1}{\sqrt{2}}, & {\eta_1}^{37} = -\frac{1}{\sqrt{2}}, & {\eta_1}^{73} = \frac{1}{\sqrt{2}}, \\
{\eta_2}^{48} = \frac{1}{\sqrt{2}}, & {\eta_2}^{84} = \frac{1}{\sqrt{2}}, & {\eta_1}^{48} = -\frac{1}{\sqrt{2}}, & {\eta_1}^{84} = \frac{1}{\sqrt{2}},
\end{array} \tag{14.174}$$

and the significant coordinates of the operators ${\eta^\Lambda}_{KL}$ will be such that

$$\begin{array}{cccc}
{\eta^3}_{12} = \frac{1}{\sqrt{2}}, & {\eta^3}_{34} = \frac{1}{\sqrt{2}}, & {\eta^4}_{12} = \frac{1}{\sqrt{2}}, & {\eta^4}_{34} = -\frac{1}{\sqrt{2}}, \\
{\eta^3}_{78} = \frac{1}{\sqrt{2}}, & {\eta^3}_{56} = \frac{1}{\sqrt{2}}, & {\eta^4}_{78} = \frac{1}{\sqrt{2}}, & {\eta^4}_{56} = -\frac{1}{\sqrt{2}}, \\
{\eta^7}_{14} = \frac{1}{\sqrt{2}}, & {\eta^7}_{23} = -\frac{1}{\sqrt{2}}, & {\eta^8}_{14} = \frac{1}{\sqrt{2}}, & {\eta^8}_{23} = \frac{1}{\sqrt{2}}, \\
{\eta^7}_{67} = \frac{1}{\sqrt{2}}, & {\eta^7}_{58} = -\frac{1}{\sqrt{2}}, & {\eta^8}_{67} = \frac{1}{\sqrt{2}}, & {\eta^8}_{58} = \frac{1}{\sqrt{2}}, \\
{\eta^6}_{13} = \frac{1}{\sqrt{2}}, & {\eta^6}_{24} = \frac{1}{\sqrt{2}}, & {\eta^5}_{13} = \frac{1}{\sqrt{2}}, & {\eta^5}_{24} = -\frac{1}{\sqrt{2}}, \\
{\eta^6}_{68} = -\frac{1}{\sqrt{2}}, & {\eta^6}_{57} = -\frac{1}{\sqrt{2}}, & {\eta^5}_{68} = -\frac{1}{\sqrt{2}}, & {\eta^5}_{57} = \frac{1}{\sqrt{2}}, \\
{\eta^2}_{15} = \frac{1}{\sqrt{2}}, & {\eta^2}_{51} = \frac{1}{\sqrt{2}}, & {\eta^1}_{15} = -\frac{1}{\sqrt{2}}, & {\eta^1}_{51} = \frac{1}{\sqrt{2}}, \\
{\eta^2}_{26} = \frac{1}{\sqrt{2}}, & {\eta^2}_{62} = \frac{1}{\sqrt{2}}, & {\eta^1}_{26} = -\frac{1}{\sqrt{2}}, & {\eta^1}_{62} = \frac{1}{\sqrt{2}}, \\
{\eta^2}_{37} = \frac{1}{\sqrt{2}}, & {\eta^2}_{73} = \frac{1}{\sqrt{2}}, & {\eta^1}_{37} = -\frac{1}{\sqrt{2}}, & {\eta^1}_{73} = \frac{1}{\sqrt{2}}, \\
{\eta^2}_{48} = \frac{1}{\sqrt{2}}, & {\eta^2}_{84} = \frac{1}{\sqrt{2}}, & {\eta^1}_{48} = -\frac{1}{\sqrt{2}}, & {\eta^1}_{84} = \frac{1}{\sqrt{2}}.
\end{array} \tag{14.175}$$

Therefore, in the same basis,

$$\begin{array}{cc}
\sqrt{2} \parallel {P_{\tilde{A}}}^{\tilde{\tilde{B}}} \parallel = & \sqrt{2} \parallel {P^{\tilde{A}}}_{\tilde{\tilde{B}}} \parallel = \\
\begin{pmatrix}
1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\
0 & 1 & 0 & 0 & 0 & -1 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & -1 & 0 \\
0 & 0 & 0 & 1 & 0 & 0 & 0 & -1 \\
0 & 0 & 0 & 1 & 0 & 0 & 0 & 1
\end{pmatrix}, &
\begin{pmatrix}
1 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\
0 & 1 & 0 & 0 & 0 & -1 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & -1 & 0 \\
0 & 0 & 0 & 1 & 0 & 0 & 0 & -1 \\
0 & 0 & 0 & 1 & 0 & 0 & 0 & 1
\end{pmatrix}.
\end{array} \tag{14.176}$$

Hence, Algorithm 8.1 will take the form

Algorithm 14.1.

1. On the first step, as the operators (6.24) and (6.30), it is used the operators

$$\underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} e^{i\frac{\pi}{8}}\delta^{a}{}_{\tilde{\tilde{a}}} & 0 & \pm ie^{i\frac{\pi}{8}}\varepsilon^{a\tilde{\tilde{p}}} & 0 \\ 0 & e^{-i\frac{\pi}{8}}\delta_{c}{}^{\tilde{\tilde{c}}} & 0 & \mp e^{-i\frac{\pi}{8}}\varepsilon_{c\tilde{\tilde{q}}} \end{pmatrix}}_{:=(\tilde{\tilde{m}}_{1_\pm})^{A}{}_{\tilde{\tilde{A}}}}\cdot$$

$$\cdot\underbrace{\begin{pmatrix} \alpha_1\varepsilon^{\tilde{\tilde{a}}\tilde{\tilde{b}}} & \eta_\alpha{}^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{d}}} & \gamma_1\delta^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{k}}} & 0 \\ -(\eta^T)_{\alpha\tilde{\tilde{c}}}{}^{\tilde{\tilde{b}}} & \beta_1\varepsilon_{\tilde{\tilde{c}}\tilde{\tilde{d}}} & 0 & \gamma_1\delta_{\tilde{\tilde{c}}}{}^{\tilde{\tilde{l}}} \\ \delta_1\delta_{\tilde{\tilde{p}}}{}^{\tilde{\tilde{b}}} & 0 & \beta_1\varepsilon_{\tilde{\tilde{p}}\tilde{\tilde{k}}} & \eta_{\alpha\tilde{\tilde{p}}}{}^{\tilde{\tilde{l}}} \\ 0 & \delta_1\delta^{\tilde{\tilde{q}}}{}_{\tilde{\tilde{d}}} & -(\eta^T)_\alpha{}^{\tilde{\tilde{q}}}{}_{\tilde{\tilde{k}}} & \alpha_1\varepsilon^{\tilde{\tilde{q}}\tilde{\tilde{l}}} \end{pmatrix}}_{\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}}\underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} 0 & e^{i\frac{\pi}{8}}\varepsilon_{\tilde{\tilde{b}}d} \\ -e^{-i\frac{\pi}{8}}\varepsilon^{\tilde{\tilde{d}}b} & 0 \\ 0 & \pm e^{i\frac{\pi}{8}}\delta^{\tilde{\tilde{k}}}{}_{d} \\ \mp e^{-i\frac{\pi}{8}}i\delta_{\tilde{\tilde{l}}}{}^{b} & 0 \end{pmatrix}}_{:=(\tilde{m}^T_{1_\pm})_{\tilde{\tilde{B}}}{}^{B}}=$$

$$=\underbrace{\begin{pmatrix} \eta_\alpha{}^{ab} & \underbrace{\frac{e^{i\frac{\pi}{4}}}{2}((-\alpha_1-i\beta_1)\pm(\gamma_1-i\delta_1))}_{:=(\phi_{1_\pm})}\delta^{a}{}_{d} \\ \underbrace{\frac{ie^{-i\frac{\pi}{4}}}{2}((-\alpha_1-i\beta_1)\mp(\gamma_1-i\delta_1))}_{:=(\psi_{1_\pm})}\delta_{c}{}^{b} & -(\eta^T)_{\alpha cd} \end{pmatrix}}_{:=\eta_{1_\pm}{}_{\tilde{\Lambda}}{}^{AB}},$$

$$\alpha_1=\tfrac{1}{2}(\eta_3+\eta_4),\quad \beta_1=\tfrac{1}{2}(\eta_3-\eta_4),\quad \gamma_1=\tfrac{1}{2}(\eta_2-\eta_1),\quad \delta_1=\tfrac{1}{2}(\eta_2+\eta_1). \tag{14.177}$$

Here, $\varepsilon^{\tilde{\tilde{a}}\tilde{\tilde{b}}}$ *is the antisymmetric metric spin-tensor for the connecting operators* $\eta_{\alpha\tilde{\tilde{p}}}{}^{\tilde{\tilde{l}}}$ *(*$a,b,...,\tilde{\tilde{a}},\tilde{\tilde{b}},...=\overline{1,2},\ \tilde{\Lambda},...,\tilde{\tilde{A}},\tilde{\tilde{B}},...=\overline{1,8},\ A,B,...=\overline{1,4}$*).*

2. On the second step, as the operators (6.24) and (6.30), it is used the operators

$$\underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} e^{i\frac{\pi}{8}} & \pm ie^{i\frac{\pi}{8}} & 0 & 0 \\ 0 & 0 & e^{-i\frac{\pi}{8}} & \pm e^{-i\frac{\pi}{8}} \end{pmatrix}}_{:=(\tilde{\tilde{m}}_{2_\pm})^{A}{}_{\tilde{\tilde{A}}}}\cdot$$

$$\cdot\underbrace{\begin{pmatrix} \alpha_2 & \gamma_2 & \phi_{1_\pm} & 0 \\ \delta_2 & \beta_2 & 0 & \phi_{1_\pm} \\ \psi_{1_\pm} & 0 & -\beta_2 & \gamma_2 \\ 0 & \psi_{1_\pm} & \delta_2 & -\alpha_2 \end{pmatrix}}_{(\eta_{1_\pm})_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}}\underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} e^{i\frac{\pi}{8}} & 0 \\ \pm e^{i\frac{\pi}{8}} & 0 \\ 0 & e^{-i\frac{\pi}{8}} \\ 0 & \mp ie^{-i\frac{\pi}{8}} \end{pmatrix}}_{:=(\tilde{m}^T_{2_\pm})_{\tilde{\tilde{B}}}{}^{B}}=$$

$$=\underbrace{\begin{pmatrix} \underbrace{\frac{e^{i\frac{\pi}{4}}}{2}((\alpha_2+i\beta_2)\pm(\gamma_2+i\delta_2))}_{:=\phi_{2_\pm}} & \underbrace{\frac{e^{i\frac{\pi}{4}}}{2}((-\alpha_1-i\beta_1)\pm(\gamma_1-i\delta_1))}_{:=\phi_{1_\pm}} \\ \underbrace{\frac{ie^{-i\frac{\pi}{4}}}{2}((-\alpha_1-i\beta_1)\mp(\gamma_1-i\delta_1))}_{:=\psi_{1_\pm}} & \underbrace{\frac{ie^{-i\frac{\pi}{4}}}{2}((\alpha_2+i\beta_2)\mp(\gamma_2+i\delta_2))}_{:=\psi_{2_\pm}} \end{pmatrix}}_{:=(\eta_{1,2,3,4})_{\tilde{\Lambda}}{}^{AB}},$$

$$\alpha_2=\tfrac{1}{2}(\eta_7+\eta_8),\quad \beta_2=\tfrac{1}{2}(\eta_7-\eta_8),\quad \gamma_2=\tfrac{1}{2}(-\eta_6-\eta_5),\quad \delta_2=\tfrac{1}{2}(\eta_6-\eta_5). \tag{14.178}$$

$(\tilde{\Lambda},...=\overline{1,8},\ \tilde{\tilde{A}},\tilde{\tilde{B}},...=\overline{1,4},\ A,B,...=\overline{1,2})$.
3. For the tangent space, the operators ${m_{\Lambda}}^{\tilde{\Lambda}}$ have the form

$$\begin{aligned} {m_{\Lambda}}^{\tilde{\Lambda}} &:= \tfrac{1}{\sqrt{2}}\begin{pmatrix} -i & 1 & 0 & 0 & 0 & 0 & 0 & 0\\ 0 & 0 & -i & -1 & 0 & 0 & 0 & 0\\ 0 & 0 & 0 & 0 & i & 1 & 0 & 0\\ 0 & 0 & 0 & 0 & 0 & 0 & i & 1 \end{pmatrix}, \\ {m^{\Lambda}}_{\tilde{\Lambda}} &:= \tfrac{1}{\sqrt{2}}\begin{pmatrix} i & 1 & 0 & 0 & 0 & 0 & 0 & 0\\ 0 & 0 & i & -1 & 0 & 0 & 0 & 0\\ 0 & 0 & 0 & 0 & -i & 1 & 0 & 0\\ 0 & 0 & 0 & 0 & 0 & 0 & -i & 1 \end{pmatrix}. \end{aligned} \tag{14.179}$$

Then the operators

$$\underbrace{\begin{pmatrix} \frac{1}{2}(\underbrace{\frac{1}{\sqrt{2}}(\eta_8+i\eta_7)}_{:=(\eta^{\mathbb{C}})_4}\mp\underbrace{\frac{1}{\sqrt{2}}(\eta_6+i\eta_5)}_{:=(\eta^{\mathbb{C}})_3}) & \frac{1}{2}(\underbrace{\frac{1}{\sqrt{2}}(-\eta_4-i\eta_3)}_{:=(\eta^{\mathbb{C}})_2}\pm\underbrace{\frac{1}{\sqrt{2}}(\eta_2-i\eta_1)}_{:=(\eta^{\mathbb{C}})_1}) \\ \frac{1}{2}(\underbrace{\frac{1}{\sqrt{2}}(-\eta_4-i\eta_3)}_{:=(\eta^{\mathbb{C}})_2}\mp\underbrace{\frac{1}{\sqrt{2}}(\eta_2-i\eta_1)}_{:=(\eta^{\mathbb{C}})_1}) & \frac{1}{2}(\underbrace{\frac{1}{\sqrt{2}}(\eta_8+i\eta_7)}_{:=(\eta^{\mathbb{C}})_4}\pm\underbrace{\frac{1}{\sqrt{2}}(\eta_6+i\eta_5)}_{:=(\eta^{\mathbb{C}})_3}) \end{pmatrix}}_{:={(\eta_{1,2,3,4})_{\Lambda}}^{AB}} \tag{14.180}$$

will be the connecting operators for the complex representation (n=4). Thus, it is constructed the transition from the real representation of the connecting operators for n = 8 to their complex counterparts for n = 4.

Thus, the operators $2({(\tilde{M}^*_K)^T)_{\tilde{\tilde{A}}}}^{A}$, according to (6.33) and Algorithm 8.1, have the form

$$\begin{pmatrix} 0 & 0 & 0 & e^{i\frac{\pi}{8}}\\ 0 & 0 & -e^{i\frac{\pi}{8}} & 0\\ 0 & -e^{-i\frac{\pi}{8}} & 0 & 0\\ e^{-i\frac{\pi}{8}} & 0 & 0 & 0\\ 0 & 0 & p_1e^{i\frac{\pi}{8}} & 0\\ 0 & 0 & 0 & p_1e^{i\frac{\pi}{8}}\\ -p_1ie^{-i\frac{\pi}{8}} & 0 & 0 & 0\\ 0 & -p_1ie^{-i\frac{\pi}{8}} & 0 & 0 \end{pmatrix} \begin{pmatrix} e^{i\frac{\pi}{8}} & 0\\ p_2e^{i\frac{\pi}{8}} & 0\\ 0 & e^{-i\frac{\pi}{8}}\\ 0 & -p_2ie^{-i\frac{\pi}{8}} \end{pmatrix} = \begin{pmatrix} 0 & -p_2i\\ 0 & -1\\ -p_2 & 0\\ 1 & 0\\ 0 & p_1\\ 0 & -p_1p_2i\\ -p_1i & 0\\ -p_1p_2i & 0 \end{pmatrix}. \tag{14.181}$$

In this case, p_1,p_2 are taking one of the two values ±1 providing the 4 different variants of $\tilde{M}^*_K$. It remains to apply these operators and the operators ${m_{\Lambda}}^{\tilde{\Lambda}}$, ${\bar{m}_{\Lambda'}}^{\tilde{\Lambda}}$ to the operators ${P_{\tilde{\Lambda}}}^{\tilde{\tilde{A}}}$

$${(P^*_K)_{\Lambda}}^{A} := \tfrac{1+i}{4}\begin{pmatrix} 0 & p_1-p_2\\ 0 & 1-p_1p_2\\ p_1-p_2 & 0\\ 1-p_1p_2 & 0 \end{pmatrix},\quad {(\bar{P}^*_K)_{\Lambda'}}^{A} := \tfrac{1-i}{4}\begin{pmatrix} 0 & p_1+p_2\\ 0 & 1+p_1p_2\\ -p_1-p_2 & 0\\ 1+p_1p_2 & 0 \end{pmatrix}, \tag{14.182}$$

$$(P_K)^{\Lambda}{}_{A} := \tfrac{1-i}{4}\begin{pmatrix} 0 & p_1-p_2 \\ 0 & 1-p_1p_2 \\ p_1-p_2 & 0 \\ 1-p_1p_2 & 0 \end{pmatrix}, \quad (\bar{P}_K)^{\Lambda'}{}_{A} := \tfrac{1+i}{4}\begin{pmatrix} 0 & p_1+p_2 \\ 0 & 1+p_1p_2 \\ -p_1-p_2 & 0 \\ 1+p_1p_2 & 0 \end{pmatrix}. \tag{14.183}$$

Note 14.2. *$(P^*_K)_{\Lambda}{}^{A} \neq (P_K)_{\Lambda}{}^{A} = g_{\Lambda\Psi}(P_K)^{\Lambda}{}_{B}\varepsilon^{AB}$ where ε^{AB} is the antisymmetric metric spinor.*

It is obvious that for the 4 variants $K = \overline{1,4}$, the two will be significantly only: $(P^*_2)_{\Lambda}{}^{A}$, $(P^*_3)_{\Lambda}{}^{A}$ and $(\bar{P}^*_1)_{\Lambda'}{}^{A}$, $(\bar{P}^*_4)_{\Lambda'}{}^{A}$ $(\Lambda, \ldots = \overline{1,4})$. Under the condition of the ordinary and covariant differentiation constancy of such the operators, the connection coefficients $\Gamma_{\tilde{\Lambda}\Psi}{}^{\Phi}$ decay into the 8 significant components

$$\Gamma_{\tilde{\Lambda}\Psi}{}^{\Phi} \to \begin{cases} (\Gamma_K)_{\Lambda A}{}^{B} := \Gamma_{\Lambda\Psi}{}^{\Phi}(P_K)^{\Psi}{}_{A}(P^*_K)_{\Phi}{}^{B}, \\ (\Gamma_K)_{\Lambda A'}{}^{B'} := \Gamma_{\Lambda\Psi}{}^{\Phi}(P_K)^{\Psi}{}_{A'}(P^*_K)_{\Phi}{}^{B'}, \\ (\bar{\Gamma}_K)_{\Lambda' A'}{}^{B'} := \bar{\Gamma}_{\Lambda'\Psi'}{}^{\Phi'}(\bar{P}_K)^{\Psi'}{}_{A'}(\bar{P}^*_K)_{\Phi'}{}^{B'}, \\ (\bar{\Gamma}_K)_{\Lambda' A}{}^{B} := \bar{\Gamma}_{\Lambda'\Psi'}{}^{\Phi'}(\bar{P}_K)^{\Psi'}{}_{A}(\bar{P}^*_K)_{\Phi'}{}^{B}. \end{cases} \tag{14.184}$$

In turn, the coefficients $\Gamma_{\tilde{\Lambda}\Psi}{}^{\Phi}$ are defined by the equation $\nabla_{\tilde{\Lambda}} m_{\Psi}{}^{\tilde{\Psi}} = 0$. Thus, Theorems 11.2, 11.3 remain valid in this case. In the case of a real inclusion, with the help of the operators $H_i{}^{\Lambda}$, under the condition of the covariant and ordinary differentiation constancy of the operators $(P^*_2)_i{}^{A}$, $(P^*_3)_i{}^{A}$ and $(P^*_1)_i{}^{A}$, $(P^*_4)_i{}^{A}$, the connection coefficients $\Gamma_{\tilde{\Lambda}i}{}^{j}$ decay into the 4 significant components

$$\Gamma_{\tilde{\Lambda}i}{}^{j} \to \begin{cases} (\Gamma_K)_{kA}{}^{B} := \Gamma_{ki}{}^{j}(P_K)^{i}{}_{A}(P^*_K)_{j}{}^{B}, \\ (\bar{\Gamma}_K)_{kA'}{}^{B'} := \Gamma_{ki}{}^{j}(\bar{P}_K)^{i}{}_{A'}(\bar{P}^*_K)_{j}{}^{B'}. \end{cases} \tag{14.185}$$

However, the real condition restricts the choice of the spinor $X^{\tilde{\tilde{A}}}$ in the determination of the operators $P_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}}$

$$X^{\tilde{\tilde{A}}} = \tilde{S}_{\tilde{\tilde{B}}}{}^{\tilde{\tilde{A}}} X^{\tilde{\tilde{B}}}, \quad X^{\tilde{\tilde{A}}} = \tilde{\tilde{S}}^{\tilde{\tilde{B}}\tilde{\tilde{A}}} X_{\tilde{\tilde{B}}}, \tag{14.186}$$

where $\tilde{S}_{\tilde{\tilde{B}}}{}^{\tilde{\tilde{A}}}$ $(S^{\tilde{\tilde{B}}\tilde{\tilde{A}}})$ is a real representation of the complex involution $\tilde{S}_B{}^{A'}$ $(\tilde{S}^{BA'})$ which is the image of the involution $S_{\Lambda}{}^{\Psi'}$ in the tangent space by Corollary 8.4. We need to add that there is the non-special orthogonal transformation

$$(-\eta_{\Lambda}\eta^{\Psi} + \delta_{\Lambda}{}^{\Psi})\eta_{\Psi}{}^{AB} = \eta_{\Lambda}{}^{BA} \tag{14.187}$$

performing the transposition of the connecting operators (14.85).

Example 14.2. *Consider the inclusion $\mathbb{R}^4_{(1,3)} \subset \mathbb{C}^4$ with*

$$\bar{H}_i{}^{\Lambda'} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -i \end{pmatrix}, \quad H^i{}_{\Lambda} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -i \end{pmatrix}, \quad S_{\Lambda}{}^{\Lambda'} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}. \tag{14.188}$$

For the values of the complex representation operators from Algorithm 14.1, the metric takes the form

$$g_{\Lambda\Psi} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad g_{ij} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}. \tag{14.189}$$

Then the definition (14.182) will have the form

$$(P^*_K)_i{}^A := \tfrac{1+i}{4} \begin{pmatrix} 0 & p_1 - p_2 \\ 0 & 1 - p_1 p_2 \\ p_1 - p_2 & 0 \\ i(1 - p_1 p_2) & 0 \end{pmatrix}, \quad (P_K)^i{}_A := \tfrac{1-i}{4} \begin{pmatrix} 0 & p_1 - p_2 \\ 0 & 1 - p_1 p_2 \\ p_1 - p_2 & 0 \\ -i(1 - p_1 p_2) & 0 \end{pmatrix}. \tag{14.190}$$

Since $x^i := (P_2)^i{}_A (X_2)^A + (P_3)^i{}_A (X_3)^A$ *then from the equations (6.33), (14.84) and Corollary (8.3) imply that* $(X_K)^A := x^i (P^*_K)_i{}^A$, $(X^*_K)_A := x^i (P_K)_{iA}$*; then from the equations (6.33), (14.84), Corollary (8.3) and the condition* $(M_K)^{iA}(M_K)_{iA} = 0$, $(M^*_K)^{iA}(M^*_K)_{iA} = 0$ *(see (14.181)) imply that* $(X^*_2)_C = (P_2)^i{}_C (P_3)_{iB} (X_3)^B$. *Therefore, we can construct the one-to-one mapping* $(a, b, ... = \overline{1,4})$

$$x^i \cdot \underbrace{((P^*_2)_i{}^A, (P^*_3)_i{}^B)}_{:=P_i{}^a} = \underbrace{((X_2)^A, (X_3)^B)}_{:=X^a}, \quad x^i \cdot \underbrace{((P_2)_{iA}, (P_3)_{iB})}_{:=P_{ia}} = \underbrace{((X^*_2)_A, (X^*_3)_B)}_{:=X_a}. \tag{14.191}$$

Then on the spin-pair $X^a = ((X_2)^A, (X_3)^B)$*, the metric and the involution of the form*

$$\begin{aligned} \varepsilon_{ab} &= \begin{pmatrix} 0 & (P_2)^i{}_A (P_3)_{iB} \\ (P_3)^i{}_C (P_2)_{iD} & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & i \\ -i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{pmatrix}, \\ S_a{}^{b'} &= \begin{pmatrix} (P_2)^i{}_A (\bar{P}^*_2)_i{}^{C'} & (P_2)^i{}_A (\bar{P}^*_3)_i{}^{D'} \\ (P_3)^i{}_B (\bar{P}^*_2)_i{}^{C'} & (P_3)^i{}_A (\bar{P}^*_3)_i{}^{D'} \end{pmatrix} = \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & -i \end{pmatrix} \end{aligned} \tag{14.192}$$

are defined. Therefor, it is possible to construct Lie pair-spin operators

$$L_x(Y)^a = x^i \nabla_i Y^a - P^j{}_b Y^b P_i{}^a \nabla_j x^i, \quad L_X(Y)^a = X^c \nabla_c Y^a - Y^c \nabla_c X^a. \tag{14.193}$$

Note, that the expansion $x^i := (P_2)^i{}_A (X_2)^A + (P_3)^i{}_A (X_3)^A$ *is an one of the vector into the two isotropic vectors. And for the isotropic vectors,* Lie spinor operators *were constructed in [22, v. 2, pp. 101-103 (eng)].*

The restricted Lorentz transformation converts the vector x^i *to the vector* $\tilde{x}^i$*. This induces the spinor transformation:* $(X_K)^A \longmapsto (\tilde{X}_K)^A$*. The spinor transformations act on the operators* $(P^*_K)_i{}^A$, $(P_K)^i{}_A$*, more precisely, on the controlling spinor* $X^{\tilde{\tilde{A}}}$ *appearing in the definition (11.59) of such the operators. In turn, this induces the spin-pair transformation of the spin-pair* $X^a = ((X_2)^A, (X_3)^B) \longmapsto \tilde{X}^a$.

15 Acknowledgments

This article consists of the answers to frequently asked questions. Therefore, the author is grateful to all those who has truly formulated these questions that has given the chance to find the answer on them. Special thanks to K. Scharnhorst (for the copies of articles, in particular), without who this publication was not.

16 Appendix

16.1 Proof of the metric tensor and involution relations

We need to prove the identity (8.19).
Case I)., the metric tensor $\varepsilon_{AB} = -\varepsilon_{BA}$.

$$\begin{gathered}
{\eta_\Lambda}^{A_1 B_1} \varepsilon_{A_1 A_{(n/2+q+1)}} \varepsilon_{B_1 B_{(n/2+q+1)}} = \\
= {\eta_\Lambda}^{A_1 B_1} \big(\varepsilon_{(n/2+q)}\big)_{A_{(n/2+q)} A_{(n/2+q+1)}} \ldots \big(\varepsilon_1\big)_{A_1 A_2} \big(\varepsilon_2\big)^{A_3 A_2} \\
\big(\varepsilon_{(n/2+q)}\big)_{B_{(n/2+q)} B_{(n/2+q+1)}} \ldots \big(\varepsilon_1\big)_{B_1 B_2} \big(\varepsilon_2\big)^{B_3 B_2} = \\
= \big(-\eta_{\Lambda B_2 A_2} + (\eta_1)_\Lambda \varepsilon_{1 A_2 B_2}\big) \big(\varepsilon_{(n/2+q)}\big)_{A_{(n/2+q)} A_{(n/2+q+1)}} \ldots \big(\varepsilon_2\big)^{A_3 A_2} \\
\big(\varepsilon_{(n/2+q)}\big)_{B_{(n/2+q)} B_{(n/2+q+1)}} \ldots \big(\varepsilon_2\big)^{B_3 B_2} = \\
= -\big(-{\eta_\Lambda}^{A_3 B_3} + (\eta_2)_\Lambda {\varepsilon_2}^{A_3 B_3}\big) \big(\varepsilon_{(n/2+q)}\big)_{A_{(n/2+q)} A_{(n/2+q+1)}} \ldots \\
\big(\varepsilon_{(n/2+q)}\big)_{B_{(n/2+q)} B_{(n/2+q+1)}} \ldots + (\eta_1)_\Lambda (\varepsilon_1)_{A_{(n/2+q+1)} B_{(n/2+q+1)}} = \\
= {\eta_\Lambda}^{A_3 B_3} \big(\varepsilon_{(n/2+q)}\big)_{A_{(n/2+q)} A_{(n/2+q+1)}} \ldots \big(\varepsilon_{(n/2+q)}\big)_{B_{(n/2+q)} B_{(n/2+q+1)}} \ldots + \\
+ \sum_{Q=1}^{2} (\eta_Q)_\Lambda (\varepsilon_Q)_{A_{(n/2+q+1)} B_{(n/2+q+1)}} = \\
= -\eta_{\Lambda B_{(n/2+q+1)} A_{(n/2+q+1)}} + \sum_{Q=1}^{n/2+q} (\eta_Q)_\Lambda (\varepsilon_Q)_{A_{(n/2+q+1)} B_{(n/2+q+1)}} = \\
= \eta_{\Lambda A_{(n/2+q+1)} B_{(n/2+q+1)}}.
\end{gathered} \tag{16.1}$$

Case I)., the metric tensor $\tilde{\varepsilon}_{AB} = \tilde{\varepsilon}_{BA}$.

$$\begin{gathered}
{\eta_\Lambda}^{A_1 B_1} \tilde{\varepsilon}_{A_1 A_{(n/2+\tilde{q}+1)}} \tilde{\varepsilon}_{B_1 B_{(n/2+\tilde{q}+1)}} = \\
= {\eta_\Lambda}^{A_1 B_1} \big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)_{A_{(n/2+\tilde{q})} A_{(n/2+\tilde{q}+1)}} \ldots \big(\varepsilon_{\tilde{1}}\big)_{A_1 A_2} \big(\varepsilon_{\tilde{2}}\big)^{A_3 A_2} \\
\big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)_{B_{(n/2+\tilde{q})} B_{(n/2+\tilde{q}+1)}} \ldots \big(\varepsilon_{\tilde{1}}\big)_{B_1 B_2} \big(\varepsilon_{\tilde{2}}\big)^{B_3 B_2} = \\
= \big(\eta_{\Lambda B_2 A_2} + (\eta_{\tilde{1}})_\Lambda \varepsilon_{\tilde{1} A_2 B_2}\big) \big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)_{A_{(n/2+\tilde{q})} A_{(n/2+\tilde{q}+1)}} \ldots \big(\varepsilon_{\tilde{2}}\big)^{A_3 A_2} \\
\big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)_{B_{(n/2+\tilde{q})} B_{(n/2+\tilde{q}+1)}} \ldots \big(\varepsilon_{\tilde{2}}\big)^{B_3 B_2} = \\
= \big({\eta_\Lambda}^{A_3 B_3} + (\eta_{\tilde{2}})_\Lambda {\varepsilon_{\tilde{2}}}^{B_3 A_3}\big) \big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)_{A_{(n/2+\tilde{q})} A_{(n/2+\tilde{q}+1)}} \ldots \big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)_{B_{(n/2+\tilde{q})} B_{(n/2+\tilde{q}+1)}} \ldots + \\
+ (\eta_{\tilde{1}})_\Lambda (\varepsilon_{\tilde{1}})_{A_{(n/2+\tilde{q}+1)} B_{(n/2+\tilde{q}+1)}} = \\
= {\eta_\Lambda}^{A_3 B_3} \big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)_{A_{(n/2+\tilde{q})} A_{(n/2+\tilde{q}+1)}} \ldots \big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)_{B_{(n/2+\tilde{q})} B_{(n/2+\tilde{q}+1)}} \ldots + \\
+ \sum_{\tilde{Q}=1}^{2} (\eta_{\tilde{Q}})_\Lambda (\varepsilon_{\tilde{Q}})_{A_{(n/2+\tilde{q}+1)} B_{(n/2+\tilde{q}+1)}} = \\
= \eta_{\Lambda B_{(n/2+\tilde{q}+1)} A_{(n/2+\tilde{q}+1)}} + \sum_{\tilde{Q}=1}^{n/2+\tilde{q}} (\eta_{\tilde{Q}})_\Lambda (\varepsilon_{\tilde{Q}})_{A_{(n/2+\tilde{q}+1)} B_{(n/2+\tilde{q}+1)}} = \\
= \eta_{\Lambda A_{(n/2+\tilde{q}+1)} B_{(n/2+\tilde{q}+1)}}.
\end{gathered} \tag{16.2}$$

Case II)., the involution ${E_A}^B {E_B}^C = -{\delta_A}^C$.

$$\begin{gathered}
{\eta_\Lambda}^{A_1B_1} {E_{A_1}}^{A_{(n/2+q)}} {E_{B_1}}^{B_{(n/2+q+1)}} = \\
= {\eta_\Lambda}^{A_1B_1} (\varepsilon_{(n/2+q)})^{A_{(n/2+q)}A_{(n/2+q+1)}} ... (\varepsilon_1)_{A_1A_2} (\varepsilon_2)^{A_3A_2} \\
(\varepsilon_{(n/2+q)})^{B_{(n/2+q)}B_{(n/2+q+1)}} ... (\varepsilon_1)_{B_1B_2} (\varepsilon_2)^{B_3B_2} = \\
= (-\eta_{\Lambda B_2A_2} + (\eta_1)_\Lambda \varepsilon_{1A_2B_2})(\varepsilon_{(n/2+q)})^{A_{(n/2+q)}A_{(n/2+q+1)}} ... (\varepsilon_2)^{A_3A_2} \\
(\varepsilon_{(n/2+q)})^{B_{(n/2+q)}B_{(n/2+q+1)}} ... (\varepsilon_2)^{B_3B_2} = \\
= -(-{\eta_\Lambda}^{A_3B_3} + (\eta_2)_\Lambda {\varepsilon_2}^{A_3B_3})(\varepsilon_{(n/2+q)})^{A_{(n/2+q)}A_{(n/2+q+1)}} ... (\varepsilon_{(n/2+q)})^{B_{(n/2+q)}B_{(n/2+q+1)}} ... - \\
-(\eta_1)_\Lambda (\varepsilon_1)^{A_{(n/2+q+1)}B_{(n/2+q+1)}} = \\
= {\eta_\Lambda}^{A_3B_3} (\varepsilon_{(n/2+q)})^{A_{(n/2+q)}A_{(n/2+q+1)}} ... (\varepsilon_{(n/2+q)})^{B_{(n/2+q)}B_{(n/2+q+1)}} ... - \\
- \sum_{Q=1}^{2} (\eta_Q)_\Lambda (\varepsilon_Q)^{A_{(n/2+q+1)}B_{(n/2+q+1)}} = \\
= {\eta_\Lambda}^{A_{(n/2+q+1)}B_{(n/2+q+1)}} - \sum_{Q=1}^{n/2+q} (\eta_Q)_\Lambda (\varepsilon_Q)^{A_{(n/2+q+1)}B_{(n/2+q+1)}} = \\
= -{\eta_\Lambda}^{B_{(n/2+q+1)}A_{(n/2+q+1)}}.
\end{gathered} \tag{16.3}$$

Case II)., the involution ${\tilde{E}_A}{}^B {\tilde{E}_B}{}^C = -{\delta_A}^C$.

$$\begin{gathered}
{\eta_\Lambda}^{A_1B_1} {\tilde{E}_{A_1}}{}^{A_{(n/2+\tilde{q}+1)}} {\tilde{E}_{B_1}}{}^{B_{(n/2+\tilde{q}+1)}} = \\
= {\eta_\Lambda}^{A_1B_1} (\varepsilon_{\widetilde{(n/2+\tilde{q})}})^{A_{(n/2+\tilde{q}+1)}A_{(n/2+\tilde{q})}} ... (\varepsilon_{\tilde{1}})_{A_1A_2} (\varepsilon_{\tilde{2}})^{A_3A_2} \\
(\varepsilon_{\widetilde{(n/2+\tilde{q})}})^{B_{(n/2+\tilde{q}+1)}B_{(n/2+\tilde{q})}} ... (\varepsilon_{\tilde{1}})_{B_1B_2} (\varepsilon_{\tilde{2}})^{B_3B_2} = \\
= (\eta_{\Lambda B_2A_2} + (\eta_{\tilde{1}})_\Lambda \varepsilon_{\tilde{1}A_2B_2})(\varepsilon_{\widetilde{(n/2+\tilde{q})}})^{A_{(n/2+\tilde{q}+1)}A_{(n/2+\tilde{q})}} ... (\varepsilon_{\tilde{2}})^{A_3A_2} \\
(\varepsilon_{\widetilde{(n/2+\tilde{q})}})^{B_{(n/2+\tilde{q}+1)}B_{(n/2+\tilde{q})}} ... (\varepsilon_{\tilde{2}})^{B_3B_2} = \\
= ({\eta_\Lambda}^{A_3B_3} + (\eta_{\tilde{2}})_\Lambda {\varepsilon_{\tilde{2}}}^{B_3A_3})(\varepsilon_{\widetilde{(n/2+\tilde{q})}})^{A_{(n/2+\tilde{q}+1)}A_{(n/2+\tilde{q})}} ... (\varepsilon_{\widetilde{(n/2+\tilde{q})}})^{B_{(n/2+\tilde{q}+1)}B_{(n/2+\tilde{q})}} ... + \\
+(\eta_{\tilde{1}})_\Lambda (\varepsilon_{\tilde{1}})_{A_{(n/2+\tilde{q}+1)}B_{(n/2+\tilde{q}+1)}} = \\
= {\eta_\Lambda}^{A_3B_3} (\varepsilon_{\widetilde{(n/2+\tilde{q})}})^{A_{(n/2+\tilde{q}+1)}A_{(n/2+\tilde{q})}} ... (\varepsilon_{\widetilde{(n/2+\tilde{q})}})^{B_{(n/2+\tilde{q}+1)}B_{(n/2+\tilde{q})}} ... + \\
+ \sum_{\tilde{Q}=1}^{2} (\eta_{\tilde{Q}})_\Lambda (\varepsilon_{\tilde{Q}})^{A_{(n/2+\tilde{q}+1)}B_{(n/2+\tilde{q}+1)}} = \\
= {\eta_\Lambda}^{A_{(n/2+\tilde{q}+1)}B_{(n/2+\tilde{q}+1)}} + \sum_{\tilde{Q}=1}^{n/2+\tilde{q}} (\eta_{\tilde{Q}})_\Lambda (\varepsilon_{\tilde{Q}})^{B_{(n/2+\tilde{q}+1)}A_{(n/2+\tilde{q}+1)}} = \\
= {\eta_\Lambda}^{B_{(n/2+\tilde{q}+1)}A_{(n/2+\tilde{q}+1)}}.
\end{gathered} \tag{16.4}$$

16.2 Proof of the connecting operator relations

We need to prove the identity (8.27)

$$\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}\eta^\Theta{}_{CB} = \frac{N}{4}(g_{\Lambda\Psi}g^{\Omega\Theta} + \delta_\Lambda{}^\Theta\delta_\Psi{}^\Omega - \delta_\Psi{}^\Theta\delta_\Lambda{}^\Omega). \tag{16.5}$$

1. First step.

$$\boxed{\boxed{\begin{array}{rcl} \eta_\Lambda{}^{AB}\eta_\Psi{}^{CD}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC} &=& \frac{N}{4}((\eta_Q)_\Psi(\eta_P)_\Lambda + (\eta_Q)_\Lambda(\eta_P)_\Psi), \quad P \neq Q. \\ \eta_{\Lambda AD}(\varepsilon_P)^{AB}(\varepsilon_Q)_{BC}\eta_\Psi{}^{CD} &=& \frac{N}{4}((\eta_Q)_\Psi(\eta_P)_\Lambda - (\eta_Q)_\Lambda(\eta_P)_\Psi), \quad P \neq Q. \end{array}}} \tag{16.6}$$

Proof.

$$\begin{aligned} \eta_\Lambda{}^{AB}\eta_\Psi{}^{CD}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC} &= (\eta_\Lambda{}^{BA} + \sum_{\tilde{P}}(\varepsilon_{\tilde{P}})^{AB}(\eta_{\tilde{P}})_\Lambda)\eta_\Psi{}^{CD}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC} = \\ &= \eta_\Lambda{}^{BA}(\eta_\Psi{}^{DC} + \sum_{\tilde{P}}(\varepsilon_{\tilde{P}})^{CD}(\eta_{\tilde{P}})_\Psi)(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC} + \\ &+ \sum_{\tilde{P}}(\varepsilon_{\tilde{P}})^{AB}(\eta_{\tilde{P}})_\Lambda\eta_\Psi{}^{CD}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC} = \eta_\Lambda{}^{BA}\eta_\Psi{}^{DC}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC} + \\ &+ \sum_{\tilde{P}}((\varepsilon_{\tilde{P}})^{AB}(\eta_{\tilde{P}})_\Lambda\eta_\Psi{}^{CD}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC} + (\varepsilon_{\tilde{P}})^{CD}(\eta_{\tilde{P}})_\Psi\eta_\Lambda{}^{BA}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC}). \end{aligned} \tag{16.7}$$

   (a) First step.

$$\begin{aligned} \eta_\Lambda{}^{BA}\eta_\Psi{}^{DC}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC} &= \tfrac{N}{2}(\eta_P)_\Lambda(\eta_Q)_\Psi - (\varepsilon_P)^{BA}\eta_\Psi{}^{DC}\eta_{\Lambda AD}(\varepsilon_Q)_{BC} = \\ &= \tfrac{N}{2}(\eta_P)_\Lambda(\eta_Q)_\Psi + (\varepsilon_Q)^{BA}\eta_\Psi{}^{DC}\eta_{\Lambda AD}(\varepsilon_P)_{BC} = \\ &= \tfrac{N}{2}((\eta_P)_\Lambda(\eta_Q)_\Psi + (\eta_Q)_\Lambda(\eta_P)_\Psi) - \eta_\Lambda{}^{BA}\eta_\Psi{}^{DC}(\varepsilon_Q)_{AD}(\varepsilon_P)_{BC}. \end{aligned} \tag{16.8}$$

$$\boxed{\begin{array}{c} \eta_\Lambda{}^{BA}\eta_\Psi{}^{DC}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC} = \\ \frac{N}{2}((\eta_P)_\Lambda(\eta_Q)_\Psi + (\eta_Q)_\Lambda(\eta_P)_\Psi) - \eta_\Lambda{}^{BA}\eta_\Psi{}^{DC}(\varepsilon_Q)_{AD}(\varepsilon_P)_{BC}. \end{array}} \tag{16.9}$$

   (b) Second step.

$$\begin{gathered} (\varepsilon_{\tilde{P}})^{AD}(\varepsilon_P)_{AB}(\varepsilon_{\tilde{Q}})^{CB}(\varepsilon_Q)_{CD} \overset{\tilde{P}\neq\tilde{Q}}{=} -(\varepsilon_{\tilde{Q}})^{AD}(\varepsilon_P)_{AB}(\varepsilon_{\tilde{P}})^{CB}(\varepsilon_Q)_{CD} = \\ -(\varepsilon_{\tilde{Q}})^{BC}(\varepsilon_P)_{BA}(\varepsilon_{\tilde{P}})^{DA}(\varepsilon_Q)_{DC}. \end{gathered} \tag{16.10}$$

$$\boxed{(\varepsilon_{\tilde{P}})^{AD}(\varepsilon_P)_{AB}(\varepsilon_{\tilde{Q}})^{CB}(\varepsilon_Q)_{CD} \overset{\tilde{P}\neq\tilde{Q}}{=} 0.} \tag{16.11}$$

   (c) Third step.

$$\begin{aligned} \eta_\Lambda{}^{AD}(\varepsilon_P)_{AB}(\varepsilon_{\tilde{P}})^{CB}(\varepsilon_Q)_{CD} &= -\eta_{\Lambda KL}(\varepsilon_{\tilde{Q}})^{KD}(\varepsilon_{\tilde{Q}})^{AL}(\varepsilon_P)_{AB}(\varepsilon_{\tilde{P}})^{CB}(\varepsilon_Q)_{CD} = \\ &= \eta_{\Lambda KL}(\varepsilon_{\tilde{Q}})^{KD}(\varepsilon_P)^{AL}(\varepsilon_{\tilde{P}})_{AB}(\varepsilon_Q)^{CB}(\varepsilon_{\tilde{Q}})_{CD} = \eta_{\Lambda KL}(\varepsilon_P)^{AL}(\varepsilon_{\tilde{P}})_{AB}(\varepsilon_Q)^{KB} = \\ &= -\eta_\Lambda{}^{AD}(\varepsilon_P)_{KD}(\varepsilon_{\tilde{P}})_{AB}(\varepsilon_Q)^{KB} = -\eta_\Lambda{}^{AD}(\varepsilon_Q)_{CD}(\varepsilon_P)_{AB}(\varepsilon_{\tilde{P}})^{CB}. \end{aligned} \tag{16.12}$$

$$\boxed{\eta_\Lambda{}^{AD}(\varepsilon_P)_{AB}(\varepsilon_{\tilde{P}})^{CB}(\varepsilon_Q)_{CD} = 0.} \tag{16.13}$$

$\square$

2. Second step.

$$\boxed{\boxed{\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}(\varepsilon_Q)_{BC} = \tfrac{N}{4}(g_{\Lambda\Psi}(\eta_Q)^\Omega + (\eta_Q)^\Lambda\delta_\Psi{}^\Omega - (\eta_Q)_\Psi\delta_\Lambda{}^\Omega).}} \tag{16.14}$$

Proof.

$$\begin{gathered}
\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}(\varepsilon_Q)_{BC} = \\
= \eta_\Lambda{}^{AB}\eta_{\Psi AD}(-\eta^\Omega{}_{KL}(\varepsilon_P)^{KD}(\varepsilon_P)^{CL} + (\eta_P)^\Omega(\varepsilon_P)^{CD})(\varepsilon_Q)_{BC} = \\
= -\eta_\Lambda{}^{AB}(-\eta_\Psi{}^{KM}(\varepsilon_P)_{MA} + (\eta_P)_\Psi\delta_A{}^K)(\varepsilon_Q)_{BC}(\varepsilon_P)^{CL}\eta^\Omega{}_{KL} + \\
+(\eta_P)^\Omega\tfrac{N}{4}((\eta_P)_\Psi(\eta_Q)_\Lambda - (\eta_P)_\Lambda(\eta_Q)_\Psi) = \\
= (-\eta_{\Lambda NM}(\varepsilon_P)^{NB} + (\eta_P)_\Lambda\delta_M{}^B)\eta_\Psi{}^{KM}(\varepsilon_Q)_{BC}(\varepsilon_P)^{CL}\eta^\Omega{}_{KL} - \\
-\tfrac{N}{4}(\eta_P)_\Psi((\eta_Q)_\Lambda(\eta_P)^\Omega - (\eta_Q)^\Omega(\eta_P)_\Lambda) + \tfrac{N}{4}(\eta_P)^\Omega((\eta_P)_\Psi(\eta_Q)_\Lambda + (\eta_P)_\Lambda(\eta_Q)_\Psi) = \\
= \eta_{\Lambda NM}\eta_\Psi{}^{KM}\eta^\Omega{}_{KL}(\varepsilon_Q)^{LN} + \tfrac{N}{4}(\eta_P)_\Lambda(\underbrace{(\eta_Q)_\Psi(\eta_P)^\Omega}_{1} - \underbrace{(\eta_P)_\Psi(\eta_Q)^\Omega}_{2}) - \\
-\tfrac{N}{4}(\eta_P)_\Psi(\underbrace{(\eta_Q)_\Lambda(\eta_P)^\Omega}_{3} - \underbrace{(\eta_Q)^\Omega(\eta_P)_\Lambda}_{2}) + \tfrac{N}{4}(\eta_P)^\Omega(\underbrace{(\eta_P)_\Psi(\eta_Q)_\Lambda}_{3} - \underbrace{(\eta_P)_\Lambda(\eta_Q)_\Psi}_{1}) = \\
= (-\eta_\Lambda{}^{AL}(\varepsilon_Q)_{AM} + (\eta_Q)_\Lambda\delta_M{}^L)\eta_\Psi{}^{KM}\eta^\Omega{}_{KL} = \\
= -\eta_\Lambda{}^{AL}(-\eta_{\Psi AD}(\varepsilon_Q)^{DK} + (\eta_Q)_\Psi\delta_A{}^K)\eta^\Omega{}_{KL} + \tfrac{N}{2}(\eta_Q)_\Lambda\delta_\Psi{}^\Omega = \\
= \eta_\Lambda{}^{AL}\eta_{\Psi AD}(-\eta^{\Omega CD}(\varepsilon_Q)_{CL} + (\eta_Q)^\Omega\delta_L{}^D) + \tfrac{N}{2}((\eta_Q)_\Lambda\delta_\Psi{}^\Omega - (\eta_Q)_\Psi\delta_\Lambda{}^\Omega) = \\
= -\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}(\varepsilon_Q)_{CB} + \tfrac{N}{2}(g_{\Lambda\Psi}(\eta_Q)^\Omega + (\eta_Q)_\Lambda\delta_\Psi{}^\Omega - (\eta_Q)_\Psi\delta_\Lambda{}^\Omega).
\end{gathered} \tag{16.15}$$

$$\boxed{\eta_\Lambda{}^{AB}\eta^\Omega{}_{CB}\eta_\Psi{}^{CD}(\varepsilon_Q)_{AD} = \tfrac{N}{4}(-g_{\Lambda\Psi}(\eta_Q)^\Omega + (\eta_Q)_\Lambda\delta_\Psi{}^\Omega + (\eta_Q)_\Psi\delta_\Lambda{}^\Omega).} \tag{16.16}$$

□

3. Third step.

$$\boxed{\boxed{\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}\eta^\Theta{}_{CB} = \tfrac{N}{4}(g_{\Lambda\Psi}g^{\Omega\Theta} + \delta_\Lambda{}^\Theta\delta_\Psi{}^\Omega - \delta_\Psi{}^\Theta\delta_\Lambda{}^\Omega).}} \tag{16.17}$$

Proof.

$$\begin{gathered}
\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}\eta^\Theta{}_{CB} = \\
= \eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}(-\eta^{\Theta KL}(\varepsilon_Q)_{KB}(\varepsilon_Q)_{LC} + (\eta_Q)^\Theta(\varepsilon_Q)_{BC}) = \\
= -\eta_\Lambda{}^{AB}\eta_{\Psi AD}(-\eta^\Omega{}_{ML}(\varepsilon_Q)^{MD} + (\eta_Q)^\Omega\delta_L{}^D)(\varepsilon_Q)_{KB}\eta^{\Theta KL} + \\
+(\eta_Q)^\Theta\tfrac{N}{4}(g_{\Lambda\Psi}(\eta_Q)^\Omega + (\eta_Q)_\Lambda\delta_\Psi{}^\Omega - (\eta_Q)_\Psi\delta_\Lambda{}^\Omega) = \\
= \eta_\Lambda{}^{AB}(-\eta_\Psi{}^{MN}(\varepsilon_Q)_{NA} + (\eta_Q)_\Psi\delta_A{}^M)\eta^\Omega{}_{ML}\eta^{\Theta KL}(\varepsilon_Q)_{KB} + \\
+\tfrac{N}{4}(\eta_Q)^\Theta(g_{\Lambda\Psi}(\eta_Q)^\Omega + (\eta_Q)_\Lambda\delta_\Psi{}^\Omega - (\eta_Q)_\Psi\delta_\Lambda{}^\Omega) - \\
-\tfrac{N}{4}(\eta_Q)^\Omega(g_{\Lambda\Psi}(\eta_Q)^\Theta + (\eta_Q)_\Lambda\delta_\Psi{}^\Theta - (\eta_Q)_\Psi\delta_\Lambda{}^\Theta) = \\
= -(-\eta_{\Lambda SN}(\varepsilon_Q)^{SB} + (\eta_Q)^\Lambda\delta_N{}^B)\eta_\Psi{}^{MN}\eta^\Omega{}_{ML}\eta^{\Theta KL}(\varepsilon_Q)_{KB} + \tfrac{N}{4}(\eta_Q)^\Theta(g_{\Lambda\Psi}(\eta_Q)^\Omega + \\
+(\eta_Q)_\Lambda\delta_\Psi{}^\Omega - (\eta_Q)_\Psi\delta_\Lambda{}^\Omega) - \tfrac{N}{4}(\eta_Q)^\Omega(g_{\Lambda\Psi}(\eta_Q)^\Theta + (\eta_Q)_\Lambda\delta_\Psi{}^\Theta - (\eta_Q)_\Psi\delta_\Lambda{}^\Theta) + \\
+\tfrac{N}{4}(\eta_Q)_\Psi(\delta_\Lambda{}^\Omega(\eta_Q)^\Theta + (\eta_Q)_\Lambda g^{\Omega\Theta} - (\eta_Q)^\Omega\delta_\Lambda{}^\Theta) = \eta_{\Lambda SN}\eta_\Psi{}^{MN}\eta^\Omega{}_{ML}\eta^{\Theta SL} + \\
+\tfrac{N}{4}(\eta_Q)^\Theta(\underbrace{g_{\Lambda\Psi}(\eta_Q)^\Omega}_{1} + \underbrace{(\eta_Q)_\Lambda\delta_\Psi{}^\Omega}_{2} - \underbrace{(\eta_Q)_\Psi\delta_\Lambda{}^\Omega}_{3}) - \tfrac{N}{4}(\eta_Q)^\Omega(\underbrace{g_{\Lambda\Psi}(\eta_Q)^\Theta}_{1} + \underbrace{(\eta_Q)_\Lambda\delta_\Psi{}^\Theta}_{4} - \\
-\underbrace{(\eta_Q)_\Psi\delta_\Lambda{}^\Theta}_{5}) + \tfrac{N}{4}(\eta_Q)_\Psi(\underbrace{\delta_\Lambda{}^\Omega(\eta_Q)^\Theta}_{3} + \underbrace{(\eta_Q)_\Lambda g^{\Omega\Theta}}_{6} - \underbrace{(\eta_Q)^\Omega\delta_\Lambda{}^\Theta}_{5}) - \tfrac{N}{4}(\eta_Q)_\Lambda(\underbrace{\delta_\Psi{}^\Omega(\eta_Q)^\Theta}_{2} + \\
+\underbrace{(\eta_Q)_\Psi g^{\Omega\Theta}}_{6} - \underbrace{(\eta_Q)^\Omega\delta_\Psi{}^\Theta}_{4}) = \\
= -\eta_{\Psi AD}\eta^{\Omega CD}\eta^\Theta{}_{CB}\eta^{\Lambda AB} + \tfrac{N}{2}(g_{\Lambda\Psi}g^{\Omega\Theta} + \delta_\Lambda{}^\Theta\delta_\Psi{}^\Omega - \delta_\Psi{}^\Theta\delta_\Lambda{}^\Omega).
\end{gathered} \tag{16.18}$$

□

16.3 Proof of the infinitesimal transformation relations

We need to prove the identity (10.5).

1. Case 1).

$$\begin{gathered}
S_\Lambda{}^\Psi \eta_\Psi{}^{AB} = -\eta_\Lambda{}^{CD} S_C{}^A E_D{}^K S_K{}^M E_M{}^B, \\
T_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \eta_\Lambda{}^{CB} T_C{}^A - \eta_\Lambda{}^{AD} E_D{}^K T_K{}^M E_M{}^B, \\
(\eta_I)^\Lambda T_\Lambda{}^\Psi \eta_\Psi{}^{AB} = (\varepsilon_I)^{CB} T_C{}^A - (\varepsilon_I)^{AD} E_D{}^K T_K{}^M E_M{}^B, \\
E_D{}^K T_K{}^M E_M{}^B = (\varepsilon_I)^{CB} T_C{}^A (\varepsilon_I)_{AD} - (\eta_I)^\Lambda T_\Lambda{}^\Psi \eta_\Psi{}^{AB} (\varepsilon_I)_{AD}, \\
T_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \eta_\Lambda{}^{CB} T_C{}^A - \eta_\Lambda{}^{AD} (\varepsilon_I)^{KB} T_K{}^M (\varepsilon_I)_{MD} + \eta_\Lambda{}^{AD} (\varepsilon_I)_{MD} (\eta_I)^\Omega T_\Omega{}^\Phi \eta_\Phi{}^{MB}, \\
T_C{}^A = -E_C{}^B T_B{}^D E_D{}^A.
\end{gathered} \tag{16.19}$$

2. Case 2).

$$\begin{gathered}
S_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \eta_\Lambda{}^{CD} S_C{}^A (\eta_I)^\Omega S_\Omega{}^\Phi \eta_\Phi{}^{KB} \varepsilon_{KL} S_X{}^L \varepsilon^{MX} (\varepsilon_I)_{MD}, \\
T_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \\
= \eta_\Lambda{}^{CB} T_C{}^A + \eta_\Lambda{}^{AD} (\eta_I)^\Omega T_\Omega{}^\Phi \eta_\Psi{}^{KB} (\varepsilon_I)_{KD} + \eta_\Lambda{}^{AD} (\varepsilon_I)^{KB} \varepsilon_{KL} T_X{}^L \varepsilon^{MX} (\varepsilon_I)_{MD}, \\
T_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \eta_\Lambda{}^{CB} T_C{}^A - \eta_\Lambda{}^{AD} (\varepsilon_I)^{KB} T_K{}^M (\varepsilon_I)_{MD} + \eta_\Lambda{}^{AD} (\varepsilon_I)_{MD} (\eta_I)^\Omega T_\Omega{}^\Phi \eta_\Phi{}^{MB}, \\
T_A{}^K = -T_C{}^D \varepsilon_{AD} \varepsilon^{KC}.
\end{gathered} \tag{16.20}$$

Substitute in this equation

$$T_C{}^A = \frac{1}{2} T^{\Theta\Phi} \eta_\Phi{}^{AB} \eta_{\Theta CB} \tag{16.21}$$

and obtain

$$\begin{gathered}
T_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \tfrac{1}{2}(\eta_\Lambda{}^{CB} T^{\Theta\Phi} \eta_\Phi{}^{AD} \eta_{\Theta CD} - \eta_\Lambda{}^{AD} (\varepsilon_I)^{KB} T^{\Theta\Phi} \eta_\Phi{}^{MN} \eta_{\Theta KN} (\varepsilon_I)_{MD}) + \\
+ \eta_\Lambda{}^{AD} (\varepsilon_I)_{MD} (\eta_I)^\Omega T_\Omega{}^\Phi \eta_\Phi{}^{MB} = \tfrac{1}{2}(T_\Lambda{}^\Phi \eta_\Phi{}^{AB} - \eta_{\Lambda CD} T^{\Theta\Phi} \eta_\Phi{}^{AD} \eta_\Theta{}^{CB} - \\
- \eta_\Lambda{}^{AD} (\eta_I)_\Theta T^{\Theta\Phi} \eta_\Phi{}^{MB} (\varepsilon_I)_{MD} + \eta_\Lambda{}^{AD} T^{\Theta\Phi} \eta_\Theta{}^{LB} (\varepsilon_I)_{LN} \eta_\Phi{}^{MN} (\varepsilon_I)_{MD}) + \\
+ \eta_\Lambda{}^{AD} (\varepsilon_I)_{MD} (\eta_I)^\Omega T_\Omega{}^\Phi \eta_\Phi{}^{MB} = \tfrac{1}{2}(T_\Lambda{}^\Phi \eta_\Phi{}^{AB} - T^\Theta{}_\Lambda \eta_\Theta{}^{AB} + T^{\Theta\Phi} \eta_{\Phi CD} \eta_\Lambda{}^{AD} \eta_\Theta{}^{CB} - \\
- \eta_\Lambda{}^{AD} (\eta_I)_\Theta T^{\Theta\Phi} \eta_\Phi{}^{MB} (\varepsilon_I)_{MD} - \eta_\Lambda{}^{AD} T^{\Theta\Phi} \eta_{\Phi LD} \eta_\Theta{}^{LB} + \\
+ \eta_\Lambda{}^{AD} T^{\Theta\Phi} \eta_\Theta{}^{LB} (\eta_I)_\Phi (\varepsilon_I)_{LD}) + \eta_\Lambda{}^{AD} (\varepsilon_I)_{MD} (\eta_I)^\Omega T_\Omega{}^\Phi \eta_\Phi{}^{MB} = T_\Lambda{}^\Psi \eta_\Psi{}^{AB}.
\end{gathered} \tag{16.22}$$

16.4 Proof of the Lie bracket identity

We need to prove the identity (11.70) (see (11.72)).

$$
\begin{gathered}
(L_\pm)_{[x,y]}(Y_K)^A = \\
= (x^\Theta\partial_\Theta y^\Omega - y^\Theta\partial_\Theta x^\Omega)\partial_\Omega(Y_K)^A - \sum_J^{2N}(Y_J)^B(P_J)^\Lambda{}_B(P^*_K)_\Theta{}^A\partial_\Lambda(x^\Xi\partial_\Xi y^\Theta - y^\Xi\partial_\Xi x^\Theta) = \\
= \underbrace{(x^\Theta\partial_\Theta y^\Omega - y^\Theta\partial_\Theta x^\Omega)\partial_\Omega(Y_K)^A}_{1} - \underbrace{\sum_J^{2N}(Y_J)^B(P_J)^\Lambda{}_B(P^*_K)_\Theta{}^A(\partial_\Lambda x^\Xi\partial_\Xi y^\Theta - \partial_\Lambda y^\Xi\partial_\Xi x^\Theta)}_{6} - \\
- \underbrace{\sum_J^{2N}(Y_J)^B(P_J)^\Lambda{}_B(P^*_K)_\Theta{}^A(x^\Xi\partial_\Lambda\partial_\Xi y^\Theta - y^\Xi\partial_\Lambda\partial_\Xi x^\Theta)}_{4}, \\
((L_\pm)_x(L_\pm)_y - (L_\pm)_y(L_\pm)_x)(Y)^A = x^\Phi\partial_\Phi(y^\Omega\partial_\Omega(Y_K)^A - \sum_J^{2N}(Y_J)^B(P_J)^\Lambda{}_B(P^*_K)_\Theta{}^A\partial_\Lambda y^\Theta) - \\
-(y^\Omega\partial_\Omega(Y_K)^C - \sum_{I,J}^{2N}(Y_J)^B(P_J)^\Lambda{}_B(P^*_I)_\Theta{}^C\partial_\Lambda y^\Theta)(P_I)^\Phi{}_C(P^*_K)_\Xi{}^A\partial_\Phi x^\Xi - \\
-y^\Phi\partial_\Phi(x^\Omega\partial_\Omega(Y_K)^A - \sum_J^{2N}(Y_J)^B(P_J)^\Lambda{}_B(P^*_K)_\Theta{}^A\partial_\Lambda x^\Theta) + \\
+(x^\Omega\partial_\Omega(Y_K)^C - \sum_{I,J}^{2N}(Y_J)^B(P_J)^\Lambda{}_B(P^*_I)_\Theta{}^C\partial_\Lambda x^\Theta)(P_I)^\Phi{}_C(P^*_K)_\Xi{}^A\partial_\Phi y^\Xi = \\
= \underbrace{x^\Phi\partial_\Phi y^\Omega\partial_\Omega(Y_K)^A}_{1} + \underbrace{x^\Phi y^\Omega\partial_\Phi\partial_\Omega(Y_K)^A}_{2} - \underbrace{x^\Phi\sum_J^{2N}\partial_\Phi(Y_J)^B(P_J)^\Lambda{}_B(P^*_K)_\Theta{}^A\partial_\Lambda y^\Theta}_{3} - \\
- \underbrace{x^\Theta\sum_J^{2N}(Y_J)^B(P_J)^\Lambda{}_B(P^*_K)_\Theta{}^A\partial_\Phi\partial_\Lambda y^\Theta}_{4} - \underbrace{y^\Omega\sum_J^{2N}\partial_\Omega(Y_J)^C(P_J)^\Phi{}_C(P^*_K)_\Xi{}^A\partial_\Phi x^\Xi}_{5} + \\
+ \underbrace{\sum_{I,J}^{2N}(Y_J)^B(P_J)^\Lambda{}_B(P^*_I)_\Theta{}^C\partial_\Lambda y^\Theta(P_I)^\Phi{}_C(P^*_K)_\Xi{}^A\partial_\Phi x^\Xi}_{6} - \\
- \underbrace{y^\Phi\partial_\Phi x^\Omega\partial_\Omega(Y_K)^A}_{1} - \underbrace{y^\Phi x^\Omega\partial_\Phi\partial_\Omega(Y_K)^A}_{2} + \underbrace{y^\Phi\sum_J^{2N}\partial_\Phi(Y_J)^B(P_J)^\Lambda{}_B(P^*_K)_\Theta{}^A\partial_\Lambda x^\Theta}_{5} + \\
+ \underbrace{y^\Phi\sum_J^{2N}(Y_J)^B(P_J)^\Lambda{}_B(P^*_K)_\Theta{}^A\partial_\Phi\partial_\Lambda x^\Theta}_{4} + \underbrace{x^\Omega\sum_J^{2N}\partial_\Omega(Y_J)^C(P_J)^\Phi{}_C(P^*_K)_\Xi{}^A\partial_\Phi y^\Xi}_{3} - \\
- \underbrace{\sum_{I,J}^{2N}(Y_J)^B(P_J)^\Lambda{}_B(P^*_I)_\Theta{}^C\partial_\Lambda x^\Theta(P_I)^\Phi{}_C(P^*_K)_\Xi{}^A\partial_\Phi y^\Xi}_{6}.
\end{gathered}
\tag{16.23}
$$

16.5 Proof of the curvature tensor relations

We need to prove the identity (12.7).
The curvature tensor in the spinor bundle is given by

$$(\mathcal{R}_K)_{\Lambda\Psi C}{}^{A} = 2(\partial_{[\Lambda}(\Gamma_K)_{\Psi]C}{}^{A} + (\Gamma_K)_{[\Lambda|L|}{}^{A}(\Gamma_K)_{\Psi]C}{}^{L}). \tag{16.24}$$

Then $\partial_\Lambda(\varepsilon_K)_{AB}{}^{CD} = 0$ under the condition (11.33) means

$$\partial_{[\Lambda}(\Gamma_K)_{\Psi]A}{}^{C} = \underbrace{-\frac{2}{n-4}((\varepsilon_K)_{AB}{}^{CD} - \frac{n-2}{N}\delta_A{}^{C}\delta_B{}^{D})}_{:=(\tilde{\varepsilon}_K)_{AB}{}^{CD}}\,\partial_{[\Lambda}(\Gamma_K)_{\Psi]D}{}^{B}. \tag{16.25}$$

In turn, $\nabla_\Lambda(\varepsilon_K)_{AB}{}^{CD} = \partial_\Lambda(\varepsilon_K)_{AB}{}^{CD} = 0$ under the condition (11.33) means

$$\begin{gathered}
(\Gamma_K)_{\Lambda A}{}^{L}(\tilde{\varepsilon}_K)_{LB}{}^{CD} + (\Gamma_K)_{\Lambda B}{}^{L}(\tilde{\varepsilon}_K)_{AL}{}^{CD} = (\Gamma_K)_{\Lambda L}{}^{C}(\tilde{\varepsilon}_K)_{AB}{}^{LD} + (\Gamma_K)_{\Lambda L}{}^{D}(\tilde{\varepsilon}_K)_{AB}{}^{CL},\\
(\Gamma_K)_{\Lambda A}{}^{L}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{LB}{}^{CD} + (\Gamma_K)_{\Lambda B}{}^{L}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{AL}{}^{CD} =\\
= (\Gamma_K)_{\Lambda L}{}^{C}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{AB}{}^{LD} + (\Gamma_K)_{\Lambda L}{}^{D}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{AB}{}^{CL},\\
(\Gamma_K)_{\Lambda A}{}^{L}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{LB}{}^{CD} + (\Gamma_K)_{\Lambda B}{}^{L}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{AL}{}^{CD} =\\
= (\Gamma_K)_{\Lambda C}{}^{A}(\Gamma_K)_{\Psi A}{}^{L}(\tilde{\varepsilon}_K)_{LB}{}^{CD} + (\Gamma_K)_{\Lambda L}{}^{D}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{AB}{}^{CL},\\
(\Gamma_K)_{[\Lambda|A|}{}^{L}(\Gamma_K)_{\Psi]C}{}^{A}(\tilde{\varepsilon}_K)_{LB}{}^{CD} = (\Gamma_K)_{[\Lambda|L|}{}^{D}(\Gamma_K)_{\Psi]B}{}^{L}.
\end{gathered} \tag{16.26}$$

Whence,

$$(\mathcal{R}_K)_{\Lambda\Psi C}{}^{M} = (\mathcal{R}_K)_{\Lambda\Psi L}{}^{D}\tilde{\varepsilon}_{CD}{}^{ML}, \tag{16.27}$$

16.6 Proof of the auxiliary identities for the theorem on the spin analogs of the curvature tensor

We need to prove the identity (12.9).

$$\boxed{\boxed{\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_2} = \tfrac{N}{2}g_{\Lambda_1[\Lambda_4}g_{\Lambda_3]\Lambda_2}.}} \tag{16.28}$$

Proof.

$$\begin{gathered}
\eta_{\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2A_1A_3} = (-\eta_{\Lambda_1B_1B_2}(\varepsilon_I)^{A_1B_2}(\varepsilon_I)^{B_1A_2} + (\eta_I)_{\Lambda_1}(\varepsilon_I)^{A_1A_2})\eta_{\Lambda_2A_1A_3} =\\
= -\eta_{\Lambda_1B_1B_2}(\varepsilon_I)^{B_1A_2}(-\eta_{\Lambda_2}{}^{A_1B_2}(\varepsilon_I)_{A_1A_3} + (\eta_I)_{\Lambda_2}\delta_{A_3}{}^{B_2}) + (\eta_I)_{\Lambda_1}(\varepsilon_I)^{A_1A_2}\eta_{\Lambda_2A_1A_3} =\\
= \eta_{\Lambda_1B_1B_2}\eta_{\Lambda_2}{}^{A_1B_2}(\varepsilon_I)^{B_1A_2}(\varepsilon_I)_{A_1A_3} - (\eta_I)_{\Lambda_2}(\varepsilon_I)^{B_1A_2}\eta_{\Lambda_1B_1A_3} + (\eta_I)_{\Lambda_1}(\varepsilon_I)^{A_1A_2}\eta_{\Lambda_2A_1A_3}.
\end{gathered} \tag{16.29}$$

Sum over I alternating on $\Lambda_1,\ \Lambda_2$

$$\begin{gathered}
\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3} = \tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2}{}_{A_1A_3},\\
\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_2} =\\
= \eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}(\tfrac{2}{n-4}\varepsilon^{B_1A_2}{}_{A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_2}) =\\
= \eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1B_3|}\eta_{\Lambda_4]}{}^{B_1B_3} =\\
= -\eta_{[\Lambda_1}{}^{A_1B_2}\eta_{\Lambda_2]A_1B_3}\eta_{[\Lambda_3}{}^{B_1B_3}\eta_{\Lambda_4]B_1B_2} + \tfrac{N}{2}(g_{\Lambda_1[\Lambda_4}g_{\Lambda_3]\Lambda_2} - g_{\Lambda_1[\Lambda_3}g_{\Lambda_4]\Lambda_2}),
\end{gathered} \tag{16.30}$$

whence the demanded follows. □

We need to prove the identity (12.10).

$$\boxed{\boxed{\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} = Ng_{[\Lambda_3|[\Lambda_2}g_{\Lambda_1]|[\Lambda_6}g_{\Lambda_5]|\Lambda_4]}.}} \tag{16.31}$$

Proof.

$$\begin{gathered}
\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} = \\
= \tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2}{}_{A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} = \\
= -\tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2A_4A_3}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} + \\
+\tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3}{}^{B_1A_2}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} = \\
= \tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2}{}_{A_4A_5}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} + \\
+\tfrac{4}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3}{}^{B_1A_2}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} = \\
= \eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4B_3|}\eta_{\Lambda_6]}{}^{B_1B_3} - \\
-\tfrac{4}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_5|}\eta_{\Lambda_4]}{}^{A_6A_5}\eta_{[\Lambda_5|A_6A_2|}\eta_{\Lambda_6]}{}^{B_1A_2} - \\
-\tfrac{8}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_5|}g_{\Lambda_4][\Lambda_6}\eta_{\Lambda_5]}{}^{B_1A_5} = \\
\left|\begin{gathered}
\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4B_3|}\eta_{\Lambda_6]}{}^{B_1B_3} = \\
= 2\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4B_3|}g_{\Lambda_6][\Lambda_1}\eta_{\Lambda_2]}{}^{A_1B_3} + \\
+\eta_{[\Lambda_1|A_4B_3|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5}{}^{B_1B_3}\eta_{\Lambda_6]B_1B_2} = \\
= 2\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4B_3|}g_{\Lambda_6][\Lambda_1}\eta_{\Lambda_2]}{}^{A_1B_3} + \\
+2\eta_{[\Lambda_2}{}^{A_1B_2}g_{\Lambda_1][\Lambda_4}\eta_{\Lambda_3]A_1A_3}\eta_{[\Lambda_5}{}^{B_1A_3}\eta_{\Lambda_6]B_1B_2} + \\
+\eta_{[\Lambda_1|A_1A_3|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4B_3}\eta_{[\Lambda_5}{}^{B_1B_3}\eta_{\Lambda_6]B_1B_2}
\end{gathered}\right| \\
= (-1 + \tfrac{4}{n-4})\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} + \\
+(\tfrac{8}{n-4} - 4)\tfrac{N}{2}g_{[\Lambda_3|[\Lambda_2}g_{\Lambda_1]|[\Lambda_5}g_{\Lambda_6]|\Lambda_4]},
\end{gathered}\tag{16.32}$$

whence the demanded follows. □

We need to prove the identity (12.11).

$$\boxed{\begin{gathered}
\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2} = \\
= \tfrac{N}{4}\big(g_{\Lambda_1[\Lambda_8}g_{\Lambda_7]\Lambda_2}g_{\Lambda_3[\Lambda_6}g_{\Lambda_5]\Lambda_4} - g_{\Lambda_1[\Lambda_6}g_{\Lambda_5]\Lambda_2}g_{\Lambda_4[\Lambda_8}g_{\Lambda_7]\Lambda_3} + \\
+g_{\Lambda_1[\Lambda_4}g_{\Lambda_3]\Lambda_2}g_{\Lambda_5[\Lambda_8}g_{\Lambda_7]\Lambda_6}\big) + N\big(g_{[\Lambda_3|[\Lambda_2}g_{\Lambda_1][\Lambda_8}g_{\Lambda_7][\Lambda_6}g_{\Lambda_5]|\Lambda_4]} + \\
+g_{[\Lambda_5|[\Lambda_4}g_{\Lambda_3][\Lambda_7}g_{\Lambda_8][\Lambda_2}g_{\Lambda_1]|\Lambda_6]} + g_{[\Lambda_3|[\Lambda_7}g_{\Lambda_8][\Lambda_5}g_{\Lambda_6][\Lambda_1}g_{\Lambda_2]|\Lambda_4]}\big).
\end{gathered}}\tag{16.33}$$

Proof.
1. First step.

$$\begin{gathered}
\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2} = \\
= \tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2}{}_{A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2} = \\
= -\tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2A_4A_3}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2} + \\
+\tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3}{}^{B_1A_2}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2} = \\
= \tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2}{}_{A_4A_5}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2} + \\
+\tfrac{4}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3}{}^{B_1A_2}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2} = \\
= \eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4A_5|}\eta_{\Lambda_6]}{}^{A_6A_5}\eta_{[\Lambda_7|A_6A_7|}\eta_{\Lambda_8]}{}^{B_1A_7} + \\
+\tfrac{4}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3}{}^{B_1A_2}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2} + \\
+\tfrac{4}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5}{}^{B_1A_2}\eta_{\Lambda_6]A_4A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2} = \\
= (1 - \tfrac{4}{n-4})\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4A_5|}\eta_{\Lambda_6]}{}^{A_6A_5}\eta_{[\Lambda_7|A_6A_7|}\eta_{\Lambda_8]}{}^{B_1A_7} + \\
+\tfrac{4}{n-4}\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2} + \\
+\tfrac{8}{n-4}\tfrac{N}{2}\big(\tfrac{1}{2}g_{\Lambda_1[\Lambda_3}g_{\Lambda_4]\Lambda_2}g_{\Lambda_5[\Lambda_8}g_{\Lambda_7]\Lambda_6} + g_{[\Lambda_3|[\Lambda_1}g_{\Lambda_2][\Lambda_8}g_{\Lambda_7][\Lambda_6}g_{\Lambda_5]|\Lambda_4]} - \\
-g_{[\Lambda_3|[\Lambda_1}g_{\Lambda_2][\Lambda_5}g_{\Lambda_6][\Lambda_7}g_{\Lambda_8]|\Lambda_4]} + \tfrac{1}{2}g_{\Lambda_5[\Lambda_8}g_{\Lambda_7]\Lambda_6}g_{\Lambda_1[\Lambda_4}g_{\Lambda_3]\Lambda_2} + \\
+g_{[\Lambda_5|[\Lambda_8}g_{\Lambda_7][\Lambda_1}g_{\Lambda_2][\Lambda_3}g_{\Lambda_4]|\Lambda_6]} - g_{[\Lambda_5|[\Lambda_8}g_{\Lambda_7][\Lambda_4}g_{\Lambda_3][\Lambda_2}g_{\Lambda_1]|\Lambda_6]}\big),
\end{gathered}\tag{16.34}$$

$$
\begin{gathered}
\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2}=\\
=\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4A_5|}\eta_{\Lambda_6]}{}^{A_6A_5}\eta_{[\Lambda_7|A_6A_7|}\eta_{\Lambda_8]}{}^{B_1A_7}
\end{gathered}
\tag{16.35}
$$

2. Second step.

$$
\begin{gathered}
\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2}=\\
=\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_1}{}^{A_8A_7}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_7|A_8A_2|}\eta_{\Lambda_8]}{}^{A_1A_2}+\\
+2\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_2|A_1A_3|}g_{\Lambda_1][\Lambda_8}\eta_{\Lambda_7]}{}^{A_1A_7}=\\
=\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_1}{}^{A_6A_5}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_5|A_6A_7|}\eta_{\Lambda_6]}{}^{A_8A_7}\eta_{[\Lambda_7|A_8A_2|}\eta_{\Lambda_8]}{}^{A_1A_2}+\\
+2(\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_2|A_1A_3|}g_{\Lambda_1][\Lambda_8}\eta_{\Lambda_7]}{}^{A_1A_7}+\\
+\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_2|A_1A_3|}g_{\Lambda_1][\Lambda_6}\eta_{\Lambda_5]}{}^{A_6A_5}\eta_{[\Lambda_7|A_6A_2|}\eta_{\Lambda_8]}{}^{A_1A_2})=\\
=-\eta_{[\Lambda_1|A_1A_3|}\eta_{\Lambda_2]}{}^{A_4A_3}\eta_{[\Lambda_3|A_4A_5|}\eta_{\Lambda_4]}{}^{A_6A_5}\eta_{[\Lambda_5|A_6A_7|}\eta_{\Lambda_6]}{}^{A_8A_7}\eta_{[\Lambda_7|A_8A_2|}\eta_{\Lambda_8]}{}^{A_1A_2}+\\
+2(\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_2|A_1A_3|}g_{\Lambda_1][\Lambda_8}\eta_{\Lambda_7]}{}^{A_1A_7}+\\
+\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_2|A_6A_2|}g_{\Lambda_1][\Lambda_6}\eta_{\Lambda_5]}{}^{A_6A_5}\eta_{[\Lambda_8|A_1A_3|}\eta_{\Lambda_7]}{}^{A_1A_2}+\\
+2\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_7|A_6A_3|}g_{\Lambda_8][\Lambda_2}g_{\Lambda_1][\Lambda_6}\eta_{\Lambda_5]}{}^{A_6A_5}+\\
+\eta_{[\Lambda_3}{}^{A_4A_3}g_{\Lambda_4][\Lambda_1}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_5|A_4A_7|}\eta_{\Lambda_6]}{}^{A_8A_7}\eta_{[\Lambda_7|A_8A_2|}\eta_{\Lambda_8]}{}^{A_1A_2})=\\
=2(\tfrac{N}{4}(g_{\Lambda_1[\Lambda_8}g_{\Lambda_7]\Lambda_2}g_{\Lambda_3[\Lambda_6}g_{\Lambda_5]\Lambda_4}-g_{\Lambda_1[\Lambda_6}g_{\Lambda_5]\Lambda_2}g_{\Lambda_4[\Lambda_8}g_{\Lambda_7]\Lambda_3}+\\
+g_{\Lambda_1[\Lambda_4}g_{\Lambda_3]\Lambda_2}g_{\Lambda_5[\Lambda_8}g_{\Lambda_7]\Lambda_6})+\tfrac{N}{2}(g_{[\Lambda_3|[\Lambda_2}g_{\Lambda_1][\Lambda_8}g_{\Lambda_7][\Lambda_6}g_{\Lambda_5]|\Lambda_4]}-\\
-g_{[\Lambda_3|[\Lambda_7}g_{\Lambda_8][\Lambda_1}g_{\Lambda_2][\Lambda_6}g_{\Lambda_5]|\Lambda_4]}+2g_{[\Lambda_5|[\Lambda_4}g_{\Lambda_3][\Lambda_7}g_{\Lambda_8][\Lambda_2}g_{\Lambda_1]|\Lambda_6]}+\\
+g_{[\Lambda_2|[\Lambda_8}g_{\Lambda_7][\Lambda_6}g_{\Lambda_5][\Lambda_3}g_{\Lambda_4]|\Lambda_1]}-g_{[\Lambda_3|[\Lambda_8}g_{\Lambda_7][\Lambda_6}g_{\Lambda_5][\Lambda_2}g_{\Lambda_1]|\Lambda_4]}-\\
-g_{[\Lambda_3|[\Lambda_7}g_{\Lambda_8][\Lambda_2}g_{\Lambda_1][\Lambda_6}g_{\Lambda_5]|\Lambda_4]}+g_{[\Lambda_3|[\Lambda_7}g_{\Lambda_8][\Lambda_5}g_{\Lambda_6][\Lambda_1}g_{\Lambda_2]|\Lambda_4]}))-\\
-\eta_{[\Lambda_1|A_1A_3|}\eta_{\Lambda_2]}{}^{A_4A_3}\eta_{[\Lambda_3|A_4A_5|}\eta_{\Lambda_4]}{}^{A_6A_5}\eta_{[\Lambda_5|A_6A_7|}\eta_{\Lambda_6]}{}^{A_8A_7}\eta_{[\Lambda_7|A_8A_2|}\eta_{\Lambda_8]}{}^{A_1A_2},
\end{gathered}
\tag{16.36}
$$

whence the demanded follows. □

Note 16.1. *It is obvious that these calculations are not applicable for* $n<8$*, since* $\frac{8}{n-4}-4$ *will not make the sense in this case.*

16.7 Proof of the twistor equation relations

We need to prove (13.4).

$$
\begin{gathered}
\nabla_\Psi X^C=\tfrac{2}{n}\eta_\Psi{}^{CD}\nabla_{AD}X^A,\\
\nabla_\Phi\nabla_\Psi X^C=\tfrac{2}{n}\eta_\Psi{}^{CD}\nabla_\Phi\nabla_{AD}X^A=\tfrac{2}{n}(\eta_\Psi{}^{CD}\nabla_{AD}\nabla_\Phi X^A-\eta_\Psi{}^{CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^AX^K)=\\
=(\tfrac{2}{n})^2\eta_\Psi{}^{CD}\nabla_{AD}\eta_\Phi{}^{AK}\nabla_{LK}X^L-\tfrac{2}{n}\eta_\Psi{}^{CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^AX^K=\\
=(\tfrac{2}{n})^2(-\eta_{\Psi AD}\nabla^{CD}\eta_\Phi{}^{AK}\nabla_{LK}X^L+\nabla_\Psi\eta_\Phi{}^{CK}\nabla_{LK}X^L)-\tfrac{2}{n}\eta_\Psi{}^{CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^AX^K=\\
=-(\tfrac{2}{n})^2\eta_{\Psi AD}\eta_\Phi{}^{AK}\nabla^{CD}\nabla_{LK}X^L+\tfrac{2}{n}\nabla_\Psi\nabla_\Phi X^C-\tfrac{2}{n}\eta_\Psi{}^{CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^AX^K.
\end{gathered}
\tag{16.37}
$$

$$
\begin{gathered}
(1-\tfrac{2}{n})\nabla_\Phi\nabla_\Psi X^C+\tfrac{2}{n}R_{\Phi\Psi K}{}^CX^K+\tfrac{2}{n}\eta_\Psi{}^{CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^AX^K=\\
=-(\tfrac{2}{n})^2\eta_{\Psi AD}\eta_\Phi{}^{AK}\nabla^{CD}\nabla_{LK}X^L,\\
(1-\tfrac{2}{n})\nabla_{(\Phi}\nabla_{\Psi)}X^C+\tfrac{2}{n}\eta_{(\Psi}{}^{CD}\eta^\Omega{}_{|AD}R_{\Omega|\Phi)K}{}^AX^K=\\
=-(\tfrac{2}{n})^2\tfrac{1}{2}g_{\Psi\Phi}\nabla^{CK}\nabla_{LK}X^L=-\tfrac{1}{n}g_{\Psi\Phi}\nabla^\Omega\nabla_\Omega X^C\\
(1-\tfrac{2}{n})\nabla^\Theta\nabla_\Theta X^C+\tfrac{2}{n}\eta^{\Phi CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^AX^K=-\nabla^\Theta\nabla_\Theta X^C,\\
\nabla^\Theta\nabla_\Theta X^C=-\tfrac{1}{n-1}\eta^{\Phi CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^AX^K.
\end{gathered}
\tag{16.38}
$$

$$\begin{gathered}
(1-\tfrac{2}{n})\nabla_{(\Phi}\nabla_{\Psi)}X^C+\tfrac{2}{n}\eta_{(\Psi}{}^{CD}\eta^{\Omega}{}_{|AD}R_{\Omega|\Phi)K}{}^{A}X^K=\tfrac{1}{n}g_{\Psi\Phi}\tfrac{1}{n-1}\eta^{\Theta CD}\eta^{\Omega}{}_{AD}R_{\Omega\Theta K}{}^{A}X^K\\
\nabla_{(\Phi}\nabla_{\Psi)}X^C=\tfrac{1}{(n-1)(n-2)}g_{\Psi\Phi}\eta^{\Theta CD}\eta^{\Omega}{}_{AD}R_{\Omega\Theta K}{}^{A}X^K-\tfrac{2}{n-2}\eta_{(\Psi}{}^{CD}\eta^{\Omega}{}_{|AD}R_{\Omega|\Phi)K}{}^{A}X^K,\\
2\nabla_{\Phi}\nabla_{\Psi}X^C=\tfrac{2}{n}\eta_{\Psi}{}^{CD}\eta^{\Gamma}{}_{AD}(R_{\Phi\Gamma K}{}^{A}X^K+\tfrac{2}{(n-1)(n-2)}g_{\Phi\Gamma}\eta^{\Theta AL}\eta^{\Omega}{}_{SL}R_{\Omega\Theta K}{}^{S}X^K-\\
-\tfrac{2}{n-2}\eta_{\Phi}{}^{AL}\eta^{\Omega}{}_{SL}R_{\Omega\Gamma K}{}^{S}X^K-\tfrac{2}{n-2}\eta_{\Gamma}{}^{AL}\eta^{\Omega}{}_{SL}R_{\Omega\Phi K}{}^{S}X^K)=\\
=\tfrac{2}{n}((-1-\tfrac{n}{n-2})\eta_{\Psi}{}^{CD}\eta^{\Omega}{}_{SD}R_{\Omega\Phi K}{}^{S}X^K+\tfrac{2}{(n-1)(n-2)}\eta_{\Psi}{}^{CD}\eta_{\Phi AD}\eta^{\Theta AL}\eta^{\Omega}{}_{SL}R_{\Omega\Theta K}{}^{S}X^K-\\
-\tfrac{2}{n-2}\eta_{\Psi}{}^{CD}\eta_{\Phi}{}^{AL}\eta^{\Omega}{}_{SL}\eta^{\Theta}{}_{AD}R_{\Omega\Theta K}{}^{S}X^K)=\\
=-\tfrac{4}{n-2}\eta_{\Psi}{}^{CD}\eta^{\Omega}{}_{SD}R_{\Omega\Phi K}{}^{S}X^K+\tfrac{4}{(n-1)(n-2)}\eta_{\Psi}{}^{CD}\eta_{\Phi AD}\eta^{\Theta AR}\eta^{\Omega}{}_{SR}R_{\Omega\Theta K}{}^{S}X^K.
\end{gathered}\tag{16.39}$$

Thus, the integrability conditions take the form

$$R_{\Phi\Psi K}{}^{C}X^K=-\tfrac{4}{n-2}\eta_{[\Psi}{}^{CD}\eta^{\Omega}{}_{|SD}R_{\Omega|\Phi]K}{}^{S}X^K+\tfrac{4}{(n-1)(n-2)}\eta_{[\Psi}{}^{CD}\eta_{\Phi]AD}\eta^{\Theta AL}\eta^{\Omega}{}_{SL}R_{\Omega\Theta K}{}^{S}X^K.\tag{16.40}$$

We assume that this condition is satisfied for any X^K. We use the Bianchi identity $2R^{\Delta[\Phi\Psi]\Lambda}=R^{\Psi\Phi\Delta\Lambda}$:

$$\begin{gathered}
(R_{\Phi\Psi K}{}^{C}=-\tfrac{4}{n-2}\eta_{[\Psi}{}^{CD}\eta^{\Omega}{}_{|SD}R_{\Omega|\Phi]K}{}^{S}+\tfrac{4}{(n-1)(n-2)}\eta_{[\Psi}{}^{CD}\eta_{\Phi]AD}\eta^{\Theta AL}\eta^{\Omega}{}_{SL}R_{\Omega\Theta K}{}^{S})\times\\
\times\tfrac{4}{N}\eta^{[\Lambda|KM|}\eta^{\Delta]}{}_{CM},
\end{gathered}$$

$$\begin{gathered}
-R^{\Phi\Psi\Lambda\Delta}=-\tfrac{4}{n-2}\tfrac{4}{N}\eta^{[\Psi|CD}\eta^{\Omega}{}_{SD}\eta^{[\Sigma|SL|}\eta^{\Theta]}{}_{KL}\eta^{[\Lambda|KM|}\eta^{\Delta]|}{}_{CM}\tfrac{1}{2}R_{\Omega}{}^{\Phi]}{}_{\Sigma\Theta}+\\
+\tfrac{4}{(n-1)(n-2)}\tfrac{4}{N}\eta^{[\Psi|CD|}\eta^{\Phi]}{}_{AD}\eta^{[\Theta|AL|}\eta^{\Omega]}{}_{SL}\eta^{[\Sigma|SN|}\eta^{\Xi]}{}_{KN}\eta^{[\Lambda|KM|}\eta^{\Delta]}{}_{CM}\tfrac{1}{2}R_{\Omega\Theta\Sigma\Xi}=\\
=-\tfrac{8}{n-2}(\tfrac{1}{2}g^{[\Psi|[\Delta}g^{\Lambda]\Theta}g^{\Sigma\Omega|}R_{\Omega}{}^{\Phi]}{}_{\Sigma\Theta}-\tfrac{1}{2}g^{\Omega[\Delta}g^{\Lambda]\Theta}g^{\Sigma[\Psi}R_{\Omega}{}^{\Phi]}{}_{\Sigma\Theta}+\tfrac{1}{4}g^{[\Psi|\Omega}g^{\Sigma[\Delta}g^{\Lambda]\Theta|}R_{\Omega}{}^{\Phi]}{}_{\Sigma\Theta})+\\
+\tfrac{8}{(n-1)(n-2)}(\tfrac{1}{4}g^{\Psi[\Delta}g^{\Lambda]\Phi}g^{\Theta\Xi}g^{\Sigma\Omega}R_{\Omega\Theta\Sigma\Xi}-\tfrac{1}{4}g^{\Psi\Xi}g^{\Sigma\Phi}g^{\Omega\Delta}g^{\Lambda\Theta}R_{\Omega\Theta\Sigma\Xi}+\tfrac{1}{4}g^{\Psi\Omega}g^{\Theta\Phi}g^{\Sigma\Delta}g^{\Lambda\Xi}R_{\Omega\Theta\Sigma\Xi}+\\
+g^{\Theta[\Phi}g^{\Psi][\Delta}g^{\Lambda]\Xi}g^{\Sigma\Omega}R_{\Omega\Theta\Sigma\Xi}+g^{\Theta[\Lambda}g^{\Delta][\Phi}g^{\Psi]\Xi}g^{\Sigma\Omega}R_{\Omega\Theta\Sigma\Xi}+g^{\Theta[\Lambda}g^{\Delta]\Sigma}g^{\Xi[\Psi}g^{\Phi]\Omega}R_{\Omega\Theta\Sigma\Xi})=\\
=-\tfrac{4}{n-2}g^{[\Delta|[\Psi}R^{\Phi]|\Lambda]}+\tfrac{4}{n-2}R^{[\Delta|[\Phi\Psi]|\Lambda]}-\tfrac{2}{n-2}R^{\Psi\Phi\Delta\Lambda}+\\
+\tfrac{2}{(n-1)(n-2)}Rg^{\Psi[\Delta}g^{\Lambda]\Phi}-\tfrac{2}{(n-1)(n-2)}R^{\Delta\Lambda\Phi\Psi}+\tfrac{2}{(n-1)(n-2)}R^{\Psi\Phi\Delta\Lambda}+\\
+\tfrac{8}{(n-1)(n-2)}R^{[\Phi|[\Lambda}g^{\Delta]|\Psi]}+\tfrac{8}{(n-1)(n-2)}g^{[\Phi|[\Delta}R^{\Lambda]|\Psi]}+\tfrac{8}{(n-1)(n-2)}R^{[\Phi|[\Lambda\Delta]|\Psi]}.
\end{gathered}\tag{16.41}$$

Finally,

$$R^{\Phi\Psi\Lambda\Delta}-\frac{4}{n-2}R^{[\Lambda\,|[\Phi}\,g^{\Psi]|\,\Delta]}+\frac{2}{(n-1)(n-2)}Rg^{[\Delta\,|[\Psi}\,g^{\Phi]|\,\Lambda]}=C^{\Phi\Psi\Lambda\Delta}=0.\tag{16.42}$$

16.8 Proof of the identities for n = 6 and n=8

16.8.1 Proof of the fundamental identities for n = 6

We need to prove (14.8).

$$A_{\alpha\beta d}{}^{c} = \eta_{[\alpha}{}^{ca}\eta_{\beta]da}, \tag{16.43}$$

$$\begin{gathered}
A_{\alpha\beta d}{}^{c}A_{\gamma}{}^{\beta}{}_{r}{}^{s} = \tfrac{1}{4}(\eta_\alpha{}^{ca}\eta_{\beta ad} - \eta_{\alpha ad}\eta_\beta{}^{ca})(\eta_\gamma{}^{sa_1}\eta^{\beta}{}_{a_1 r} - \eta_{\gamma a_1 r}\eta^{\beta s a_1}) = \\
= \tfrac{1}{4}(\eta_\alpha{}^{ca}\eta_\gamma{}^{sa_1}\varepsilon_{a_1 rad} - \eta_{\alpha ad}\eta_\gamma{}^{sa_1}\varepsilon_{a_1 r}{}^{ca} - \eta_\alpha{}^{ca}\eta_{\gamma a_1 r}\varepsilon_{ad}{}^{sa_1} + \eta_{\alpha ad}\eta_{\gamma a_1 r}\varepsilon^{casa_1}) = \\
= \tfrac{1}{4}(\tfrac{1}{2}\eta_\alpha{}^{ca}\eta_{\gamma kl}\varepsilon^{klsa_1}\varepsilon_{a_1 rad} - \eta_{\alpha ad}\eta_\gamma{}^{sa_1}(\delta_{a_1}{}^{c}\delta_r{}^{a} - \delta_{a_1}{}^{a}\delta_r{}^{c}) - \\
-\eta_\alpha{}^{ca}\eta_{\gamma a_1 r}(\delta_a{}^{s}\delta_d{}^{a_1} - \delta_a{}^{a_1}\delta_d{}^{s}) + \tfrac{1}{2}\eta_\alpha{}^{kl}\eta_{\gamma a_1 r}\varepsilon_{klad}\varepsilon^{casa_1}) = \\
= \tfrac{1}{4}(-3\eta_\alpha{}^{ca}\eta_{\gamma[ra}\delta_{d]}{}^{s} - \eta_{\alpha rd}\eta_\gamma{}^{sc} + \eta_{\alpha ad}\eta_\gamma{}^{sa}\delta_r{}^{c} - \\
-\eta_\alpha{}^{cs}\eta_{\gamma dr} + \eta_\alpha{}^{ca}\eta_{\gamma ar}\delta_d{}^{s} - 3\eta_\alpha{}^{[cs}\delta_d{}^{a_1]}\eta_{\gamma a_1 r} = \\
= \tfrac{1}{4}(-\eta_\alpha{}^{ck}\eta_{\gamma rk}\delta_d{}^{s} + \eta_\alpha{}^{cs}\eta_{\gamma rd} - \eta_\alpha{}^{ck}\eta_{\gamma kd}\delta_r{}^{s} - \eta_{\alpha rd}\eta_\gamma{}^{sc} + \eta_{\alpha ad}\eta_\gamma{}^{sa}\delta_r{}^{c} - \\
-\eta_\alpha{}^{cs}\eta_{\gamma dr} + \eta_\alpha{}^{ca}\eta_{\gamma ar}\delta_d{}^{s} - \eta_\alpha{}^{cs}\eta_{\gamma dr} + \eta_\alpha{}^{ck}\eta_{\gamma kr}\delta_d{}^{s} - \eta_\alpha{}^{sk}\eta_{\gamma kr}\delta_d{}^{c}) = \\
= \tfrac{1}{4}(3\eta_\alpha{}^{cs}\eta_{\gamma rd} + 3\eta_\alpha{}^{ck}\eta_{\gamma kr}\delta_d{}^{s} + \eta_\alpha{}^{ck}\eta_{\gamma dk}\delta_r{}^{s} + \eta_{\alpha dr}\eta_\gamma{}^{sc} + \eta_{\alpha kd}\eta_\gamma{}^{sk}\delta_r{}^{c} + \eta_\alpha{}^{sk}\eta_{\gamma rk}\delta_d{}^{c}) =
\end{gathered} \tag{16.44}$$

$$\left|\begin{gathered}
\eta_{\alpha dr}\eta_\gamma{}^{sc} + \eta_{\alpha kd}\eta_\gamma{}^{sk}\delta_r{}^{c} = \tfrac{1}{4}(\eta_\alpha{}^{mm_1}\eta_{\gamma ll_1}\varepsilon_{mm_1 dr}\varepsilon^{ll_1 sc} + \eta_\alpha{}^{mm_1}\eta_{\gamma ll_1}\varepsilon_{mm_1 kd}\varepsilon^{ll_1 sk}\delta_r{}^{c}) = \\
= 6\eta_\alpha{}^{mm_1}\eta_{\gamma ll_1}\delta_{[m}{}^{l}\delta_{m_1}{}^{l_1}\delta_d{}^{s}\delta_{r]}{}^{c} - \tfrac{3}{2}\eta_\alpha{}^{mm_1}\eta_{\gamma ll_1}\delta_{[m}{}^{l}\delta_{m_1}{}^{l_1}\delta_{d]}{}^{s}\delta_r{}^{c} = \\[1em]
= -g_{\alpha\gamma}\delta_r{}^{s}\delta_d{}^{c} + \eta_\alpha{}^{kc}\eta_{\gamma kd}\delta_r{}^{s} + \eta_\alpha{}^{ck}\eta_{\gamma kr}\delta_d{}^{s} + \eta_\alpha{}^{ks}\eta_{\gamma kr}\delta_d{}^{c} + \eta_\alpha{}^{sc}\eta_{\gamma dr}.
\end{gathered}\right| \tag{16.45}$$

$$= \eta_\alpha{}^{cs}\eta_{\gamma rd} + \eta_\alpha{}^{ck}\eta_{\gamma kr}\delta_d{}^{s} + \tfrac{1}{2}(\eta_\alpha{}^{ck}\eta_{\gamma dk}\delta_r{}^{s} + \eta_\alpha{}^{sk}\eta_{\gamma rk}\delta_d{}^{c}) - \tfrac{1}{4}g_{\alpha\gamma}\delta_r{}^{s}\delta_d{}^{c}. \tag{16.46}$$

Contract with $g^{\alpha\gamma}$

$$\begin{gathered}
A_{\alpha\beta d}{}^{c}A^{\alpha\beta}{}_{r}{}^{s} = \varepsilon_{rd}{}^{cs} + \varepsilon_{kr}{}^{ck}\delta_d{}^{s} + \tfrac{1}{2}(\varepsilon^{ck}{}_{dk}\delta_r{}^{s} + \varepsilon^{sk}{}_{rk}\delta_d{}^{c}) - \tfrac{3}{2}\delta_r{}^{s}\delta_d{}^{c} = \\
= \delta_r{}^{c}\delta_d{}^{s} - \delta_r{}^{s}\delta_d{}^{c} - 3\delta_r{}^{c}\delta_d{}^{s} + \tfrac{3}{2}\delta_d{}^{c}\delta_r{}^{s} + \tfrac{3}{2}\delta_d{}^{c}\delta_r{}^{s} - \tfrac{3}{2}\delta_r{}^{s}\delta_d{}^{c} = \tfrac{1}{2}\delta_r{}^{s}\delta_d{}^{c} - 2\delta_r{}^{c}\delta_d{}^{s}.
\end{gathered} \tag{16.47}$$

$$\begin{gathered}
A_{\alpha\beta d}{}^{c}A^{\lambda\mu}{}_{c}{}^{d} = \tfrac{1}{2}\eta_{[\alpha}{}^{ca}\eta_{\beta]}{}^{nn_1}\varepsilon_{adnn_1}\cdot\tfrac{1}{2}\eta^{[\lambda}{}_{ll_1}\eta^{\mu]}{}_{a_1 c}\varepsilon^{da_1 ll_1} = \\
= \tfrac{1}{4}\eta_{[\alpha}{}^{ca}\eta_{\beta]}{}^{nn_1}\eta^{[\lambda}{}_{ll_1}\eta^{\mu]}{}_{a_1 c}\cdot(-6\delta_{[a}{}^{a_1}\delta_n{}^{l}\delta_{n_1]}{}^{l_1}) = \\
= -\tfrac{1}{2}(\eta_{[\alpha}{}^{ca}\eta_{\beta]}{}^{nn_1}\eta^{[\lambda}{}_{nn_1}\eta^{\mu]}{}_{ac} - 2\eta_{[\alpha}{}^{ca}\eta_{\beta]}{}^{a_1 k_1}\eta^{[\lambda}{}_{ak_1}\eta^{\mu]}{}_{a_1 c}) = \\
= -2\delta_{[\alpha}{}^{\lambda}\delta_{\beta]}{}^{\mu} + (\eta_\alpha{}^{ca}\eta_\beta{}^{a_1 k_1}\eta^{[\lambda}{}_{ak_1}\eta^{\mu]}{}_{a_1 c} - \eta_\beta{}^{ca}\eta_\alpha{}^{a_1 k_1}\eta^{[\lambda}{}_{ak_1}\eta^{\mu]}{}_{a_1 c}) = \\
= 2\delta_{[\alpha}{}^{\mu}\delta_{\beta]}{}^{\lambda} + (\eta_\alpha{}^{ca}\eta_\beta{}^{a_1 k_1}\eta^{[\lambda}{}_{ak_1}\eta^{\mu]}{}_{a_1 c} - \eta_\beta{}^{a_1 k_1}\eta_\alpha{}^{ca}\eta^{[\lambda}{}_{k_1 a}\eta^{\mu]}{}_{ca_1}) = 2\delta_{[\alpha}{}^{\mu}\delta_{\beta]}{}^{\lambda}.
\end{gathered} \tag{16.48}$$

$$\begin{gathered}
A^{\alpha\beta}{}_{m}{}^{n}T_{\alpha\beta} = A^{\alpha\beta}{}_{m}{}^{n}A_{\alpha\beta k}{}^{l}T_l{}^{k} = \\
= (\tfrac{1}{2}\delta_m{}^{n}\delta_k{}^{l} - 2\delta_k{}^{n}\delta_m{}^{l})T_l{}^{k} = -2T_m{}^{n}, \quad T_{\alpha\beta} = -T_{\beta\alpha}.
\end{gathered} \tag{16.49}$$

16.8.2 Proof of the 4-vector identities

We need to prove (14.9).

$$e_{\alpha\beta\gamma\delta}=e_{[\alpha\beta\gamma\delta]}=A_{\alpha\beta b}{}^{a}A_{\gamma\delta d}{}^{c}e_{a}{}^{b}{}_{c}{}^{d},\quad e_{k}{}^{k}{}_{c}{}^{d}=e_{a}{}^{b}{}_{k}{}^{k}=0,\quad e_{a}{}^{b}{}_{c}{}^{d}=e_{c}{}^{d}{}_{a}{}^{b}. \tag{16.50}$$

Contract with $A^{\alpha\beta}{}_{m}{}^{n}A^{\gamma\delta}{}_{r}{}^{s}$

$$\begin{gathered}
A^{\alpha\beta}{}_{m}{}^{n}A^{\gamma\delta}{}_{r}{}^{s}A_{\alpha\beta b}{}^{a}A_{\gamma\delta d}{}^{c}e_{a}{}^{b}{}_{c}{}^{d}=\\
=(\tfrac{1}{2}\delta_{r}{}^{s}\delta_{d}{}^{c}-2\delta_{r}{}^{c}\delta_{d}{}^{s})(\tfrac{1}{2}\delta_{b}{}^{a}\delta_{m}{}^{n}-2\delta_{b}{}^{n}\delta_{m}{}^{a})e_{a}{}^{b}{}_{c}{}^{d}=4e_{m}{}^{n}{}_{r}{}^{s}=\\
=-A^{\beta\alpha}{}_{m}{}^{n}A^{\gamma\delta}{}_{r}{}^{s}A_{\gamma\alpha b}{}^{a}A_{\beta\delta d}{}^{c}e_{a}{}^{b}{}_{c}{}^{d}=\\
=-(\eta_{\gamma}{}^{an}\eta^{\beta}{}_{mb}+\eta_{\gamma}{}^{ak}\eta^{\beta}{}_{km}\delta_{b}{}^{n}+\tfrac{1}{2}\eta_{\gamma}{}^{ak}\eta^{\beta}{}_{bk}\delta_{m}{}^{n})\cdot\\
\cdot(\eta_{\beta}{}^{cs}\eta^{\gamma}{}_{rd}+\eta_{\beta}{}^{ct}\eta^{\gamma}{}_{tr}\delta_{d}{}^{s}+\tfrac{1}{2}\eta_{\beta}{}^{ct}\eta^{\gamma}{}_{dt}\delta_{r}{}^{s})e_{a}{}^{b}{}_{c}{}^{d}=\\
=-(\varepsilon_{rd}{}^{an}\varepsilon_{mb}{}^{cs}+\varepsilon_{mb}{}^{ck}\varepsilon_{kr}{}^{an}\delta_{d}{}^{s}+\tfrac{1}{2}\varepsilon_{mb}{}^{ck}\varepsilon_{dk}{}^{an}\delta_{r}{}^{s}+\\
+\varepsilon_{rd}{}^{ak}\varepsilon_{km}{}^{cs}\delta_{b}{}^{n}+\varepsilon^{ak}{}_{tr}\varepsilon^{ct}{}_{km}\delta_{d}{}^{s}\delta_{b}{}^{n}+\tfrac{1}{2}\varepsilon^{ak}{}_{dt}\varepsilon^{ct}{}_{km}\delta_{r}{}^{s}\delta_{b}{}^{n}+\\
+\tfrac{1}{2}\varepsilon_{rd}{}^{ak}\varepsilon_{bk}{}^{cs}\delta_{m}{}^{n}+\tfrac{1}{2}\varepsilon_{bk}{}^{ct}\varepsilon_{tr}{}^{ak}\delta_{m}{}^{n}\delta_{d}{}^{s}+\tfrac{1}{4}\varepsilon_{bk}{}^{ct}\varepsilon_{dt}{}^{ak}\delta_{m}{}^{n}\delta_{r}{}^{s})e_{a}{}^{b}{}_{c}{}^{d}=\\
=-(\delta_{r}{}^{a}\delta_{d}{}^{n}\delta_{m}{}^{c}\delta_{b}{}^{s}-\delta_{d}{}^{a}\delta_{r}{}^{n}\delta_{m}{}^{c}\delta_{b}{}^{s}+\delta_{d}{}^{a}\delta_{r}{}^{n}\delta_{b}{}^{c}\delta_{m}{}^{s}-\delta_{r}{}^{a}\delta_{d}{}^{n}\delta_{b}{}^{c}\delta_{m}{}^{s}+\\
+\delta_{m}{}^{c}\delta_{b}{}^{k}\delta_{k}{}^{a}\delta_{r}{}^{n}\delta_{d}{}^{s}-\delta_{m}{}^{k}\delta_{b}{}^{c}\delta_{k}{}^{a}\delta_{r}{}^{n}\delta_{d}{}^{s}+\delta_{m}{}^{k}\delta_{b}{}^{c}\delta_{k}{}^{n}\delta_{r}{}^{a}\delta_{d}{}^{s}-\delta_{m}{}^{c}\delta_{b}{}^{k}\delta_{k}{}^{n}\delta_{r}{}^{a}\delta_{d}{}^{s}+\\
+\tfrac{1}{2}\delta_{m}{}^{c}\delta_{b}{}^{k}\delta_{d}{}^{a}\delta_{k}{}^{n}\delta_{r}{}^{s}-\tfrac{1}{2}\delta_{m}{}^{k}\delta_{b}{}^{c}\delta_{d}{}^{a}\delta_{k}{}^{n}\delta_{r}{}^{s}+\tfrac{1}{2}\delta_{m}{}^{k}\delta_{b}{}^{c}\delta_{d}{}^{n}\delta_{k}{}^{a}\delta_{r}{}^{s}-\tfrac{1}{2}\delta_{m}{}^{c}\delta_{b}{}^{k}\delta_{d}{}^{n}\delta_{k}{}^{a}\delta_{r}{}^{s}+\\
+\delta_{r}{}^{a}\delta_{d}{}^{k}\delta_{k}{}^{c}\delta_{m}{}^{s}\delta_{b}{}^{n}-\delta_{r}{}^{k}\delta_{d}{}^{a}\delta_{k}{}^{c}\delta_{m}{}^{s}\delta_{b}{}^{n}+\delta_{r}{}^{k}\delta_{d}{}^{a}\delta_{k}{}^{s}\delta_{m}{}^{c}\delta_{b}{}^{n}-\delta_{r}{}^{a}\delta_{d}{}^{k}\delta_{k}{}^{s}\delta_{m}{}^{c}\delta_{b}{}^{n}+\\
+\delta_{m}{}^{a}\delta_{r}{}^{c}\delta_{d}{}^{s}\delta_{b}{}^{n}+2\delta_{m}{}^{c}\delta_{r}{}^{a}\delta_{d}{}^{s}\delta_{b}{}^{n}-\tfrac{1}{2}\delta_{m}{}^{a}\delta_{d}{}^{c}\delta_{r}{}^{s}\delta_{b}{}^{n}-\delta_{m}{}^{c}\delta_{d}{}^{a}\delta_{r}{}^{s}\delta_{b}{}^{n}+\\
+\tfrac{1}{2}\delta_{r}{}^{a}\delta_{d}{}^{k}\delta_{b}{}^{c}\delta_{k}{}^{s}\delta_{m}{}^{n}-\tfrac{1}{2}\delta_{r}{}^{k}\delta_{d}{}^{a}\delta_{b}{}^{c}\delta_{k}{}^{s}\delta_{m}{}^{n}+\tfrac{1}{2}\delta_{r}{}^{k}\delta_{d}{}^{a}\delta_{b}{}^{s}\delta_{k}{}^{c}\delta_{m}{}^{n}-\tfrac{1}{2}\delta_{r}{}^{a}\delta_{d}{}^{k}\delta_{b}{}^{s}\delta_{k}{}^{c}\delta_{m}{}^{n}-\\
-\tfrac{1}{2}\delta_{b}{}^{a}\delta_{r}{}^{c}\delta_{m}{}^{n}\delta_{d}{}^{s}-\delta_{b}{}^{c}\delta_{r}{}^{a}\delta_{m}{}^{n}\delta_{d}{}^{s}+\tfrac{1}{4}\delta_{b}{}^{a}\delta_{d}{}^{c}\delta_{m}{}^{n}\delta_{r}{}^{s}+\tfrac{1}{2}\delta_{b}{}^{c}\delta_{d}{}^{a}\delta_{m}{}^{n}\delta_{r}{}^{s})e_{a}{}^{b}{}_{c}{}^{d}=\\
=-(e_{r}{}^{s}{}_{m}{}^{n}-e_{t}{}^{s}{}_{m}{}^{t}\delta_{r}{}^{n}+e_{k}{}^{t}{}_{t}{}^{k}\delta_{r}{}^{n}\delta_{m}{}^{s}-e_{r}{}^{t}{}_{t}{}^{n}\delta_{m}{}^{s}-e_{m}{}^{t}{}_{t}{}^{s}\delta_{r}{}^{n}+e_{r}{}^{t}{}_{t}{}^{s}\delta_{m}{}^{n}-\\
-e_{r}{}^{n}{}_{m}{}^{s}+\tfrac{1}{2}e_{t}{}^{n}{}_{m}{}^{t}\delta_{r}{}^{s}-\tfrac{1}{2}e_{p}{}^{t}{}_{t}{}^{p}\delta_{m}{}^{n}\delta_{r}{}^{s}+\tfrac{1}{2}e_{m}{}^{t}{}_{t}{}^{n}\delta_{r}{}^{s}-e_{t}{}^{n}{}_{r}{}^{t}\delta_{m}{}^{s}+e_{t}{}^{n}{}_{m}{}^{t}\delta_{r}{}^{s}-\\
-e_{r}{}^{n}{}_{m}{}^{s}+e_{m}{}^{n}{}_{r}{}^{s}+2e_{r}{}^{n}{}_{m}{}^{s}-e_{t}{}^{n}{}_{m}{}^{t}\delta_{r}{}^{s}+\tfrac{1}{2}e_{r}{}^{t}{}_{t}{}^{s}\delta_{m}{}^{n}-\\
-\tfrac{1}{2}e_{t}{}^{p}{}_{p}{}^{t}\delta_{r}{}^{s}\delta_{m}{}^{n}+\tfrac{1}{2}e_{t}{}^{s}{}_{r}{}^{t}\delta_{m}{}^{n}-e_{r}{}^{t}{}_{t}{}^{s}\delta_{m}{}^{n}+\tfrac{1}{2}e_{t}{}^{p}{}_{p}{}^{t}\delta_{m}{}^{n}\delta_{r}{}^{s}).
\end{gathered} \tag{16.51}$$

$$e_{m}{}^{s}:=\tfrac{1}{3}e_{m}{}^{t}{}_{t}{}^{s}\quad\Rightarrow\quad 2e_{m}{}^{n}{}_{r}{}^{s}=2(e_{m}{}^{s}\delta_{r}{}^{n}+e_{r}{}^{n}\delta_{m}{}^{s})-(e_{r}{}^{s}\delta_{m}{}^{n}+e_{m}{}^{n}\delta_{r}{}^{s}). \tag{16.52}$$

16.8.3 Proof of the 6-vector identities

We need to prove (14.10).

$$\begin{gathered}
e_{\alpha\beta\gamma\delta\rho\sigma}=e_{[\alpha\beta\gamma\delta\rho\sigma]}=A_{\alpha\beta b}{}^{a}A_{\gamma\delta d}{}^{c}A_{\rho\sigma s}{}^{r}e_{a}{}^{b}{}_{c}{}^{d}{}_{r}{}^{s},\\
6e_{m}{}^{n}{}_{p}{}^{q}{}_{r}{}^{s}=2(e_{m}{}^{s}{}_{p}{}^{q}\delta_{r}{}^{n}+e_{r}{}^{n}{}_{p}{}^{q}\delta_{m}{}^{s})-(e_{r}{}^{s}{}_{p}{}^{q}\delta_{m}{}^{n}+e_{m}{}^{n}{}_{p}{}^{q}\delta_{r}{}^{s})=\\
=2(e_{m}{}^{q}{}_{r}{}^{s}\delta_{p}{}^{n}+e_{p}{}^{n}{}_{r}{}^{s}\delta_{m}{}^{q})-(e_{p}{}^{q}{}_{r}{}^{s}\delta_{m}{}^{n}+e_{m}{}^{n}{}_{r}{}^{s}\delta_{p}{}^{q}),\quad e_{m}{}^{s}{}_{p}{}^{q}:=e_{m}{}^{k}{}_{k}{}^{s}{}_{p}{}^{q}=e_{k}{}^{s}{}_{p}{}^{q}{}_{m}{}^{k}.
\end{gathered} \tag{16.53}$$

1. First contraction.
   Contract with $\delta_{q}{}^{p}$

$$\begin{gathered}
0=6e_{m}{}^{n}{}_{r}{}^{s}{}_{p}{}^{p}=2(e_{m}{}^{n}{}_{r}{}^{s}+e_{m}{}^{n}{}_{r}{}^{s})-(e_{p}{}^{p}{}_{r}{}^{s}\delta_{m}{}^{n}+4e_{m}{}^{n}{}_{r}{}^{s})=-(e_{p}{}^{p}{}_{r}{}^{s})\cdot\delta_{m}{}^{n}=0,\\
e_{p}{}^{p}{}_{r}{}^{s}=e_{r}{}^{s}{}_{p}{}^{p}=0.
\end{gathered} \tag{16.54}$$

2. Second contraction.
   Contract with $\delta_{n}{}^{r}$

$$\begin{gathered}
8e_{m}{}^{s}{}_{p}{}^{q}+2e_{t}{}^{t}{}_{p}{}^{q}\delta_{m}{}^{s}-e_{m}{}^{s}{}_{p}{}^{q}-e_{m}{}^{s}{}_{p}{}^{q}=2(e_{m}{}^{q}{}_{p}{}^{s}+e_{p}{}^{k}{}_{k}{}^{s}\delta_{m}{}^{q})-(e_{p}{}^{q}{}_{m}{}^{s}+e_{m}{}^{k}{}_{k}{}^{s}\delta_{p}{}^{q}),\\
6e_{m}{}^{s}{}_{p}{}^{q}+e_{p}{}^{q}{}_{m}{}^{s}-2e_{m}{}^{q}{}_{p}{}^{s}=2e_{p}{}^{t}{}_{t}{}^{s}\delta_{m}{}^{q}-e_{m}{}^{k}{}_{k}{}^{s}\delta_{p}{}^{q}.
\end{gathered} \tag{16.55}$$

(a) First contraction.
Contract with $\delta_q{}^m$

$$6e_k{}^s{}_p{}^k + e_p{}^k{}_k{}^s = 8e_p{}^t{}_t{}^s - e_p{}^k{}_k{}^s, \quad e_p{}^t{}_t{}^q = e_t{}^q{}_p{}^t. \tag{16.56}$$

(b) Second contraction.
Contract with $\delta_s{}^m$

$$10\tilde{e} := e_t{}^k{}_k{}^t, \quad -2e_k{}^q{}_p{}^k = 2e_p{}^t{}_t{}^q - e_t{}^k{}_k{}^t\delta_p{}^q, \quad e_p{}^t{}_t{}^q = \tfrac{5\tilde{e}}{2}\delta_p{}^q. \tag{16.57}$$

3. Third contraction.
Contract with $\delta_s{}^p$

$$\begin{gathered}2e_m{}^k{}_k{}^q\delta_r{}^n + 2e_r{}^n{}_m{}^q - e_r{}^k{}_k{}^q\delta_m{}^n - e_m{}^n{}_r{}^q = 2e_m{}^q{}_r{}^n + 2e_k{}^n{}_r{}^k\delta_m{}^q - e_k{}^q{}_r{}^k\delta_m{}^n - e_m{}^n{}_r{}^q,\\ e_r{}^n{}_m{}^q = e_m{}^q{}_r{}^n.\end{gathered} \tag{16.58}$$

$$\begin{gathered}7e_m{}^s{}_p{}^q - 2e_m{}^q{}_p{}^s = 5\tilde{e}\delta_m{}^q\delta_p{}^s - \tfrac{5\tilde{e}}{2}\delta_m{}^s\delta_p{}^q, \quad 9e_m{}^{[s}{}_p{}^{q]} = \tfrac{15\tilde{e}}{2}\delta_m{}^{[q}\delta_p{}^{s]},\\ 5e_m{}^s{}_p{}^q + 4e_m{}^{[s}{}_p{}^{q]} = 5\tilde{e}\delta_m{}^q\delta_p{}^s - \tfrac{5\tilde{e}}{2}\delta_m{}^s\delta_p{}^q, \quad 6e_m{}^s{}_p{}^q = \tilde{e}(4\delta_m{}^q\delta_p{}^s - \delta_m{}^s\delta_p{}^q),\\ e_m{}^n{}_r{}^s{}_p{}^q = \tilde{e}(2((4\delta_p{}^n\delta_r{}^q - \delta_p{}^q\delta_r{}^n)\delta_m{}^s + (4\delta_p{}^s\delta_m{}^q - \delta_p{}^q\delta_m{}^s)\delta_r{}^n) -\\ -((4\delta_p{}^n\delta_m{}^q - \delta_p{}^q\delta_m{}^n)\delta_r{}^s + (4\delta_p{}^s\delta_r{}^q - \delta_p{}^q\delta_r{}^s)\delta_m{}^n)).\end{gathered} \tag{16.59}$$

Calculate $\tilde{e}$

$$\begin{gathered}720 = e^{\alpha\beta\gamma\delta\mu\nu}e_{\alpha\beta\gamma\delta\mu\nu} = A^{\alpha\beta}{}_{b_1}{}^{a_1}A_{\alpha\beta b}{}^{a}A^{\gamma\delta}{}_{d_1}{}^{c_1}A_{\gamma\delta d}{}^{c}A^{\mu\nu}{}_{n_1}{}^{m_1}A_{\mu\nu n}{}^{m},\\ e_a{}^b{}_c{}^d{}_m{}^n e_{a_1}{}^{b_1}{}_{c_1}{}^{d_1}{}_{m_1}{}^{n_1} = 8(\tfrac{1}{4}\delta_{b_1}{}^{a_1}\delta_b{}^a - \delta_{b_1}{}^a\delta_b{}^{a_1})(\tfrac{1}{4}\delta_{d_1}{}^{c_1}\delta_d{}^c - \delta_{d_1}{}^c\delta_d{}^{c_1})\cdot\\ \cdot(\tfrac{1}{4}\delta_{n_1}{}^{m_1}\delta_n{}^m - \delta_{n_1}{}^m\delta_n{}^{m_1})e_a{}^b{}_c{}^d{}_m{}^n e_{a_1}{}^{b_1}{}_{c_1}{}^{d_1}{}_{m_1}{}^{n_1} = -8e_a{}^b{}_c{}^d{}_m{}^n e_b{}^a{}_d{}^c{}_n{}^m =\\ = -8\tilde{e}^2(8\delta_m{}^b\delta_c{}^n\delta_a{}^d - 4\delta_m{}^n\delta_c{}^b\delta_a{}^d + 8\delta_m{}^d\delta_a{}^n\delta_c{}^b - 4\delta_m{}^b\delta_a{}^n\delta_c{}^d + 2\delta_m{}^n\delta_a{}^b\delta_c{}^d - 4\delta_m{}^d\delta_c{}^n\delta_a{}^b)\\ (8\delta_n{}^a\delta_d{}^m\delta_b{}^c - 4\delta_n{}^m\delta_d{}^a\delta_b{}^c + 8\delta_n{}^c\delta_b{}^m\delta_d{}^a - 4\delta_n{}^a\delta_b{}^m\delta_d{}^c + 2\delta_n{}^m\delta_b{}^a\delta_d{}^c - 4\delta_n{}^c\delta_d{}^m\delta_b{}^a) =\\ = -720\cdot 64\tilde{e}^2.\end{gathered} \tag{16.60}$$

Whence,

$$\tilde{e} = \frac{1}{8}i. \tag{16.61}$$

16.8.4 Proof of the Bianchi identity

The Bianchi identity has the form

$$\begin{gathered}R_{\alpha\beta\gamma\delta}+R_{\alpha\gamma\delta\beta}+R_{\alpha\delta\beta\gamma}=0\\ (A_{\alpha\beta d}{}^{c}A_{\gamma\delta r}{}^{s}+A_{\alpha\gamma d}{}^{c}A_{\delta\beta r}{}^{s}+A_{\alpha\delta d}{}^{c}A_{\beta\gamma r}{}^{s})R_{c}{}^{d}{}_{s}{}^{r}=0.\end{gathered}\tag{16.62}$$

Let's contract it with $A^{\alpha\beta}{}_{t}{}^{l}A^{\gamma\delta}{}_{m}{}^{n}$

$$\left|\begin{array}{c}A_{\alpha\beta d}{}^{c}A_{\gamma\delta r}{}^{s}R_{c}{}^{d}{}_{s}{}^{r}A^{\alpha\beta}{}_{t}{}^{l}A^{\gamma\delta}{}_{m}{}^{n}R_{c}{}^{d}{}_{s}{}^{r}=\\ =(\frac{1}{2}\delta_{d}{}^{c}\delta_{t}{}^{l}-2\delta_{d}{}^{l}\delta_{t}{}^{c})(\frac{1}{2}\delta_{r}{}^{s}\delta_{m}{}^{n}-2\delta_{r}{}^{n}\delta_{m}{}^{s})R_{c}{}^{d}{}_{s}{}^{r}=4R_{t}{}^{l}{}_{m}{}^{n}\end{array}\right|$$

$$\left|\begin{array}{c}(A_{\alpha\gamma d}{}^{c}A_{\delta\beta r}{}^{s}+A_{\alpha\delta d}{}^{c}A_{\beta\gamma r}{}^{s})R_{c}{}^{d}{}_{s}{}^{r}A^{\alpha\beta}{}_{t}{}^{l}A^{\gamma\delta}{}_{m}{}^{n}=-2A_{\delta\beta r}{}^{s}A^{\delta\gamma}{}_{m}{}^{n}A_{\alpha\gamma d}{}^{c}A^{\alpha\beta}{}_{t}{}^{l}R_{c}{}^{d}{}_{s}{}^{r}=\\ =-2(\eta_{\beta}{}^{sn}\eta^{\gamma}{}_{mr}+\eta_{\beta}{}^{sk}\eta^{\gamma}{}_{km}\delta_{r}{}^{n}+\frac{1}{2}\eta_{\beta}{}^{sk}\eta^{\gamma}{}_{rk}\delta_{m}{}^{n})\cdot\\ \cdot(\eta_{\gamma}{}^{cl}\eta^{\beta}{}_{td}+\eta_{\gamma}{}^{ck_1}\eta^{\beta}{}_{k_1t}\delta_{d}{}^{l}+\frac{1}{2}\eta_{\gamma}{}^{ck_1}\eta^{\beta}{}_{dk_1}\delta_{t}{}^{l})R_{c}{}^{d}{}_{s}{}^{r}=\\ =-2(\varepsilon_{td}{}^{sn}\varepsilon_{mr}{}^{cl}+\varepsilon^{ck}{}_{mr}\varepsilon_{kt}{}^{sn}\delta_{d}{}^{l}+\frac{1}{2}\varepsilon_{mr}{}^{ck}\varepsilon_{dk}{}^{sn}\delta_{t}{}^{l}+\varepsilon_{td}{}^{sk}\varepsilon_{km}{}^{cl}\delta_{r}{}^{n}+\\ +\varepsilon^{ck_1}{}_{km}\varepsilon_{k_1t}{}^{sk}\delta_{d}{}^{l}\delta_{r}{}^{n}+\frac{1}{2}\varepsilon^{ck_1}{}_{km}\varepsilon_{dk_1}{}^{sk}\delta_{t}{}^{l}\delta_{r}{}^{n}+\frac{1}{2}(\varepsilon^{cl}{}_{rk}\varepsilon_{td}{}^{sk}\delta_{m}{}^{n}+\varepsilon^{sk}{}_{k_1t}\varepsilon^{ck_1}{}_{rk}\delta_{d}{}^{l}\delta_{m}{}^{n})+\\ +\frac{1}{4}\varepsilon^{ck_1}{}_{rk}\varepsilon^{sk}{}_{dk_1}\delta_{t}{}^{l}\delta_{m}{}^{n})R_{c}{}^{d}{}_{s}{}^{r}=-2(R_{m}{}^{n}{}_{t}{}^{l}+R_{k}{}^{r}{}_{r}{}^{k}\delta_{t}{}^{n}\delta_{m}{}^{l}-R_{m}{}^{k}{}_{k}{}^{l}\delta_{t}{}^{n}-R_{t}{}^{k}{}_{k}{}^{n}\delta_{m}{}^{l}-\\ -R_{m}{}^{l}{}_{t}{}^{n}-R_{k}{}^{l}{}_{m}{}^{k}\delta_{t}{}^{n}+R_{k}{}^{l}{}_{t}{}^{k}\delta_{m}{}^{n}+\frac{1}{2}(R_{m}{}^{k}{}_{k}{}^{n}\delta_{t}{}^{l}+R_{k}{}^{n}{}_{m}{}^{k}\delta_{t}{}^{l}-R_{k}{}^{r}{}_{r}{}^{k}\delta_{t}{}^{l}\delta_{m}{}^{n})+\\ +R_{m}{}^{k}{}_{k}{}^{n}\delta_{t}{}^{l}-R_{m}{}^{l}{}_{t}{}^{n}-R_{t}{}^{k}{}_{k}{}^{n}\delta_{m}{}^{l}+2R_{m}{}^{l}{}_{t}{}^{n}+R_{t}{}^{l}{}_{m}{}^{n}-\frac{1}{2}\cdot 2R_{m}{}^{k}{}_{k}{}^{n}\delta_{t}{}^{l}+\frac{1}{2}(R_{t}{}^{k}{}_{k}{}^{l}\delta_{m}{}^{n}+\\ +R_{k}{}^{l}{}_{t}{}^{k}\delta_{m}{}^{n}-R_{k}{}^{r}{}_{r}{}^{k}\delta_{m}{}^{n}\delta_{t}{}^{l})-R_{t}{}^{k}{}_{k}{}^{l}\delta_{m}{}^{n}+\frac{1}{2}R_{k}{}^{r}{}_{r}{}^{k}\delta_{t}{}^{l}\delta_{m}{}^{n}=-2(2R_{m}{}^{n}{}_{t}{}^{l}-\\ -2R_{k}{}^{l}{}_{m}{}^{k}\delta_{t}{}^{n}-2R_{t}{}^{k}{}_{k}{}^{n}\delta_{m}{}^{l}+R_{k}{}^{l}{}_{t}{}^{k}\delta_{m}{}^{n}+R_{m}{}^{k}{}_{k}{}^{n}\delta_{t}{}^{l}+R_{k}{}^{r}{}_{r}{}^{k}\delta_{t}{}^{n}\delta_{m}{}^{l}-\frac{1}{2}R_{k}{}^{r}{}_{r}{}^{k}\delta_{m}{}^{n}\delta_{t}{}^{l})\end{array}\right|$$

$$4R_{k}{}^{l}{}_{m}{}^{k}\delta_{t}{}^{n}+4R_{t}{}^{k}{}_{k}{}^{n}\delta_{m}{}^{l}-2R_{k}{}^{l}{}_{t}{}^{k}\delta_{m}{}^{n}-2R_{m}{}^{k}{}_{k}{}^{n}\delta_{t}{}^{l}-2R_{k}{}^{r}{}_{r}{}^{k}\delta_{t}{}^{n}\delta_{m}{}^{l}+R_{k}{}^{r}{}_{r}{}^{k}\delta_{m}{}^{n}\delta_{t}{}^{l}=0.\tag{16.63}$$

Contract with $\delta_{n}{}^{t}$

$$\begin{gathered}16R_{k}{}^{l}{}_{m}{}^{k}+4R_{t}{}^{k}{}_{k}{}^{t}\delta_{m}{}^{l}-2R_{k}{}^{l}{}_{m}{}^{k}-2R_{m}{}^{k}{}_{k}{}^{l}-8R_{k}{}^{r}{}_{r}{}^{k}\delta_{m}{}^{l}+R_{k}{}^{r}{}_{r}{}^{k}\delta_{m}{}^{l}=0,\\ R_{k}{}^{l}{}_{m}{}^{k}=\frac{1}{4}R_{k}{}^{r}{}_{r}{}^{k}\delta_{m}{}^{l}=-\frac{1}{8}R\delta_{m}{}^{l}\end{gathered}\tag{16.64}$$

that finishes the proof.

16.8.5 Proof of the identities related to the Weyl tensor

Since,

$$\begin{gathered}T_{[\alpha\beta]}=\frac{1}{4}\eta_{\alpha}{}^{aa_1}\eta_{\beta}{}^{bb_1}\cdot\frac{3}{2}(T_{a[a_1bb_1]}-T_{[bb_1a]a_1})=\\ =\frac{1}{4}\eta_{\alpha}{}^{aa_1}\eta_{\beta}{}^{bb_1}\cdot\frac{1}{2}(T_{ak}{}^{kd}\varepsilon_{a_1dbb_1}-T^{kd}{}_{ka_1}\varepsilon_{adbb_1})=\frac{1}{2}\eta_{\alpha}{}^{ca}\eta_{\beta}{}^{bk_1}\varepsilon_{adbk_1}\cdot\\ \cdot\frac{1}{4}(T^{kd}{}_{kc}-T_{kc}{}^{kd})=A_{\alpha\beta d}{}^{c}\cdot\frac{1}{4}(T^{kd}{}_{kc}-T_{kc}{}^{kd})\end{gathered}\tag{16.65}$$

then it will be true the following relation

$$\begin{gathered}R_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]}=A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\frac{1}{16}(P^{kdnr}\varepsilon_{kcns}-P_{kc}{}^{nr}\varepsilon^{kd}{}_{ns}+P_{kcns}\varepsilon^{kdnr}-P^{kd}{}_{ns}\varepsilon_{kc}{}^{nr})=\\ =A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\frac{1}{16}(P^{kd}{}_{mm_1}\frac{1}{2}\varepsilon^{mm_1nr}\varepsilon_{kcns}+P_{kc}{}^{mm_1}\frac{1}{2}\varepsilon_{mm_1ns}\varepsilon^{kdnr}-\\ -P_{kc}{}^{nr}(\delta_{n}{}^{k}\delta_{s}{}^{d}-\delta_{n}{}^{d}\delta_{s}{}^{k})-P^{kd}{}_{ns}(\delta_{k}{}^{n}\delta_{c}{}^{r}-\delta_{k}{}^{r}\delta_{c}{}^{n}))=\\ =A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\frac{1}{16}(P^{kd}{}_{kc}\delta_{s}{}^{r}-P^{kd}{}_{ks}\delta_{c}{}^{r}+P^{rd}{}_{cs}+\\ +P_{kc}{}^{kd}\delta_{s}{}^{r}-P_{kc}{}^{kr}\delta_{s}{}^{d}+P_{sc}{}^{dr}-P_{kc}{}^{kr}\delta_{s}{}^{d}+P_{sc}{}^{dr}-P^{kd}{}_{ks}\delta_{c}{}^{r}+P^{rd}{}_{cs})=\\ =A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\frac{1}{4}(P_{sc}{}^{dr}-\frac{1}{2}P^{kd}{}_{ks}\delta_{c}{}^{r}+\frac{1}{2}P_{kc}{}^{kd}\delta_{s}{}^{r}-\frac{1}{2}P_{kc}{}^{kr}\delta_{s}{}^{d})=\\ =A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\frac{1}{4}(P_{sc}{}^{dr}-\frac{1}{2}R\delta_{s}{}^{d}\delta_{c}{}^{r}+\frac{1}{4}R\delta_{s}{}^{r}\delta_{c}{}^{d}).\end{gathered}\tag{16.66}$$

Similarly,

$$\begin{gathered} g_{[\alpha}{}^{[\gamma} g_{\beta]}{}^{\delta]} = A_{\alpha\beta d}{}^{c} A^{\gamma\delta}{}_{r}{}^{s} \tfrac{1}{4}(\varepsilon_{sc}{}^{dr} - \tfrac{1}{2}\varepsilon^{kd}{}_{ks}\delta_{c}{}^{r} + \tfrac{1}{2}\varepsilon^{kd}{}_{kc}\delta_{s}{}^{r} - \tfrac{1}{2}\varepsilon_{kc}{}^{kr}\delta_{s}{}^{d}) = \\ = A_{\alpha\beta d}{}^{c} A^{\gamma\delta}{}_{r}{}^{s} \tfrac{1}{4}(\tfrac{1}{2}\delta_{s}{}^{r}\delta_{c}{}^{d} - 2\delta_{s}{}^{d}\delta_{c}{}^{r}). \end{gathered} \tag{16.67}$$

Thus, from

$$C_{\alpha\beta}{}^{\gamma\delta} := R_{\alpha\beta}{}^{\gamma\delta} - R_{[\alpha}{}^{[\gamma}\, g_{\beta]}{}^{\delta]} + 1/10 R g_{[\alpha}{}^{[\gamma}\, g_{\beta]}{}^{\delta]}, \tag{16.68}$$

it follows

$$\begin{gathered} C_{\alpha\beta}{}^{\gamma\delta} = A_{\alpha\beta d}{}^{c} A^{\gamma\delta}{}_{r}{}^{s}(R_{c}{}^{d}{}_{s}{}^{r} - \tfrac{1}{4}P_{sc}{}^{dr} + \tfrac{1}{8}R\delta_{s}{}^{d}\delta_{c}{}^{r} - \tfrac{1}{16}R\delta_{s}{}^{r}\delta_{c}{}^{d} + \tfrac{1}{40}R(\tfrac{1}{2}\delta_{s}{}^{r}\delta_{c}{}^{d} - \\ -2\delta_{s}{}^{d}\delta_{c}{}^{r})) = A_{\alpha\beta d}{}^{c} A^{\gamma\delta}{}_{r}{}^{s}(R_{(c}{}^{d}{}_{s)}{}^{r} + R_{[c}{}^{k}{}_{|k|}{}^{[r}\delta_{s]}{}^{d]} + \tfrac{1}{40}R(3\delta_{s}{}^{d}\delta_{c}{}^{r} - 2\delta_{s}{}^{r}\delta_{c}{}^{d})). \end{gathered} \tag{16.69}$$

Then

$$C_{\alpha\beta}{}^{\gamma\delta} = A_{\alpha\beta d}{}^{c} A^{\gamma\delta}{}_{r}{}^{s}(R_{(c}{}^{d}{}_{s)}{}^{r} + \tfrac{1}{80}R\delta_{s}{}^{d}\delta_{c}{}^{r}). \tag{16.70}$$

Contract it with $A^{\alpha\beta}{}_{k}{}^{l} A_{\gamma\delta t}{}^{n}$

$$\begin{gathered} 4C_{k}{}^{l}{}_{t}{}^{n} := A^{\alpha\beta}{}_{k}{}^{l} A_{\gamma\delta t}{}^{n} C_{\alpha\beta}{}^{\gamma\delta} = A^{\alpha\beta}{}_{k}{}^{l} A_{\alpha\beta d}{}^{c} A^{\gamma\delta}{}_{r}{}^{s} A_{\gamma\delta t}{}^{n}(R_{(c}{}^{d}{}_{s)}{}^{r} + \tfrac{1}{80}R\delta_{s}{}^{d}\delta_{c}{}^{r}) = \\ = (\tfrac{1}{2}\delta_{k}{}^{l}\delta_{d}{}^{c} - 2\delta_{k}{}^{c}\delta_{d}{}^{l})(\tfrac{1}{2}\delta_{r}{}^{s}\delta_{t}{}^{n} - 2\delta_{r}{}^{n}\delta_{t}{}^{s})(\tfrac{1}{2}R_{c}{}^{d}{}_{s}{}^{r} + \tfrac{1}{2}R_{s}{}^{d}{}_{c}{}^{r} + \tfrac{1}{80}R\delta_{s}{}^{d}\delta_{c}{}^{r}) = \\ = (2R_{k}{}^{l}{}_{t}{}^{n} + \tfrac{1}{8}R_{k_1}{}^{r_1}{}_{r_1}{}^{k_1}\delta_{k}{}^{l}\delta_{t}{}^{n} - \tfrac{1}{2}R_{k_1}{}^{l}{}_{k}{}^{k_1}\delta_{t}{}^{n} - \tfrac{1}{2}R_{t}{}^{k_1}{}_{k_1}{}^{n}\delta_{k}{}^{l} + \\ +2R_{t}{}^{l}{}_{k}{}^{n} + \tfrac{1}{80}R(\delta_{t}{}^{n}\delta_{k}{}^{l} - \delta_{t}{}^{n}\delta_{k}{}^{l} - \delta_{t}{}^{n}\delta_{k}{}^{l} + 4\delta_{t}{}^{l}\delta_{k}{}^{n}) = \\ = (4R_{(k}{}^{l}{}_{t)}{}^{n} - \tfrac{1}{16}R\delta_{k}{}^{l}\delta_{t}{}^{n} + \tfrac{1}{16}R\delta_{k}{}^{l}\delta_{t}{}^{n} + \tfrac{1}{16}R\delta_{t}{}^{n}\delta_{k}{}^{l} + \tfrac{1}{80}R(4\delta_{t}{}^{l}\delta_{k}{}^{n} - \delta_{t}{}^{n}\delta_{k}{}^{l})) = \\ = 4(R_{(k}{}^{l}{}_{t)}{}^{n} + \tfrac{1}{40}R\delta_{(k}{}^{l}\delta_{t)}{}^{n}), \end{gathered} \tag{16.71}$$

that finishes the proof.

16.8.6 Proof of the relations associated with the metric induced in the cross section of the cone K_6

Let the cone section of K_6

$$T^2 + V^2 - W^2 - X^2 - Y^2 - Z^2 = 0 \tag{16.72}$$

by the plane V+W=1 be set. We make a stereographic projection of the resulting hyperboloid onto the plane (V = 0, W = 1).

$$\begin{gathered} \tfrac{t}{T} = \tfrac{x}{X} = \tfrac{y}{Y} = \tfrac{z}{Z} = -\tfrac{\frac{1}{2}}{V-\frac{1}{2}}, \quad x^2 + y^2 + z^2 - t^2 = \tfrac{1}{2V-1}, \\ T = -t(2V-1), \quad X = -x(2V-1), \quad Y = -y(2V-1), \quad Z = -z(2V-1), \\ dT = -(dt\ (2V-1) + 2tdV), \quad dX = -(dx\ (2V-1) + 2xdV), \\ dY = -(dy\ (2V-1) + 2ydV), \quad dZ = -(dz\ (2V-1) + 2zdV), \\ dS^2 = dT^2 + dV^2 - dW^2 - dX^2 - dY^2 - dZ^2 = |dV = -dW| = \\ = dT^2 - dX^2 - dY^2 - dZ^2 = (dt^2 - dx^2 - dy^2 - dz^2)(2V-1)^2 + \\ +4(tdt - xdx - ydy - zdz)dV(2V-1) + 4(t^2 - x^2 - y^2 - z^2)dV^2 = \\ = |tdt - xdx - ydy - zdz = -\tfrac{1}{2}d(\tfrac{1}{2V-1})| = \tfrac{dt^2 - dx^2 - dy^2 - dz^2}{(t^2 - x^2 - y^2 - z^2)^2}. \end{gathered} \tag{16.73}$$

We make the substitution

$$\begin{gathered}\varsigma=\tfrac{-iX+Y}{2V-1}=-y+ix,\quad \omega=-\tfrac{i(T+Z)}{2V-1}=i(t+z),\quad \eta=\tfrac{i(Z-T)}{2V-1}=i(t-z),\\ \varsigma\bar{\varsigma}+\eta\omega=\tfrac{1}{2V-1},\\ d\varsigma=(\tfrac{dY}{2V-1}-\tfrac{2YdV}{(2V-1)^2})-i(\tfrac{dX}{2V-1}-\tfrac{2XdV}{(2V-1)^2}),\quad d\bar{\varsigma}=(\tfrac{dY}{2V-1}-\tfrac{2YdV}{(2V-1)^2})+i(\tfrac{dX}{2V-1}-\tfrac{2XdV}{(2V-1)^2}),\\ d\omega=-i(\tfrac{dT}{2V-1}-\tfrac{2TdV}{(2V-1)^2}+\tfrac{dZ}{2V-1}-\tfrac{2ZdV}{(2V-1)^2}),\quad d\eta=-i(\tfrac{dT}{2V-1}-\tfrac{2TdV}{(2V-1)^2}-\tfrac{dZ}{2V-1}+\tfrac{2ZdV}{(2V-1)^2}),\\ d\varsigma d\bar{\varsigma}+d\omega d\eta=\tfrac{(dX^2+dY^2+dZ^2-dT^2)}{(2V-1)^2}-4\tfrac{dV}{(2V-1)^3}(XdX+YdY+ZdZ-TdT)+\\ +4\tfrac{dV^2}{(2V-1)^4}(X^2+Y^2+Z^2-T^2)=-\tfrac{dT^2-dX^2-dY^2-dZ^2}{(T^2-X^2-Y^2-Z^2)^2},\\ ds^2:=dT^2-dX^2-dY^2-dZ^2=-\tfrac{d\varsigma d\bar{\varsigma}+d\eta d\omega}{(\varsigma\bar{\varsigma}+\eta\omega)^2},\quad x^2+y^2+z^2-t^2=\tfrac{1}{2V-1}.\end{gathered}\tag{16.74}$$

Therefore, there is a reason to set

$$X:=\begin{pmatrix}\omega & \varsigma\\ -\bar{\varsigma} & \eta\end{pmatrix},\quad dX:=\begin{pmatrix}d\omega & d\varsigma\\ -d\bar{\varsigma} & d\eta\end{pmatrix},\quad \frac{\partial}{\partial X}:=\begin{pmatrix}\frac{\partial}{\partial\omega} & \frac{\partial}{\partial\varsigma}\\ -\frac{\partial}{\partial\bar{\varsigma}} & \frac{\partial}{\partial\eta}\end{pmatrix}.\tag{16.75}$$

Then

$$ds^2=-\frac{det(dX)}{(det(X))^2},\quad \bar{X}^T+X=0.\tag{16.76}$$

16.8.7 Proof of the first invariant relations

Consider the fractional linear group L

$$\tilde{X}=(AX+B)(CX+D)^{-1},\quad S:=\begin{pmatrix}A & B\\ C & D\end{pmatrix},\quad detS=1.\tag{16.77}$$

Let there be two consecutive transformations

$$\tilde{X}=(AX+B)(CX+D)^{-1},\quad \tilde{\tilde{X}}=(\tilde{A}\tilde{X}+\tilde{B})(\tilde{C}\tilde{X}+\tilde{D})^{-1},\tag{16.78}$$

then

$$\begin{gathered}\tilde{\tilde{X}}=(\tilde{\tilde{A}}X+\tilde{\tilde{B}})(\tilde{\tilde{C}}X+\tilde{\tilde{D}})^{-1}=\\ =(\tilde{A}(AX+B)+\tilde{B}(CX+D))(\tilde{C}(AX+B)+\tilde{D}(CX+D))^{-1}=\\ =((\tilde{A}A+\tilde{B}C)X+\tilde{A}B+\tilde{B}D)((\tilde{C}A+\tilde{D}C)X+\tilde{C}B+\tilde{D}D)^{-1},\\ S:=\begin{pmatrix}A & B\\ C & D\end{pmatrix},\quad \tilde{S}:=\begin{pmatrix}\tilde{A} & \tilde{B}\\ \tilde{C} & \tilde{D}\end{pmatrix},\quad \tilde{\tilde{S}}:=\begin{pmatrix}\tilde{\tilde{A}} & \tilde{\tilde{B}}\\ \tilde{\tilde{C}} & \tilde{\tilde{D}}\end{pmatrix},\quad \tilde{\tilde{S}}=\tilde{S}S.\end{gathered}\tag{16.79}$$

For the unitary fractional-linear transformations, we have

$$\begin{gathered}X^*+X\equiv\tilde{X}^*+\tilde{X}=0,\\ 0=(AX+B)(CX+D)^{-1}+(CX+D)^{*-1}(AX+B)^*,\\ 0=(CX+D)^*(AX+B)+(AX+B)^*(CX+D)=\\ =X^*(A^*C+C^*A)X+X^*(A^*D+C^*B)+(B^*C+D^*A)X+D^*B+B^*D\equiv\\ \equiv X^*+X.\end{gathered}\tag{16.80}$$

Whence,

$$\begin{gathered}A^*C+C^*A=0,\quad B^*D+D^*B=0,\quad A^*D+C^*B=E,\quad S^*\hat{E}S=\hat{E},\\ \hat{E}:=\begin{pmatrix}0 & E\\ E & 0\end{pmatrix}.\end{gathered}\tag{16.81}$$

Let

$$X = YZ^{-1}, \quad \tilde{Y} = AY + BZ, \quad \tilde{Z} = CY + DZ. \tag{16.82}$$

Set

$$X := \begin{pmatrix} x_1 & x_2 \\ x_3 & x_4 \end{pmatrix}, \quad \hat{X} := \begin{pmatrix} \hat{x}_1 & \hat{x}_2 \\ \hat{x}_3 & \hat{x}_4 \end{pmatrix}, \quad dX := \begin{pmatrix} dx_1 & dx_2 \\ dx_3 & dx_4 \end{pmatrix}, \quad \hat{d}X := \begin{pmatrix} d\hat{x}_1 & d\hat{x}_2 \\ d\hat{x}_3 & d\hat{x}_4 \end{pmatrix},$$

$$\tfrac{\partial}{\partial X} := \begin{pmatrix} \dfrac{\partial}{\partial x_1} & \dfrac{\partial}{\partial x_2} \\ \dfrac{\partial}{\partial x_3} & \dfrac{\partial}{\partial x_4} \end{pmatrix}, \quad \tfrac{\partial}{\partial \hat{X}} := \begin{pmatrix} \dfrac{\partial}{\partial \hat{x}_1} & \dfrac{\partial}{\partial \hat{x}_2} \\ \dfrac{\partial}{\partial \hat{x}_3} & \dfrac{\partial}{\partial \hat{x}_4} \end{pmatrix}. \tag{16.83}$$

Then

$$\begin{gathered} dX + d\hat{X} = d(X + \hat{X}), \\ d(X\hat{X}) = d \begin{pmatrix} x_1\hat{x}_1 + x_2\hat{x}_3 & x_1\hat{x}_2 + x_2\hat{x}_1 \\ x_3\hat{x}_1 + x_4\hat{x}_3 & x_3\hat{x}_2 + x_4\hat{x}_4 \end{pmatrix} = \\ = \begin{pmatrix} dx_1\hat{x}_1 + dx_2\hat{x}_3 & dx_1\hat{x}_2 + dx_2\hat{x}_1 \\ dx_3\hat{x}_1 + dx_4\hat{x}_3 & dx_3\hat{d}x_2 + x_4\hat{x}_4 \end{pmatrix} + \begin{pmatrix} x_1d\hat{x}_1 + x_2d\hat{x}_3 & x_1d\hat{x}_2 + x_2d\hat{x}_1 \\ x_3d\hat{x}_1 + x_4d\hat{x}_3 & x_3d\hat{x}_2 + x_4d\hat{x}_4 \end{pmatrix} = \\ = (dX)\ \hat{X} + X\ (d\hat{X}). \end{gathered} \tag{16.84}$$

Therefore, the identities

$$\hat{\hat{X}} = AX + B \ \ \Rightarrow \ \ d\hat{\hat{X}} = AdX, \qquad \hat{\hat{X}} = X^{-1} \ \ \Rightarrow \ \ d\hat{\hat{X}} = -X^{-1}\ dX\ X^{-1}. \tag{16.85}$$

are true. The proof of the second is that

$$\begin{gathered} \hat{\hat{X}}X = 1, \ \ (d\hat{\hat{X}})\ X + \hat{\hat{X}}\ (dX) = 0, \\ (d\hat{\hat{X}})\ X + X^{-1}\ (dX) = 0, \ \ d\hat{\hat{X}} = -X^{-1}\ (dX)\ X^{-1}. \end{gathered} \tag{16.86}$$

Then we will obtain the identity chain

$$\begin{gathered} \tilde{X}^* + \tilde{X} = 0, \ \ -\tilde{X}^* = \tilde{X}, \\ -(CX + D)^{*-1}(AX + B)^* = (AX + B)(CX + D)^{-1}, \\ -(AX + B)^* = (CX + D)^*(AX + B)(CX + D)^{-1}. \end{gathered} \tag{16.87}$$

Multiply both parts by CdX

$$\begin{gathered} (-X^*A^* - B^*)CdX = (CX + D)^*(AX + B)(CX + D)^{-1}CdX, \\ (-X^*A^*C - B^*C)dX = (CX + D)^*(AX + B)(CX + D)^{-1}CdX, \\ (X^*C^*A + D^*A - E)dX = (CX + D)^*(AX + B)(CX + D)^{-1}CdX, \\ (CX + D)^*AdX - (CX + D)^*(AX + B)(CX + D)^{-1}CdX = dX, \\ A(dX)(CX + D)^{-1} - (AX + B)(CX + D)^{-1}C(dX)(CX + D)^{-1} = \\ = (CX + D)^{*-1}(dX)(CX + D)^{-1}, \\ (d(AX + B))\ (CX + D)^{-1} + (AX + B)d((CX + D)^{-1}) = \\ = (CY + DZ)^{*-1}Z^*(dX)Z(CY + DZ)^{-1}, \\ \tilde{Z}^*\ d\tilde{X}\tilde{Z} = Z^*\ dXZ. \end{gathered} \tag{16.88}$$

Thus, the first invariant turns out.

16.8.8 Proof of the second invariant relations

Let

$$\hat{X} = AX + B, \tag{16.89}$$

or in the componentwise record

$$\begin{aligned} \hat{x}_1 &= a_1x_1 + a_2x_3 + b_1, \quad & \hat{x}_2 &= a_1x_2 + a_2x_4 + b_2, \\ \hat{x}_3 &= a_3x_1 + a_4x_3 + b_3, \quad & \hat{x}_4 &= a_3x_2 + a_4x_4 + b_4. \end{aligned} \tag{16.90}$$

Then

$$\begin{aligned} \tfrac{\partial}{\partial x_1} &= a_1\tfrac{\partial}{\partial \hat{x}_1} + a_3\tfrac{\partial}{\partial \hat{x}_3}, \qquad & \tfrac{\partial}{\partial x_2} &= a_1\tfrac{\partial}{\partial \hat{x}_2} + a_3\tfrac{\partial}{\partial \hat{x}_4}, \\ \tfrac{\partial}{\partial x_3} &= a_2\tfrac{\partial}{\partial \hat{x}_1} + a_4\tfrac{\partial}{\partial \hat{x}_3}, \qquad & \tfrac{\partial}{\partial x_4} &= a_2\tfrac{\partial}{\partial \hat{x}_2} + a_4\tfrac{\partial}{\partial \hat{x}_4}, \end{aligned} \tag{16.91}$$

or in the abbreviated form

$$\tfrac{\partial}{\partial X} = A^T\tfrac{\partial}{\partial \hat{X}} \quad \Rightarrow \quad \tfrac{\partial}{\partial \hat{X}^T} = \tfrac{\partial}{\partial X^T}A^{-1} \ (det(A) \neq 0). \tag{16.92}$$

Now let

$$\hat{X} = X^{-1}, \tag{16.93}$$

or in the componentwise record

$$\hat{x}_1 = \tfrac{x_4}{x_1x_4-x_2x_3} \quad \hat{x}_2 = -\tfrac{x_2}{x_1x_4-x_2x_3}, \quad \hat{x}_3 = -\tfrac{x_3}{x_1x_4-x_2x_3} \quad \hat{x}_4 = \tfrac{x_1}{x_1x_4-x_2x_3}. \tag{16.94}$$

Then

$$\begin{aligned} \tfrac{\partial}{\partial x_1} &= -\tfrac{x_4x_4}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_1} + \tfrac{x_2x_4}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_2} + \tfrac{x_3x_4}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_3} - \tfrac{x_2x_3}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_4}, \\ \tfrac{\partial}{\partial x_2} &= \tfrac{x_4x_3}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_1} - \tfrac{x_1x_4}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_2} - \tfrac{x_3x_3}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_3} + \tfrac{x_1x_3}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_4}, \\ \tfrac{\partial}{\partial x_3} &= \tfrac{x_4x_2}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_1} - \tfrac{x_2x_2}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_2} - \tfrac{x_1x_4}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_3} + \tfrac{x_1x_2}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_4}, \\ \tfrac{\partial}{\partial x_4} &= -\tfrac{x_2x_3}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_1} + \tfrac{x_1x_2}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_2} + \tfrac{x_1x_3}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_3} - \tfrac{x_1x_1}{(x_1x_4-x_2x_3)^2}\tfrac{\partial}{\partial \hat{x}_4}, \end{aligned} \tag{16.95}$$

or in the abbreviated form

$$\tfrac{\partial}{\partial X} = -X^{-1T}\tfrac{\partial}{\partial \hat{X}}X^{-1T} \quad \Leftrightarrow \quad \tfrac{\partial}{\partial \hat{X}^T} = -X\tfrac{\partial}{\partial X^T}X. \tag{16.96}$$

Assume that $det(C) \neq 0$, then

$$\begin{gathered} \tilde{X} = (AX+B)(CX+D)^{-1} = AC^{-1} + (B - AC^{-1}D)(CX+D)^{-1}, \\ \tfrac{\partial}{\partial \tilde{X}^T} = \tfrac{\partial}{\partial (CX+D)^{T-1}}(B - AC^{-1}D)^{-1} = \\ = -(CX+D)\tfrac{\partial}{\partial (CX+D)^T}(CX+D)(B - AC^{-1}D)^{-1} = \\ = -(CX+D)\tfrac{\partial}{\partial X^T}C^{-1}(CX+D)(B - AC^{-1}D)^{-1} = \\ = -(CX+D)\tfrac{\partial}{\partial X^T}(XC^* + C^{-1}DC^*)(BC^* - AC^{-1}DC^*)^{-1} = \\ = (CX+D)\tfrac{\partial}{\partial X^T}(CX+D)^*(BC^* + AD^*)^{-1} = (CX+D)\tfrac{\partial}{\partial X^T}(CX+D)^*. \end{gathered} \tag{16.97}$$

Whence,

$$\tilde{Z}^{-1}\frac{\partial}{\partial \tilde{X}^T}\tilde{Z}^{*-1} = Z^{-1}\frac{\partial}{\partial X^T}Z^{*-1}. \tag{16.98}$$

If now $det(C) = 0$ then set

$$\tilde{\tilde{X}} = \tilde{X}^{-1} = (CX + D)(AX + B)^{-1}, \quad \tilde{\tilde{Z}} = AY + BZ, \quad \tilde{\tilde{C}} = A, \quad det(\tilde{\tilde{C}}) \neq 0. \tag{16.99}$$

If $det(A) \neq 0$ then

$$\begin{gathered} Z^{-1}\frac{\partial}{\partial\tilde{X}^T}Z^{*-1} = \tilde{\tilde{Z}}^{-1}\frac{\partial}{\partial\tilde{\tilde{X}}^T}\tilde{\tilde{Z}}^{*-1} = \\ = -(AY + BZ)^{-1}\tilde{X}\frac{\partial}{\partial\tilde{X}^T}\tilde{X}(AY + BZ)^{*-1} = |\tilde{X}^* = -\tilde{X}| = \\ = (AY + BZ)^{-1}(AY + BZ)(CY + DZ)^{-1}\frac{\partial}{\partial\tilde{X}^T}\times \\ \times(CY + DZ)^{*-1}(AY + BZ)^{*}(AY + BZ)^{*-1} = \tilde{Z}^{-1}\frac{\partial}{\partial\tilde{X}^T}\tilde{Z}^{*-1}. \end{gathered} \tag{16.100}$$

If $det(A) = 0$ then set

$$\tilde{\tilde{X}} = \tilde{X} + \tilde{D}, \quad \tilde{\tilde{\tilde{X}}} = \tilde{\tilde{X}}^{-1}, \quad \tilde{\tilde{\tilde{X}}} = (\tilde{\tilde{\tilde{A}}}X + \tilde{\tilde{\tilde{B}}})(\tilde{\tilde{\tilde{C}}}X + \tilde{\tilde{\tilde{D}}})^{-1}, \quad \tilde{\tilde{\tilde{C}}} = A + \tilde{D}C, \quad det(\tilde{\tilde{\tilde{C}}}) \neq 0. \tag{16.101}$$

It is obvious that $\tilde{D}$ can always be chosen so that $det(\tilde{\tilde{\tilde{C}}}) \neq 0$. So

$$Z^{-1}\frac{\partial}{\partial X^T}Z^{*-1} = \tilde{\tilde{\tilde{Z}}}^{-1}\frac{\partial}{\partial\tilde{\tilde{\tilde{X}}}^T}\tilde{\tilde{\tilde{Z}}}^{*-1} = \tilde{\tilde{Z}}^{-1}\frac{\partial}{\partial\tilde{\tilde{X}}^T}\tilde{\tilde{Z}}^{*-1} = \tilde{Z}^{-1}\frac{\partial}{\partial\tilde{X}^T}\tilde{Z}^{*-1}, \tag{16.102}$$

that will give the second invariant.

16.8.9 Proof of the octave alternative identity

Let us prove the identity (14.95). If $P_{\Psi A} = \eta_{\Psi XA}X^X$ then

$$\begin{gathered} \tfrac{1}{2}\eta_{(\Lambda\Psi)}{}^{\Theta}\eta_{\Theta\Phi}{}^{\Upsilon} = \eta_{(\Lambda}{}^{AB}P_{\Psi)A}P^{\Theta}{}_{B}\eta_{\Theta CD}P_{\Phi}{}^{C}P^{\Upsilon D} = \\ = (-\eta_{(\Lambda}{}^{AB}\eta_{\Psi)AX} + 2\eta_{(\Lambda}{}^{AB}\eta_{\Psi)(AX)})X^{X}P^{\Theta}{}_{B}\eta_{\Theta CD}P_{\Phi}{}^{C}P^{\Upsilon D} = \\ = -\tfrac{1}{2}g_{\Lambda\Psi}X^{B}X^{Y}\varepsilon_{YBCD}P_{\Phi}{}^{C}P^{\Upsilon D} + \tfrac{1}{4\sqrt{2}}\eta_{(\Psi}{}^{K}{}_{|K}\eta_{\Theta\Phi}{}^{\Upsilon}P^{\Theta}{}_{B|}P_{\Lambda)}{}^{B} = \\ = -\tfrac{1}{2}g_{\Lambda\Psi}\delta_{\Phi}{}^{\Upsilon} + \tfrac{1}{4\sqrt{2}}\eta_{(\Psi}{}^{K}{}_{|K|}\eta_{\Lambda)\Phi}{}^{\Upsilon}, \end{gathered}$$

$$\begin{gathered} \tfrac{1}{2}\eta_{(\Lambda|\Theta|}{}^{\Upsilon}\eta_{\Psi)\Phi}{}^{\Theta} = \eta_{(\Lambda}{}^{AB}\eta_{\Psi)}{}^{CD}P_{\Theta A}P^{\Upsilon}{}_{B}P_{\Phi C}P^{\Theta}{}_{D} = \\ = (-\eta_{(\Lambda}{}^{AB}\eta_{\Psi)A}{}^{C} + 2\eta_{(\Lambda}{}^{AB}\eta_{\Psi)K}{}^{K}\cdot\tfrac{1}{8}\delta_{A}{}^{C})P^{\Upsilon}{}_{B}P_{\Phi C} = \\ = -\tfrac{1}{2}g_{\Lambda\Psi}\delta_{\Phi}{}^{\Upsilon} + \tfrac{1}{4}\eta_{(\Lambda}{}^{CB}\eta_{\Psi)K}{}^{K}P^{\Upsilon}{}_{B}P_{\Phi C} = -\tfrac{1}{2}g_{\Lambda\Psi}\delta_{\Phi}{}^{\Upsilon} + \tfrac{1}{4\sqrt{2}}\eta_{(\Psi}{}^{K}{}_{|K|}\eta_{\Lambda)\Phi}{}^{\Upsilon}, \end{gathered}$$

$$\begin{gathered} \tfrac{1}{2}\eta_{\Lambda\Theta}{}^{\Phi}\eta_{(\Upsilon\Psi)}{}^{\Theta} = \eta_{\Lambda}{}^{AB}P_{\Theta A}P^{\Phi}{}_{B}\eta_{(\Psi}{}^{CD}P_{\Upsilon)C}P^{\Theta}{}_{D} = \eta_{\Lambda D}{}^{B}\eta_{(\Psi}{}^{CD}P_{\Upsilon)C}P^{\Phi}{}_{B} = \\ = \eta_{\Lambda D}{}^{B}(-\eta_{(\Psi}{}^{CD}\eta_{\Upsilon)CX}X^{X} + \tfrac{1}{4}P_{(\Psi}{}^{D}\eta_{\Upsilon)K}{}^{K})P^{\Phi}{}_{B} = -\tfrac{1}{2}g_{\Upsilon\Psi}\delta_{\Lambda}{}^{\Phi} + \tfrac{1}{4\sqrt{2}}\eta_{\Lambda(\Psi}{}^{\Phi}\eta_{\Upsilon)K}{}^{K}, \end{gathered}$$

$$\begin{gathered} \tfrac{1}{2}\eta_{\Lambda(\Psi}{}^{\Theta}\eta_{|\Theta|\Upsilon)}{}^{\Phi} = \eta_{\Lambda}{}^{AB}P_{(\Psi|A}P^{\Theta}{}_{B}\eta_{\Theta|}{}^{CD}P_{\Upsilon)C}P^{\Phi}{}_{D} = \eta_{\Lambda}{}^{AB}P_{(\Psi|A|}P_{\Upsilon)C}\varepsilon_{XB}{}^{CD}X^{X}P^{\Phi}{}_{D} = \\ = (\eta_{\Lambda}{}^{CD}X^{A} - \eta_{\Lambda X}{}^{B}X^{X}\varepsilon^{A}{}_{B}{}^{CD})P_{(\Psi|A|}P_{\Upsilon)C}P^{\Phi}{}_{D} = -\tfrac{1}{2}g_{\Upsilon\Psi}\delta_{\Lambda}{}^{\Phi} + \tfrac{1}{4\sqrt{2}}\eta_{\Lambda(\Psi}{}^{\Phi}\eta_{\Upsilon)K}{}^{K}. \end{gathered} \tag{16.103}$$

16.8.10 Proof of the central Moufang identity

Let us prove the identity (14.96). Note that $r^{\Phi}\eta_{\Phi\Theta}{}^{\Omega}$ and $\eta_{\Omega\Upsilon}{}^{\Gamma}r^{\Upsilon}$ are orthogonal transformations.

$$\begin{gathered}
r^{\Phi}\eta_{\Phi\Theta}{}^{\Omega}\eta_{\Lambda\Psi}{}^{\Theta}\eta_{\Omega\Upsilon}{}^{\Gamma}r^{\Upsilon} = r^{\Phi}\eta_{\Phi\Lambda}{}^{\Theta}\eta_{\Theta\Omega}{}^{\Gamma}\eta_{\Psi\Upsilon}{}^{\Omega}r^{\Upsilon},\\
r^{\Phi}\eta_{\Phi\Lambda}{}^{\Theta}\eta_{\Theta\Delta}{}^{\Gamma} = r^{\Phi}\eta_{\Phi\Theta}{}^{\Omega}\eta_{\Lambda\Psi}{}^{\Theta}\eta_{\Omega\Upsilon}{}^{\Gamma}r^{\Upsilon}\eta^{\Psi}{}_{\Xi\Delta}r^{\Xi}.
\end{gathered} \tag{16.104}$$

Since $r^{\Lambda} = \eta^{\Lambda}{}_{AB}X^{A}Y^{B} = P^{\Lambda}{}_{B}Y^{B}$, then

$$\begin{gathered}
R_S{}^{Q}P^{\Theta}{}_{Q}\eta_{\Theta}{}^{CD} = \underbrace{R_B{}^{K}}_{[=\varepsilon_B{}^{KMN}X_MY_N]}\underbrace{P^{\Lambda}{}_{S}\eta_{\Lambda}{}^{AB}}_{[=X^P\varepsilon_{PS}{}^{AB}]}\underbrace{P^{\Omega}{}_{K}\eta_{\Omega}{}^{VD}Y_V}_{[=\varepsilon_K{}^{LDV}X_LY_V]}\underbrace{P^{\Psi}{}_{A}\eta_{\Psi}{}^{YC}Y_Y}_{[\varepsilon_A{}^{XCY}X_XY_Y]},\\
X_XY_Y\varepsilon_S{}^{QXY}\varepsilon_{RQ}{}^{CD}X^R = \varepsilon_B{}^{KMN}X_MY_NX^P\varepsilon_{PS}{}^{AB}\varepsilon_K{}^{LDV}X_LY_V\varepsilon_A{}^{XCY}X_XY_Y.
\end{gathered} \tag{16.105}$$

1. Right path.

$$\begin{gathered}
\varepsilon_B{}^{KMN}X_MY_NX^P\varepsilon_{PS}{}^{AB}\varepsilon_K{}^{LDV}X_LY_V\varepsilon_A{}^{XCY}X_XY_Y =\\
= X^P\varepsilon_{PS}{}^{AB}\varepsilon_A{}^{XCY}X_XY_Y\varepsilon_B{}^{KMN}X_MY_N\varepsilon_K{}^{LDV}X_LY_V =\\
= X^P\varepsilon_{PS}{}^{AB}\varepsilon_A{}^{XCY}X_XY_Y(-\varepsilon^{KMN}{}_B + \delta_B{}^{M}\varepsilon^{KN})X_MY_N\varepsilon_K{}^{LDV}X_LY_V =\\
= X^P\varepsilon_{PS}{}^{AB}\varepsilon_A{}^{XCY}X_XY_Y(-\varepsilon^{DVN}{}_BY_VY_N + \varepsilon^{NLDV}X_BY_NY_VX_L) =\\
= X^P(\varepsilon_{PS}\varepsilon^{AB} + \delta_P{}^{A}\delta_S{}^{B} - \delta_P{}^{B}\delta_S{}^{A} - \varepsilon^B{}_S{}^A{}_P)\varepsilon_A{}^{XCY}X_XY_Y\\
(-Y^DY_B + \delta_B{}^{D} + X_LY^LY^DX_B - X^DX_B) =\\
= (X^S\varepsilon^{BXCY}X_XY_Y + \delta_S{}^{B}Y^C - X^B\varepsilon_S{}^{XCY}X_XY_Y - \varepsilon^B{}_S{}^{CY}Y_Y)\\
(-Y^DY_B + \delta_B{}^{D} + X_LY^LY^DX_B - X^DX_B) =\\
= -X_SY^DX_XY_BY_Y\varepsilon^{BXCY} + X_S\varepsilon^{DXCY}Y_YX_X + X_SY^LX_LY^DX_B\varepsilon^{BXCY}X_XY_Y -\\
- X_SX^DX_B\varepsilon^{BXCY}X_XY_Y - Y^CY^DY_S + Y^C\delta_S{}^{D} + Y^CY^DX_SX_LY^L - Y^CX^DX_S +\\
+ X^B\varepsilon_S{}^{XCY}X_XY_YY^DY_B - X^D\varepsilon_S{}^{XCY}X_XY_Y - Y^LX_LY^DX_BX^B\varepsilon_S{}^{XCY}X_XY_Y +\\
+ X^BX_BX^D\varepsilon_S{}^{XCY}X_XY_Y + Y^DY_B\varepsilon^B{}_S{}^{CY}Y_Y - \varepsilon^D{}_S{}^{CY}Y_Y -\\
- Y^LX_LY^DX_B\varepsilon^B{}_S{}^{CY}Y_Y + X^DX_B\varepsilon^B{}_S{}^{CY}Y_Y =\\
= -X_SY^DX_X(\underbrace{Y^XY^C}_{1} - \varepsilon^{CX}) + X_S\varepsilon^{DXCY}Y_YX_X + \underbrace{X_SY^CY^DY^LX_L}_{2} -\\
- \underbrace{X_SX^DY^C}_{3} - \underbrace{Y^CY^DY_S}_{4} + Y_C\delta_S{}^{D} + \underbrace{Y^CY^DX_SX_LY^L}_{1} - Y^CX^DX_S +\\
+ \underbrace{X^BY_BY^D\varepsilon_S{}^{XCY}X_XY_Y}_{2} - \underbrace{X^D\varepsilon_S{}^{XCY}X_XY_Y}_{3} - \underbrace{2X^LY_LY^D\varepsilon_S{}^{XCY}X_XY_Y}_{2} +\\
+ 2\underbrace{X^D\varepsilon_S{}^{XCY}X_XY_Y}_{3} + Y^D(\underbrace{Y_SY^C}_{4} - \delta_S{}^{C}) - \varepsilon^D{}_S{}^{CY}Y_Y -\\
- \underbrace{Y^LX_LY^DX_B\varepsilon^B{}_S{}^{CY}Y_Y}_{1} + \underbrace{X^DX_B\varepsilon^B{}_S{}^{CY}Y_Y}_{3} =\\
= X_S\varepsilon^{DXCY}Y_YX_X + X_SY^DX^C - X_SX^DY^C - \varepsilon^D{}_S{}^{CY}Y_Y + Y^C\delta_S{}^{D} - Y^D\delta_S{}^{C} =\\
= X_S\varepsilon^{CDXY}Y_YX_X - \varepsilon^{CD}{}_S{}^{Y}Y_Y.
\end{gathered} \tag{16.106}$$

2. Left path.

$$\begin{gathered}
X_XY_Y\varepsilon_S{}^{QXY}\varepsilon_{RQ}{}^{CD}X^R = (\delta_S{}^{X}\varepsilon^{QY} - \varepsilon^{XQ}{}_S{}^{Y})\varepsilon_{RQ}{}^{CD}X^RX_XY_Y =\\
= X_SY^QX^R\varepsilon_{RQ}{}^{CD} - \varepsilon_S{}^{YCD}Y_Y.
\end{gathered} \tag{16.107}$$

This is direct proof. The simpler proof based on the formula (10.1, case 2).

16.8.11 Proof of the A-operator identities for n = 8

Let us prove the identity (14.99).

1. First.

$$\begin{gathered}
\eta_{\Lambda}{}^{KA}\eta_{\Psi K}{}^{B}\eta^{\Lambda}{}_{LC}\eta^{\Psi L}{}_{D} = \varepsilon_{LC}{}^{KA}\varepsilon_{K}{}^{BL}{}_{D} = \\
= (\varepsilon_{LC}\varepsilon^{KA} - \delta_{L}{}^{A}\delta_{C}{}^{K} + \varepsilon^{K}{}_{LC}{}^{A})(\delta_{K}{}^{B}\delta_{D}{}^{L} - \varepsilon_{KD}\varepsilon^{BL} + \varepsilon_{K}{}^{L}{}_{D}{}^{B}) = \\
= \varepsilon_{CD}\varepsilon^{AB} - \delta_{D}{}^{A}\delta_{C}{}^{B} + \varepsilon^{B}{}_{DC}{}^{A} - \delta_{D}{}^{A}\delta_{C}{}^{B} + \varepsilon_{CD}\varepsilon^{AB} - \varepsilon_{D}{}^{B}{}_{C}{}^{A} + \\
+ \varepsilon^{A}{}_{CD}{}^{B} - \varepsilon_{C}{}^{A}{}_{D}{}^{B} + 4\varepsilon_{C}{}^{A}{}_{D}{}^{B} = 2(\varepsilon_{CD}\varepsilon^{AB} + \delta_{C}{}^{A}\delta_{D}{}^{B} - \delta_{D}{}^{A}\delta_{C}{}^{B}).
\end{gathered} \tag{16.108}$$

2. Second.

$$\begin{gathered}
\eta_{\Lambda}{}^{KA}\eta_{\Psi K}{}^{B}\eta^{\Lambda}{}_{CL}\eta^{\Psi}{}_{D}{}^{L} = \varepsilon_{CL}{}^{KA}\varepsilon_{K}{}^{B}{}_{D}{}^{L} = (\delta_{C}{}^{K}\delta_{L}{}^{A} - \varepsilon^{K}{}_{LC}{}^{A})(\varepsilon_{KD}\varepsilon^{BL} - \varepsilon_{K}{}^{L}{}_{D}{}^{B}) = \\
= \varepsilon_{CD}\varepsilon^{AB} - \varepsilon_{D}{}^{B}{}_{C}{}^{A} - \varepsilon_{C}{}^{A}{}_{D}{}^{B} + 4\varepsilon_{C}{}^{A}{}_{D}{}^{B} = \varepsilon_{CD}\varepsilon^{AB} + 2\varepsilon_{C}{}^{A}{}_{D}{}^{B}.
\end{gathered} \tag{16.109}$$

3. Third.
   On the basis of the identity (8.27)

$$\eta_{\Lambda}{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}\eta^{\Theta}{}_{CB} = 2(g_{\Lambda\Psi}g^{\Omega\Theta} + \delta_{\Lambda}{}^{\Theta}\delta_{\Psi}{}^{\Omega} - \delta_{\Psi}{}^{\Theta}\delta_{\Lambda}{}^{\Omega}), \tag{16.110}$$

   we obtain

$$\begin{gathered}
\eta_{[\Lambda}{}^{C}{}_{|A|}\eta_{\Psi]CB}\eta_{\Upsilon K}{}^{A}\eta_{\Omega}{}^{KB} = 4g_{\Upsilon[\Lambda}g_{\Psi]\Omega}, \\
\eta_{[\Lambda}{}^{C}{}_{|A|}\eta_{\Psi]CB}\eta_{\Upsilon}{}^{A}{}_{K}\eta_{\Omega}{}^{BK} = 4g_{\Upsilon[\Lambda}g_{\Psi]\Omega} + 8\eta_{[\Upsilon}g_{\Omega][\Lambda}\eta_{\Psi]}, \\
\eta_{[\Lambda|A|}{}^{C}\eta_{\Psi]BC}\eta_{\Upsilon}{}^{A}{}_{K}\eta_{\Omega}{}^{BK} = 4g_{\Upsilon[\Lambda}g_{\Psi]\Omega}, \\
\eta_{[\Lambda|A|}{}^{C}\eta_{\Psi]BC}\eta_{\Upsilon K}{}^{A}\eta_{\Omega}{}^{KB} = 4g_{\Upsilon[\Lambda}g_{\Psi]\Omega} + 8\eta_{[\Upsilon}g_{\Omega][\Lambda}\eta_{\Psi]}.
\end{gathered} \tag{16.111}$$

In addition,

$$\begin{gathered}
\eta_{[\Lambda}{}^{C}{}_{|A|}\eta_{\Psi]CB}\eta_{\Upsilon K}{}^{A}\eta_{\Omega}{}^{KB} = 8g_{\Upsilon[\Lambda}g_{\Psi]\Omega} + \eta_{\Upsilon}{}^{C}{}_{A}\eta_{[\Psi|CB|}\eta_{\Lambda]}{}^{KB}\eta_{\Omega K}{}^{A} = \\
= 8g_{\Upsilon[\Lambda}g_{\Psi]\Omega} + (\eta_{[\Lambda}{}^{BK}\eta_{\Psi]BC} + 2\eta_{[\Lambda}{}^{K}{}_{|C|}\eta_{\Psi]})(\eta_{[\Upsilon|A|}{}^{C}\eta_{\Omega]}{}^{A}{}_{K} + 2\eta_{[\Omega|K|}{}^{C}\eta_{\Upsilon]}) = \\
= 8g_{\Upsilon[\Lambda}g_{\Psi]\Omega} - \eta_{[\Lambda}{}^{C}{}_{|A|}\eta_{\Psi]CB}\eta_{\Upsilon K}{}^{A}\eta_{\Omega}{}^{KB} - 2\eta_{[\Psi}{}^{K}{}_{|C|}\eta_{\Lambda]}\eta_{[\Upsilon|A|}{}^{C}\eta_{\Omega]}{}^{A}{}_{K} + \\
+ 2\eta_{[\Omega|K|}{}^{C}\eta_{\Upsilon]}\eta_{[\Lambda}{}^{BK}\eta_{\Psi]BC} - 16\eta_{[\Lambda}g_{\Psi][\Omega}\eta_{\Upsilon]}.
\end{gathered} \tag{16.112}$$

References

[1] К. В. Андреев [K.V. Andreev]: Спинорный формализм и геометрия шестимерных римановых пространств [Spinornyĭ formalizm i geometriya shestimernykh rimanovykh prostranstv]. Кандидатская диссертация [Kandidatskaya dissertatsiya], Уфа [Ufa], 1997, [in Russian: Spinor formalism and the geometry of six-dimensional Riemannian spaces. Ph. D. Thesis], arXiv:1204.0194v1.

[2] К. В. Андреев [K.V. Andreev]: О структуре тензора кривизны 6-мерных римановых пространств [O strukture tenzora krivizny 6-mernykh rimanovykh prostranstv]. Вестик Башкирского унверситета [Vestnik Bashkirskogo Universiteta], 2(1996)44-47, [in Russian: On the structure of the curvature tensor of 6-dimensional Riemannian spaces].

[3] К. В. Андреев [K.V. Andreev]: О спинорном формализме при n=6 [O spinornom formalizme pri n=6]. Изв. Высш. Учебн. Завед., Матем. [Izv. Vyssh. Uchebn. Zaved., Mat.], 1(2001)11-23 (The article is freely available online at the Mathnet site: `http://mi.mathnet.ru/eng/ivm838`, the two figures of the article are missing in the electronic version, however). English translation: K. V. Andreyev: On spinor formalism for n = 6. Russian Mathematics (Iz. VUZ), 45:1(2001)9-20.]

[4] К. В. Андреев [K.V. Andreev]: О структурных константах алгебры октав. Уравнение Клифформа [O strukturnykh konstantakh algebry oktav. Uravnenie Klifforda]. Изв. Высш. Учебн. Завед., Матем. [Izv. Vyssh. Uchebn. Zaved. Mat.], 3(2001)3-6. English translation: K. V. Andreyev: Structure constants of the algebra of octaves. The Clifford equation. Russian Mathematics (Iz. VUZ), 45:3(2001)1-4 (Mathnet URL: `http://mi.mathnet.ru/ivm856`).

[5] К. В. Андреев [K.V. Andreev]: О твисторах 6-мерного пространства [O tvistorakh 6-mernogo prostranstva]. ВИНИТИ-2469-В-98 [VINITI-2469-V-98], август [Aug] 1998, 23 с [23pp. Paper deponed on Aug 3, 1998 at VINITI (Moscow), ref. №2469-V 98], [in Russian: On twistors of 6-dimensional space].

[6] К. В. Андреев [K.V. Andreev]: Принцип тройственности для двух квадрик [Printsip troĭstvennosti dlya dvykh kvadrik]. ВИНИТИ-2470-В-98 [VINITI-2470-V-98], август [Aug] 1998. 19 с. [19pp. Paper deponed on Aug 3, 1998 at VINITI (Moscow), ref. №2470-V 98.], [in Russian: Triality principle for two quadrics].

[7] К. В. Андреев [K.V. Andreev]: О внутренних геометриях многообразия плоских образующих 6-мерной квадрики [O vnutrennikh geometriyakh mnogoobraziya ploskikh obrazuyushchikh 6-mernoĭ kvadriki], Изв. Высш. Учебн. Завед., Матем. [Izv. Vyssh. Uchebn. Zaved. Mat.], 6(1998)3-8. English translation: K. V. Andreyev: On intrinsic geometries of the manifold of plane generators of a 6-dimensional quadric, Izv. Vyssh. Uchebn. Zaved. Mat., 42:6(1998)1-6 (Mathnet URL: `http://mi.mathnet.ru/ivm436`).

[8] John C. Baez: The octonions, Bull. Amer. Math. Soc. (N.S.) 39(2002)145–205, arXiv:math.RA/0105155v4. Russian translation: Баэз Джон С. [Baèz Dzhon S]: Октонионы [Oktoniony]. Гиперкомплексные числа в геометрии и физике [Giperkomleksnye chisla v geometrii i fizike], 5:1(vol. 3)(2006)120-177.

[9] Р.Ф. Билялов, Б.С. Никитин: Спиноры в произвольных реперах. Ковариантная производная и производная Ли спиноров [Spinory v proizvol'nykh reperakh. Kovariantnaya proizvodnaya Li spinorov]. Изв. Высш. Учебн. Завед., Матем. [Izv. Vyssh. Uchebn. Zaved. Mat.], 6(1998)9-19. English translation: R. F. Bilyalov, B. S. Nikitin. Spinors in arbitrary frames. The covariant derivative and the Lie derivative of spinors. Izv. Vyssh. Uchebn. Zaved. Mat., 42:6(1998)7-15 (Mathnet URL: `http://mi.mathnet.ru/ivm440`).

[10] Berger, Marcel: Geometry. II. Translated from the French by M. Cole and S. Levy. Universitext. Springer-Verlag, Berlin, 1987. Berger, Marcel: Geometry. I. Translated from the French by M. Cole and S. Levy. Universitext. Springer-Verlag, Berlin, 1987. Russian translation: М. Бергер [M. Berger]. Геометрия [Geometriya], т. 1, 2 [v. 1, 2]. Мир [Mir], Москва [Moskva], 1984. Russian translation by Ю.Н. Сударев [Yu.N. Sudarev], А.В. Пажитнов [A.V. Pazhitnov] and С.В. Чмутов [S.V. Chmutov] under edition И.Х. Сабитов [I.Kh. Sabitov].

[11] H.A. Buchdahl: On the calculus of for-spinors. Proceedings of the Royal Society of London, Series A, Mathematical and Physical Sciences, 303(1968)355-378 (DOI:10.1098/rspa.1968.0055)

[12] Dietmar Salamon: Spin Geometry and Seiberg-Witten invariants. Warwick, 1996.

[13] F. S. Klotz: Twistors and the conformal group. Journal of Mathematical Phisics, 15(1974)2242-2247, (DOI:10.1063/1.1666606).

[14] А.П. Котельников [A.P. Kotel'nikov]: Винтовое счисление и некоторые приложения его к геометрии и механике [Vintovoe schislenie i nekotorye prilozheniya ego k geometrii i mekhanike]. Типо-литография Императорского Университета [Tipo-litografiya Imperatorskogo Universiteta], Казань [Kazan'], 1895. [in Russian: Screw Calculus and Some of Its Applications to Geometry and Mechanics]. Second reprint: А.П. Котельников [A.P. Kotel'nikov]: Винтовое счисление и некоторые приложения его к геометрии и механике [Vintovoe schislenie i nekotorye prilozheniya ego k geometrii i mekhanike]. КомКнига [Komkniga], Москва [Moskva], 2006.

[15] Э.Г. Нейфельд [È.G. Neĭfel'd]: Об инволюциях в комплексных пространствах [Ob involyutsiyakh v kompleksnykh prostranstvakh]. Тр. Геом. Семин. [Tr. Geom. Semin.], Казанский университет [Kazanskiĭ universitet] (Выпуск [Vypusk]) 19(1989)71-82 (Mathnet URL: `http://mi.mathnet.ru/eng/kutgs98`). [in Russian: Involutions in complex spaces].

[16] Э.Г. Нейфельд [È.G. Neĭfel'd]: Нормализация комплексных грассманианов и квадрик [Normalizatsiya kompleksnykh grassmanianov i kvadrik]. Тр. Геом. Семин. [Tr. Geom. Semin.], Казанский университет [Kazanskiĭ universitet] (Выпуск [Vypusk]) 20(1990)58-69 (Mathnet URL: `http://mi.mathnet.ru/kutgs82`). [in Russian: Normalization of complex Grassmannians and quadrics].

[17] А.П. Норден [A.P. Norden]: О комплексном представлении тензоров пространства Лоренца [O kompleksnom predstavlenii tenzorov prostranstva Lorentsa]. Изв. Высш. Учебн. Завед., Матем. [Izv. Vyssh. Uchebn. Zaved., Mat.] (1959) No. 1 (8), 156-164 (Mathnet URL: `http://mi.mathnet.ru/eng/ivm2415`). [in Russian: On a complex representation of the tensors of Lorentz space].

[18] М.М. Постников. [M.M. Postnikov] Группы и алгебры Ли [Gruppy i algebry Li]. Наука [Nauka], Москва [Moskva], 1986. [in Russian: M. Postnikov: Lie Groups and Lie Algebras. Lectures in Geometry, Semester 5. Mir, Moscow, 1986; URSS Publishing, Moscow, 1994]. The main ideas of the hypercomplex number constraction on the base of the Bott periodicity are given in the lectures 13-16.

[19] Б.А. Розенфельд [B.A. Rozenfel'd]: Проективная геометрия как метрическая геометрия [Proektivnaya geometriya kak metricheskaya geometriya]. Труды семинара по векторному и тензорному анализу с их приложениями к геометрии, механике и физике [Trudy seminara po vektornomu i tenzornomu analizu s ikh prilozheniyami k geometrii, mekhanike, fizike], Москва [Moskva], 8(1950)328-354, [in Russian: Projective geometry as the metric geometry. Proceedings of the Seminar on Vector and tensor analysis with their applications.]

[20] Б.А. Розенфельд [B.A. Rozenfel'd]: Неевклидовы геометрии [Neevklidovy geometrii]. ГИТТО [GITTO], Москва [Moskva], 1955. [in Russian: Non-Euclidean Geometries]. The Cartan triality principle is given on the page 534.

[21] A. Zee: Quantum field theory in a nutshell. Princeton university press, 2003. Russian translation: Зи Энтони [Zi Èntoni]. Квантовая теория в двух словах [Kvantovaya teoriya v dvukh slovakh]. РХД [RKHD], Москва [Moskva], Ижевск [Izhevsk], 2009. Russian translation by В.Г. Войткевич [V.G. Voĭtkevich] and Ю.В. Колесниченко [Yu.V. Kolesnichenko] under edition И.В. Полюбина [I.V. Polyubina].

[22] R. Penrose, W. Rindler: Spinors and Space-Time. Vol. 1: Two-Spinor Calculus and Relativistic Fields. Vol. 2: Spinor and Twistor Methods in Space-Time Geometry. Cambridge Monogr. Math. Phys. Cambridge University Press, Cambridge, Vol. 1: 1984, Vol. 2: 1986. Russian translation: Р. Пенроуз [R. Penrouz], В. Риндлер [V. Rindler]: Спиноры и пространство-время [Spinory i prostranstvo-vremya]. Мир [Mir], Москва [Moskva], т. 1 [Tom 1]: 1987, т. 2 [Tom 2]: 1988. Russian translation by Е.М. Серебрянный [E.M. Serebryannyĭ] and З.А. Штейнгард [Z.A. Sheyĭngard] under edition Д.М. Гальцов [D.M. Gal'tsov].

[23] Д.М.Синцов [D.M. Sintsov]: Теория коннексов в пространстве в связи с теорией дифференциальных уравнений в частных производных первого порядка. [Teoriya konneksov v prostranstve v svyazi s teorieĭ differentsial'nykh uravneniĭ v chastnykh proizvodnykh pervogo poryadka]. Типо-литография Императорского Университета [Tipo-litografiya Imperatorskogo Universiteta], Казань [Kazan'], 1894. [in Russian: Theory of connexes in space in relation to the theory of first order partial differential equations].

[24] Ю.П. Степановский [Yu. P. Stepanpovskiĭ]: Спиноры 6-мерного пространства и их применение к описанию поляризованных частиц со спином 1/2 [Spinory 6-mernogo prostranstva i ikh primenenie k opisaniyu polyarizovannykh chastits so spinom 1/2]. Проблемы Ядерной Физики и Космических Лучей [Problemy Yadernoĭ Fiziki i Kosmicheskikh Lucheĭ], Харьков [Khar'kov], 4(1976) 9-21, [in Russian: Spinors of a sixdimensional space and their application to the description of polarized particles with spin 1/2].

[25] Ю.П. Степановський [Yu. P. Stepanpovskiĭ]. Алгебра матриць Дірака у шестімірному вигляді [Algebra matrits Diraka u shestimirnomu vyglyadi]. Український Фізичний Журнал [Ukrains'kyĭ Fizychnyĭ Zhurnal]. Київ [Kyiv], т. XI, 8(1968) 813-824, in Ukraine: Yu. P. Stepanovskii: Algebra of Dirac matrices in six-dimensional form. Ukrainian Journal of Physics, Kiev, v.XI, 8(1968) 813-824.

[26] J.A. Schouten, J. Haantjes: Konforme Feldtheorie II; R_6 und Spinuraum [Conformal field theory II and spin space]. Annali della R.Scuola Normale Superiore di Piza, Scienze Fisiche e Matematiche, 2. Series, 4(1935) 175-189.

[27] Dale Husemoller: Fibre bundles. McGraw-Hill Book Company, 1966. Russian translation: Д. Хьюзмоллер [D. Kh'yuzmoller]. Расслоенные пространства [Rassloennye prostranstva]. Мир [Mir], Москва [Moskva], 1970. Russian translation by В.А. Исаковских [V.A. Isakovskikx] under edition М.М. Постников [M.M. Postnikov].

[28] K. Scharnhorst and J.-W. van Holten: Nonlinear Bogolyubov-Valatin transformations: 2 modes. Annals of Physics (New York), 326(2011)2868-2933 [arXiv:1002.2737v3, NIKHEF preprint NIKHEF/2010-005] (DOI: 10.1016/j.aop.2011.05.001

[29] É. Cartan: Leçons sur la Théorie des Spineurs, 2 Vols.. Vol. I: Les Spineurs de l'Espace a Trois Dimensions. Actual. Sci. Ind., Vol. 643, Exposés Géom., Vol. 9. Vol. II: Les Spineurs de l'Espace a n > 3 dimensions. Les Spineurs en Géométrie Riemanienne. Actual. Sci. Ind., Vol. 701, Exposés Géom., Vol. 11. Hermann, Paris, 1938. English translation: The Theory of Spinors. Hermann, Paris, 1966. Reprinted: Dover Publications, Inc., New York, 1981. Russian translation: Э. Картан [È Kartan]: Теория спиноров [Teoriya spinorov]. Платон [Platon], Москва [Moskva], 1997. Russian translation by П. А. Широков [P.A. Shirokov].

[30] W.V.D. Hodge, D. Pedoe: Methods of Algebraic Geometry, Vol. 2. Cambridge University Press, Cambridge, 1952. Russian edition: В.Д. Ходж [V. D. Khodzh], Д. Пидо [D. Pido]: Методы алгебраической геометрии [Metody algebraicheskoĭ geometrii]. Т. 2 [Tom 2]. ИЛ [IL], Москва [Moskva], 1954. Russian translation by А.И. Узков [A.I. Uzkov].

[31] A.A. Albert: Quadratic Forms Permitting Composition. Ann. of Math. 43(1942)161-177.

[32] Хуа Ло-гэн [Khua Lo-gèn], Б.А. Розенфельд [B.A. Rozenfel'd]: Геометрия прямоугольных матриц и ее применение к вещественной проективной и неевклидовой геометрии. [Geometriya pryamougol'nykh matrits i ee primenenie k veshchestvennoĭ proektivnoĭ i neevklidovoĭ geometrii]. Изв. Высш. Учебн. Завед., Матем. [Izv. Vyssh. Uchebn. Zaved., Matem.] (1957) No. 1, 233-247 (Mathnet URL: `http://mi.mathnet.ru/ivm3038`). English translation: Hua Loo-geng (Hua Loo-keng), B.A. Rozenfel'd: The geometry of rectangular matrices and its application to real-projective and non-euclidean geometry. Chin. Math. 8(1966)726-737.

[33] К. В. Андреев [K.V. Andreev]: О спинорном формализме при четной размерности базового пространства [O spinornom formalizme pri chetnoĭ razmernosti bazovogo prostranstva]. ВИНИТИ-298-В-11 [VINITI-298-V-11], июнь [Jun] 2011, 138 с. [138pp. Paper deponed on Jun 16, 2011 at VINITI (Moscow), ref. №298-V 11], [in Russian: On the spinor formalism for the base space of even dimension].

МИНИСТЕРСТВО ОБРАЗОВАНИЯ И НАУКИ
РОССИЙСКОЙ ФЕДЕРАЦИИ
ГОСУДАРСТВЕННОЕ ОБРАЗОВАТЕЛЬНОЕ УЧРЕЖДЕНИЕ
ВЫСШЕГО ПРОФЕССИОНАЛЬНОГО ОБРАЗОВАНИЯ
БАШКИРСКИЙ ГОСУДАРСТВЕННЫЙ УНИВЕРСИТЕТ

УДК 514.744.2

К.В. Андреев

О спинорном формализме при четной размерности базового пространства.

УФА-2011

II

Русская редакция

Содержание

1 Введение **128(5)**

2 Алгебра Клиффорда **129(6)**

3 Комплексная и вещественные реализации **133(10)**

4 Вещественные вложения. Инволюции **138(15)**

5 Элементарные преобразования ортогональной группы **140(17)**

6 Уравнение Клиффорда. Двулистное накрытие **150(27)**

7 Кватернионы **162(39)**

8 Частные решения уравнения Клиффорда **166(43)**

9 О структурных константах алгебр без деления. Седенионы **178(54)**

10 Инфинитизимальные преобразования **183(59)**

11 Комплексная и вещественная связности **184(60)**

12 О классификации тензоров с симметриями тензора кривизны. Спиноры кривизны **193(69)**

13 Твисторное уравнение **197(73)**

14 Спинорный формализм для n=6 и n=8 **200(76)**

- 14.1 Основные изоморфизмы . . . . 200(76)
- 14.2 О классификации тензоров с симметриями тензора кривизны для n=6. Спиноры кривизны . . . . 202(78)
- 14.3 Геометрическое представление твистора в $\mathbb{R}^6_{(2,4)}$ . . . . 205(81)

14.4 О структурных константах алгебры октонионов. Спинорный формализм для n = 8 . . . 211(87)

14.5 Геометрия индуктивного перехода от n=6 к n=8 . . . 213(89)

14.5.1 Нуль-пары Розенфельда . . . 213(89)

14.5.2 Построение квадрик $\mathbb{C}Q_6$ и $\mathbb{C}\tilde{Q}_6$ . . . 214(90)

14.5.3 Соответствие $\mathbb{C}Q_6 \longmapsto \mathbb{C}\tilde{Q}_6$ . . . 215(91)

14.5.4 Геометрия перехода к связующим операторам ${\eta_\Lambda}^{KL}$ для n=8 . . . 218(94)

14.5.5 Соответствие $\mathbb{C}\tilde{Q}_6 \longmapsto \mathbb{C}Q_6$ . . . 223(99)

14.5.6 О майорановских спинорах. . . . 225(101)

14.5.7 Теорема о двух квадриках. . . . 227(103)

14.6 Явное построение аналогов операторов Ли при n=4. . . . 229(105)

15 Благодарности **234(110)**

16 Приложение **235(111)**

16.1 Доказательство формул о метрических тензорах и инволюции . . . 235(111)

16.2 Доказательство формул о связующих операторах . . . 237(113)

16.3 Доказательство формул об инфинитезимальных преобразованиях . . . 239(115)

16.4 Доказательство тождества о скобке Ли . . . 240(116)

16.5 Доказательство формул о тензоре кривизны . . . 241(117)

16.6 Доказательство вспомогательных тождеств для теоремы о спинорных аналогах тензора кривизны . . . 241(117)

16.7 Доказательство формул о твисторном уравнении. . . . 244(120)

16.8 Доказательство тождеств для n=6 . . . 246(122)

16.8.1 Доказательство основных тождеств для n=6 . . . 246(122)

16.8.2 Доказательство тождеств о 4-векторе . . . 247(123)

16.8.3 Доказательство формул о 6-векторе . . . 247(123)

16.8.4 Доказательство тождества Бианки . . . 249(125)

16.8.5 Доказательство тождеств, касающихся тензора Вейля. . . . 249(125)

16.8.6 Доказательство формул, связанных с метрикой, индуцированной в сечении конуса K_6 . . . 250(126)

16.8.7 Доказательство формул о первом инварианте . . . 251(127)

16.8.8 Доказательство формул о втором инварианте . . . . 253(129)
16.8.9 Доказательство тождества альтернативности для алгебры октав . 255(131)
16.8.10 Доказательство центрального тождества Муфанг . . . . 256(132)
16.8.11 Доказательство тождеств для оператора A для n=8 . . . . 257(133)

17 Добавление **262**

Список таблиц

1 Вид матрицы тензора S для соответствующих действительных вложений 2-мерных пространств в комплексное для $r^i = (T, Z)$. . . . . 176(52)
2 Вид матрицы тензора S для соответствующих действительных вложений 4-мерных пространств в комплексное для $r^i = (T, X, Y, Z)$. . . . . 176(52)
3 Вид матрицы тензора S для соответствующих действительных вложений 6-мерных пространств в комплексное для $r^i = (T, X, V, W, Y, Z)$. . . . . . . 177(53)

Список рисунков

1 Соответствие $\forall CP_2 \subset CP_3 \leftrightarrow R$ . . . . 217(93)
2 Соответствие $CP_3 \supset CP_1 \leftrightarrow CP_1 \subset K_6$ . . . . 217(93)

1 Введение

В статье[3] рассматриваются ответы на следующие вопросы при произвольном четном $n \geq 4$:

1. Что такое алгебра Клиффорда и как ее построить?
2. Что такое вещественная и комплексная реализация?
3. Что такое инволюция и как она помогает при переходе к вещественным вложениям?
4. Как построить вещественную и комплексную реализацию образующих (связующих операторов) алгебры Клиффорда?
5. Как инволюция действует на образующие (связующие операторы) алгебры Клиффорда?
6. Как привести комплексную ортогональную матрицу к блочно-диагональному виду?
7. Как построить основные изоморфизмы (в т.ч. двулистные накрытия) и иные соотношения в явном виде с помощью связующих операторов?
8. Как построить частные решения уравнения Клиффорда для связующих операторов при четном $n \geq 4$?
9. Как по связующим операторам построить структурные константы алгебр (без деления) гиперкомплексных чисел (седенионов) для $n \; mod \; 8 = 0$?
10. Как ввести и согласовать связности в касательном и спинорном расслоении?
11. Как построить аналоги операторов Ли и каковы условия их построения?
12. Как построить спиноры кривизны?
13. Какова связь между твисторным уравнением, деривационным уравнением нормализованного грассманиана и конформным уравнением Киллинга?
14. Почему отличается спинорный формализм при $n \leq 8$ от $n > 8$ и как его построить для малых размерностей?
15. Как построить геометрическое представление спинора для $\mathbb{R}^6_{(2,4)}$?
16. Как построить обобщение принципа тройственности Картана с соответствием Клейна и какова геометрия такого обобщения при $n = 8$?
17. Как построить спинорные аналоги операторов Ли при $n = 4$ в явном виде?

---

[3] Статья приведена с исправлениями и оригинальной нумерацией страниц.

2 Алгебра Клиффорда

В этом параграфе рассказывается о том, как построить вещественную 2n-мерную алгебру Клиффорда с произвольным функционалом G согласно [18, лекция 13, с. 258-299]. В случае, когда функционал G в подходящем базисе имеет диагональный вид с одинаковым количеством $\ll + \gg$ *и* $\ll - \gg$ *на главной диагонали, образующие такой алгебры Клиффорда порождают образующие комплексной n-мерной алгебры Клиффорда, которая изоморфна алгебре комплексных матриц* $\mathbb{C}(2^{\frac{n}{2}})$. *Вывод всех результатов этого параграфа сделан на основе [18].*

Пусть $\mathcal{V}$ - векторное пространство над $\mathbb{R}$. Тогда $\mathcal{V}$ будет модулем над $\mathbb{R}$: 1). a(x+y)=ax+ay, 2). (a+b)x=ax+bx, 3). (ab)x=a(bx), 4). 1$\cdot$ x=x, где x,y $\in$ $\mathcal{V}$, a,b $\in$ $\mathbb{R}$. Рассмотрим в алгебре $T_0(\mathcal{V}) = {T_0}^0(\mathcal{V}) \oplus ... \oplus {T_0}^q(\mathcal{V}) \oplus ...$ идеал

$$J(G) := \{T \otimes (x \otimes x - \frac{1}{2}G(x));\ T \in T_0(\mathcal{V}), x \in {T_0}^1(\mathcal{V})\} \tag{2.1}$$

и определим алгебру Клиффорда как $CL(G^{2n}) := T_0(\mathcal{V})/J(G)$. Построим представление такой алгебры в алгебру многочленов

$$\begin{aligned} &\alpha : \mathcal{V} \longmapsto \mathcal{A}, &&\tilde{\alpha} : T_0(\mathcal{V}) \longmapsto \mathcal{A}, \\ &\alpha(\mathcal{V}) = B^{\tilde{\Lambda}} x_{\tilde{\Lambda}}, &&\tilde{\alpha}(T_0(\mathcal{V})) = A + B^{\tilde{\Lambda}} x_{\tilde{\Lambda}} + C^{\tilde{\Lambda}\tilde{\Psi}} x_{\tilde{\Lambda}} x_{\tilde{\Psi}} + ... \ . \end{aligned} \tag{2.2}$$

Здесь и далее $\tilde{\Lambda}$, $\tilde{\Psi}$, $... = \overline{1, 2n}$. Пусть теперь выполнено $\hat{\alpha}(J(G)) = 0$ для некоторого отображения $\hat{\alpha} : CL(G) \longmapsto C\mathcal{A}$. Это будет означать, что

$$x_{\tilde{\Lambda}} x_{\tilde{\Psi}} + x_{\tilde{\Psi}} x_{\tilde{\Lambda}} = G(x_{\tilde{\Lambda}}, x_{\tilde{\Psi}}). \tag{2.3}$$

Поэтому форма $G(x_{\tilde{\Lambda}}, x_{\tilde{\Psi}})$ в подходящем базисе имеет диагональный вид. Пусть для такого базиса выполняются соотношения

$$x_{\tilde{\Lambda}} x_{\tilde{\Psi}} + x_{\tilde{\Psi}} x_{\tilde{\Lambda}} = 0, \ \ (\tilde{\Lambda} \neq \tilde{\Psi}); \ \ {x_{\tilde{\Lambda}}}^2 = \pm 1, \tag{2.4}$$

что определит отображение

$$\hat{\alpha}(Cl(G^{2n})) = A + B^{\tilde{\Lambda}} x_{\tilde{\Lambda}} + C^{[\tilde{\Lambda}\tilde{\Psi}]} x_{\tilde{\Lambda}} x_{\tilde{\Psi}} + ... \tag{2.5}$$

В свою очередь, это означает, что алгебра Клиффорда конечна и $dim\, CL(G^{2n}) = 2^{2n}$, $dim\, \mathcal{V} = 2n$. Построим теперь комплексную алгебру Клиффорда $CL(g^n)$. Для этого рассмотрим вещественную алгебру Клиффорда $CL(G^{2n}_{(n,n)})$. Будем считать, что $(x_{2\Lambda})^2 = 1$, $(x_{2\Lambda-1})^2 = -1$. Зададим

$$\sqrt{2} z_\Lambda = x_{2\Lambda} + i x_{2\Lambda-1}, \ \ \sqrt{2} \bar{z}_\Lambda = x_{2\Lambda} - i x_{2\Lambda-1}. \tag{2.6}$$

Здесь и далее $\Lambda, \quad \Psi, \quad ... = \overline{1,n}$. Тогда

$$\begin{aligned}
2(z_\Lambda z_\Psi + z_\Psi z_\Lambda) &= (x_{2\Lambda} + i x_{2\Lambda-1})(x_{2\Psi} + i x_{2\Psi-1}) + (x_{2\Psi} + i x_{2\Psi-1})(x_{2\Lambda} + i x_{2\Lambda-1}) = \\
&= x_{2\Lambda}x_{2\Psi} - x_{2\Lambda-1}x_{2\Psi-1} + i(x_{2\Lambda-1}x_{2\Psi} + x_{2\Lambda}x_{2\Psi-1}) + \\
&+ x_{2\Psi}x_{2\Lambda} - x_{2\Psi-1}x_{2\Lambda-1} + i(x_{2\Psi-1}x_{2\Lambda} + x_{2\Psi}x_{2\Lambda-1}) = \\
&= \begin{cases} 4, & \Lambda = \Psi, \\ 0, & \Lambda \neq \Psi, \end{cases} \\
2(z_\Lambda \bar{z}_\Psi + \bar{z}_\Psi z_\Lambda) &= (x_{2\Lambda} + i x_{2\Lambda-1})(x_{2\Psi} - i x_{2\Psi-1}) + (x_{2\Psi} - i x_{2\Psi-1})(x_{2\Lambda} + i x_{2\Lambda-1}) = \\
&= x_{2\Lambda}x_{2\Psi} + x_{2\Lambda-1}x_{2\Psi-1} - i(-x_{2\Lambda-1}x_{2\Psi} + x_{2\Lambda}x_{2\Psi-1}) + \\
&+ x_{2\Psi}x_{2\Lambda} + x_{2\Psi-1}x_{2\Lambda-1} - i(x_{2\Psi-1}x_{2\Lambda} - x_{2\Psi}x_{2\Lambda-1}) = 0,
\end{aligned} \tag{2.7}$$

$$z_\Lambda z_\Psi + z_\Psi z_\Lambda = g(z_\Lambda, z_\Psi). \tag{2.8}$$

Поэтому факторизация комплексной алгебры $T_0(\mathcal{V}^{\mathbb{C}}) = T_0(\mathcal{V}^{\mathbb{C}})(z)$ по идеалу $J(g) = \{T \otimes (z \otimes z - \frac{1}{2}g(z)); \quad T \in T_0(\mathcal{V}^{\mathbb{C}}), \quad z \in {T_0}^1(\mathcal{V}^{\mathbb{C}})\}$ определит комплексную алгебру $CL(g^n) \cong T_0(\mathcal{V}^{\mathbb{C}})/J(g)$.

Теорема 2.1. *Для алгебр Клиффорда существует разложение*

$$\begin{gathered}
CL(G^{(2n+2)}_{(n+1,n+1)}) = CL(G^{2n}_{(n,n)}) \otimes CL(G^2_{(1,1)}), \\
CL(G^{(2n+2)}_{(2n+2,0)}) = CL(G^{2n}_{(2n,0)}) \otimes CL(G^2_{(2,0)}), \quad CL(g^n) = CL(g^{n-2}) \otimes CL(g^2)
\end{gathered} \tag{2.9}$$

с умножением $a \cdot b = (\tilde{a}, \quad \tilde{\tilde{a}}) \cdot (\tilde{b}, \quad \tilde{\tilde{b}}) = (\tilde{a}\tilde{b}, \quad \tilde{\tilde{a}}\tilde{\tilde{b}})$.

Доказательство. Пусть $x_{\tilde{\Lambda}}$ - базис $\mathcal{V}^{2n}$, а x_{2n+1}, x_{2n+2} - базис в $\mathcal{V}^2$

$$\begin{gathered}
x_{\tilde{\Lambda}} x_{\tilde{\Psi}} + x_{\tilde{\Psi}} x_{\tilde{\Lambda}} = 0, \ (\tilde{\Lambda} \neq \tilde{\Psi}); \quad (x_{2\Lambda})^2 = 1; \ (x_{2\Lambda-1})^2 = -1, \\
x_{2n+1}x_{2n+2} + x_{2n+2}x_{2n+1} = 0; \quad (x_{2n+2})^2 = -1; \ (x_{2n+1})^2 = 1.
\end{gathered} \tag{2.10}$$

Это даст возможность определить базис в $\mathcal{V}^{2n+2}$ как

$$X_{\tilde{\Lambda}} = x_{\tilde{\Lambda}} x_{2n+1} x_{2n+2}, \quad X_{2n+1} = x_{2n+2}, \quad X_{2n+2} = x_{2n+1} \tag{2.11}$$

с условиями

$$\begin{gathered}
X_{\tilde{\Lambda}} X_{\tilde{\Psi}} + X_{\tilde{\Psi}} X_{\tilde{\Lambda}} = 0, \ (\tilde{\Lambda} \neq \tilde{\Psi}), \quad (X_{2\Lambda})^2 = 1, \quad (X_{2\Lambda-1})^2 = -1, \\
\text{в этом уравнении } \Lambda = \overline{1, n+1}, \quad \tilde{\Lambda} = \overline{1, 2n+2}.
\end{gathered} \tag{2.12}$$

Тогда для любого $a \in CL(G^{2n+2}_{(n+1,n+1)})$ выполнено разложение

$$\begin{gathered}
a = a_1 X_{2n+2} X_{2n+1} + a_2 X_{2n+2} + a_3 X_{2n+1} + a_4 = \\
= a_1 \otimes x_{2n+1} x_{2n+2} + a_2 \otimes x_{2n+1} + a_3 \otimes x_{2n+2} + a_4 \otimes 1.
\end{gathered} \tag{2.13}$$

Проведем замену базиса

$$\begin{gathered}y_{2n+1} = (x_{2n+1} + x_{2n+2})/2, \quad y_{2n+2} = (x_{2n+1} - x_{2n+2})/2, \\ (y_{2n+1})^2 = 0, \quad (y_{2n+2})^2 = 0, \quad 2y_{2n+1}y_{2n+2} = 1 - x_{2n+1}x_{2n+2}.\end{gathered} \tag{2.14}$$

Тогда разложение (2.13) примет вид

$$a = (-a_1 + a_4)y_{2n+1}y_{2n+2} + (a_2 + a_3)y_{2n+1} + (a_2 - a_3)y_{2n+2} + (a_1 + a_4)y_{2n+2}y_{2n+1}. \tag{2.15}$$

Следовательно, любой элемент $a \in CL(G^2_{(1,1)})$ разлагается как

$$\begin{gathered}a = a_1x_{2n+1}x_{2n+2} + a_2x_{2n+1} + a_3x_{2n+2} + a_4 = \\ = \begin{pmatrix} y_{2n+1}y_{2n+2} & y_{2n+2} \end{pmatrix} \begin{pmatrix} -a_1 + a_4 & a_2 + a_3 \\ a_2 - a_3 & a_1 + a_4 \end{pmatrix} \begin{pmatrix} y_{2n+1}y_{2n+2} \\ y_{2n+1} \end{pmatrix},\end{gathered} \tag{2.16}$$

а для любых двух элементов $a, b \in CL(G^{2n+2}_{(n+1,n+1)})$ выполнено

$$\begin{aligned} c := a \cdot b \;=\; & (a_1 \otimes x_{2n+1}x_{2n+2} + a_2 \otimes x_{2n+1} + a_3 \otimes x_{2n+2} + a_4 \otimes 1)\cdot \\ & \cdot(b_1 \otimes x_{2n+1}x_{2n+2} + b_2 \otimes x_{2n+1} + b_3 \otimes x_{2n+2} + b_4 \otimes 1) = \\ = & \underbrace{(a_2b_3 - a_3b_2 + a_1b_4 + a_4b_1)}_{:=c_1} \otimes x_{2n+1}x_{2n+2} + \underbrace{(a_2b_4 + a_4b_2 - a_1b_3 + a_3b_1)}_{:=c_2} \otimes x_{2n+1} + \\ + & \underbrace{(a_3b_4 + a_4b_3 - a_1b_2 + a_2b_1)}_{:=c_3} \otimes x_{2n+2} + \underbrace{(a_1b_1 + a_4b_4 + a_2b_2 - a_3b_3)}_{:=c_4} \end{aligned} \tag{2.17}$$

В то же время для любых двух элементов $a, b \in CL(G^2_{(1,1)})$ верно тождество

$$\begin{gathered} c := a \cdot b = \begin{pmatrix} -a_1 + a_4 & a_2 + a_3 \\ a_2 - a_3 & a_1 + a_4 \end{pmatrix} \cdot \begin{pmatrix} -b_1 + b_4 & b_2 + b_3 \\ b_2 - b_3 & b_1 + b_4 \end{pmatrix} = \\ = \begin{pmatrix} a_1b_1 - a_1b_4 - a_4b_1 + a_4b_4 + a_2b_2 + a_3b_2 - a_2b_3 - a_3b_3 & -a_1b_2 + a_4b_2 - a_1b_3 + a_4b_3 + a_2b_1 + a_3b_1 + a_2b_4 + a_3b_4 \\ -a_2b_1 + a_3b_1 + a_2b_4 - a_3b_4 + a_1b_2 + a_4b_2 - a_1b_3 - a_4b_3 & a_2b_2 - a_3b_2 + a_2b_3 - a_3b_3 + a_1b_1 + a_4b_1 + a_1b_4 + a_4b_4 \end{pmatrix} = \\ = \begin{pmatrix} -c_1 + c_4 & c_2 + c_3 \\ c_2 - c_3 & c_1 + c_4 \end{pmatrix}. \end{gathered} \tag{2.18}$$

Что даст возможность поставить в соответствие образующим алгебры Клиффорда $CL(G^2_{(1,1)})$ их матричное представление

$$x_{2n+1} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad x_{2n+2} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \tag{2.19}$$

Это означает, что для любых двух элементов $a, b \in CL(G^{2n+2}_{(n+1,n+1)})$ в аналогичном матричном представлении $c_1, c_2, c_3, c_4 \in CL(G^{2n}_{(n,n)})$. Аналогично доказываются и два других представления. □

Теорема 2.2. *Комплексная алгебра Клиффорда при четном n изоморфна алгебре комплексных матриц вида*

$$CL(g) \cong \mathbb{C}(2^{n/2}). \tag{2.20}$$

Доказательство. Доопределим, согласуя с доказательством теоремы 2.1,

$$\begin{gathered}
\sqrt{2}z_{n+1} = x_{2n+2} + ix_{2n+1}, \quad \sqrt{2}z_{n+2} = x_{2n+4} + ix_{2n+3}, \\
z_{n+1}z_{n+1} = 1, \quad z_{n+1}\bar{z}_{n+1} = ix_{2n+1}x_{2n+2} = -\bar{z}_{n+1}z_{n+1}, \\
z_{n+2}z_{n+2} = 1, \quad z_{n+2}\bar{z}_{n+2} = ix_{2n+3}x_{2n+4} = -\bar{z}_{n+2}z_{n+2}, \\
z_{n+1}z_{n+2} + z_{n+2}z_{n+1} = 0.
\end{gathered} \tag{2.21}$$

Поставим в соответствие образующим $CL(G^4_{(2,2)})$ их матричное представление

$$\begin{gathered}
x_{2n+2} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad x_{2n+1} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \\
x_{2n+4} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \quad x_{2n+3} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}.
\end{gathered} \tag{2.22}$$

Тогда для образующих $CL(g^2)$ в базисе, подобном базису примера 3.2[4], получим

$$\sqrt{2}z_{n+1} = \begin{pmatrix} 0 & 1+i \\ 1-i & 0 \end{pmatrix}, \quad \sqrt{2}z_{n+2} = \begin{pmatrix} 0 & 1-i \\ 1+i & 0 \end{pmatrix}, \tag{2.23}$$

что и докажет нашу теорему. В общем виде данное соответствие будет построено ниже. □

Таким образом вещественная алгебра Клиффорда $CL(G^{2n}_{(n,n)})$ изоморфна вещественной алгебре матриц размерности n

$$CL(G^{2n}_{(n,n)}) \cong \mathbb{R}(2^n). \tag{2.24}$$

Поэтому существует отображение

$$\gamma : \quad \mathbb{R}(2^n) \longmapsto (\mathcal{V}^{\mathbb{R}})^{2n}. \tag{2.25}$$

---

[4] С точностью до множителя $x_{\tilde{\Lambda}} \longmapsto \frac{1}{\sqrt{2}}x_{\tilde{\Lambda}}$, $z_{\Lambda} \longmapsto \frac{1}{\sqrt{2}}z_{\Lambda}$.

На основании (2.3) операторы $\gamma_{\tilde{\Lambda}}$ должны необходимо удовлетворять уравнению

$$\gamma_{\tilde{\Lambda}}\gamma_{\tilde{\Psi}} + \gamma_{\tilde{\Psi}}\gamma_{\tilde{\Lambda}} = G_{\tilde{\Lambda}\tilde{\Psi}}, \tag{2.26}$$

которое и называется уравнением Клиффорда. В то же время комплексная алгебра Клиффорда представляется матричной алгеброй вида $\mathbb{C}(2^{n/2})$. Умножение в комплексной алгебре Клиффорда делегирует матричное умножение в $\mathbb{C}(2^{n/2})$. Поэтому можно построить отображение

$$\gamma: \quad \mathbb{C}(2^{n/2}) \longmapsto (\mathcal{V}^{\mathbb{C}})^{n}. \tag{2.27}$$

Из (2.8) следует, что операторы γ_{Λ} должны необходимо удовлетворять комплексному уравнению Клиффорда

$$\gamma_{\Lambda}\gamma_{\Psi} + \gamma_{\Psi}\gamma_{\Lambda} = g_{\Lambda\Psi}. \tag{2.28}$$

3 Комплексная и вещественные реализации

В этом параграфе рассказывается о том, как по вещественной реализации 2n-мерного псевдориманова многообразия со слоями касательного расслоения, изоморфными $\mathbb{R}^{2n}_{(n,n)}$*, локально построить его n-мерное комплексное представление. Для этого строится соответствующая комплексная репараметризация атласа в окрестности некоторой точки, что индуцирует в касательном расслоении операторы Нейфельда [15], связанные с аффинором Нордена [17]. Соответственно, в спинорном расслоении связующими операторами* $\gamma_{\tilde{\Lambda}}$ *индуцируются с точностью до знака аналоги операторов Нейфельда. Это даст возможность перейти к комплексной реализации связующих операторов* γ_{Λ}*. Вывод всех результатов этого параграфа сделан на основе [15], [17].*

Под комплексным аналитическим римановым пространством $\mathbb{C}V_n$ в дальнейшем будем понимать аналитическое комплексное многообразие, снабженное аналитической квадратичной метрикой, т.е. метрикой, определенной с помощью симметрического невырожденного тензора $g_{\Lambda\Psi}$ (здесь $\Lambda,\ \ \Psi,\ \ ... = \overline{1,n}$; $\tilde{\Lambda},\ \ \tilde{\Psi},\ \ ... = \overline{1,2n}$), координаты которого - аналитические функции координат точки. Этому тензору соответствует комплексная риманова связность без кручения, коэффициенты которой определяются символами Кристофеля и поэтому являются аналитическими функциями. Касательное расслоение этого многообразия $\tau^{\mathbb{C}}(\mathbb{C}V_n)$ имеет слои $\tau^{\mathbb{C}}_{x} \cong \mathbb{C}\mathbb{R}^{n}$, то есть слои, изоморфные n-мерному комплексному евклидовому пространству, метрика которого определяется значением метрического тензора в данной точке. Вещественная реализация V_{2n} нашего $\mathbb{C}V_n$ имеет касательное расслоение $\tau^{\mathbb{R}}(V_{2n})$ со слоями, изоморфными $\mathbb{R}^{2n}_{(n,n)}$. Пусть на V_{2n} задан атлас $(U; x^{\tilde{\Lambda}})$. Рассмотрим репараметризацию этого атласа $(U; w^\Lambda)$ такую, что $w^\Lambda = \frac{1}{\sqrt{2}}(u^\Lambda(x^{\tilde\Lambda}) + iv^\Lambda(x^{\tilde\Lambda}))$, которая локально разрешима как $x^{\tilde\Lambda} = x^{\tilde\Lambda}(u^\Lambda, v^\Lambda)$. Положим

$$m^\Lambda{}_{\tilde\Lambda} := \frac{1}{\sqrt{2}}(\frac{\partial u^\Lambda}{\partial x^{\tilde\Lambda}} + i\frac{\partial v^\Lambda}{\partial x^{\tilde\Lambda}}) =: \frac{\partial w^\Lambda}{\partial x^{\tilde\Lambda}}, \quad m_\Lambda{}^{\tilde\Lambda} := \frac{1}{\sqrt{2}}(\frac{\partial x^{\tilde\Lambda}}{\partial u^\Lambda} - i\frac{\partial x^{\tilde\Lambda}}{\partial v^\Lambda}) =: \frac{\partial x^{\tilde\Lambda}}{\partial w^\Lambda}. \tag{3.1}$$

Тогда

$$\Delta_{\tilde\Lambda}{}^{\tilde\Psi} := m^\Lambda{}_{\tilde\Lambda} m_\Lambda{}^{\tilde\Psi} = \frac{1}{2}((\underbrace{\frac{\partial u^\Lambda}{\partial x^{\tilde\Lambda}}\ \frac{\partial x^{\tilde\Psi}}{\partial u^\Lambda} + \frac{\partial v^\Lambda}{\partial x^{\tilde\Lambda}}\ \frac{\partial x^{\tilde\Psi}}{\partial v^\Lambda}}_{=\delta_{\tilde\Lambda}{}^{\tilde\Psi}}) + i(\underbrace{\frac{\partial v^\Lambda}{\partial x^{\tilde\Lambda}}\ \frac{\partial x^{\tilde\Psi}}{\partial u^\Lambda} - \frac{\partial u^\Lambda}{\partial x^{\tilde\Lambda}}\ \frac{\partial x^{\tilde\Psi}}{\partial v^\Lambda}}_{=:f_{\tilde\Lambda}{}^{\tilde\Psi}})), \tag{3.2}$$

что определит комплексную структуру f на V_{2n}, где $\Delta_{\tilde\Lambda}{}^{\tilde\Psi}$ - аффинор Нордена. Определим $\bar{m}^{\Psi'}{}_{\tilde\Psi} := \overline{m^\Psi{}_{\tilde\Psi}}$ и положим

$$0 =: m^\Lambda{}_{\tilde\Lambda} \bar{m}_{\Psi'}{}^{\tilde\Lambda} = \frac{1}{2}((\underbrace{\frac{\partial u^\Lambda}{\partial x^{\tilde\Lambda}}\ \frac{\partial x^{\tilde\Lambda}}{\partial u^{\Psi'}} - \frac{\partial v^\Lambda}{\partial x^{\tilde\Lambda}}\ \frac{\partial x^{\tilde\Lambda}}{\partial v^{\Psi'}}}_{\equiv 0}) + i\ (\frac{\partial u^\Lambda}{\partial x^{\tilde\Lambda}}\ \frac{\partial x^{\tilde\Lambda}}{\partial v^{\Psi'}} + \frac{\partial v^\Lambda}{\partial x^{\tilde\Lambda}}\ \frac{\partial x^{\tilde\Lambda}}{\partial u^{\Psi'}})), \tag{3.3}$$

$$\delta_\Psi{}^\Lambda =: m^\Lambda{}_{\tilde\Lambda} m_\Psi{}^{\tilde\Lambda} = \frac{1}{2}((\underbrace{\frac{\partial u^\Lambda}{\partial x^{\tilde\Lambda}}\ \frac{\partial x^{\tilde\Lambda}}{\partial u^\Psi} + \frac{\partial v^\Lambda}{\partial x^{\tilde\Lambda}}\ \frac{\partial x^{\tilde\Lambda}}{\partial v^\Psi}}_{=2\delta_\Psi{}^\Lambda}) - i\ (\frac{\partial u^\Lambda}{\partial x^{\tilde\Lambda}}\ \frac{\partial x^{\tilde\Lambda}}{\partial v^\Psi} - \frac{\partial v^\Lambda}{\partial x^{\tilde\Lambda}}\ \frac{\partial x^{\tilde\Lambda}}{\partial u^\Psi})). \tag{3.4}$$

Откуда $\frac{\partial u^\Lambda}{\partial x^{\tilde\Lambda}}\ \frac{\partial x^{\tilde\Lambda}}{\partial v^\Psi} \equiv 0$ (поскольку $v^{\Psi'} = \delta_\Psi{}^{\Psi'} v^\Psi$ и $u^{\Psi'} = \delta_\Psi{}^{\Psi'} u^\Psi$, а тензор $\delta_\Psi{}^{\Psi'}$ имеет единичную матрицу). Следовательно

$$\frac{\partial u^\Lambda}{\partial x^{\tilde\Lambda}} f_{\tilde\Psi}{}^{\tilde\Lambda} = \frac{\partial u^\Lambda}{\partial x^{\tilde\Lambda}}\ (\frac{\partial v^\Psi}{\partial x^{\tilde\Psi}}\ \frac{\partial x^{\tilde\Lambda}}{\partial u^\Psi} - \frac{\partial u^\Psi}{\partial x^{\tilde\Psi}}\ \frac{\partial x^{\tilde\Lambda}}{\partial v^\Psi}) = \frac{\partial v^\Lambda}{\partial x^{\tilde\Psi}} \tag{3.5}$$

есть условия Коши-Римана, если в данном базисе комплексная структура имеет канонический вид

$$f_{\tilde\Lambda}{}^{\tilde\Psi} = \begin{pmatrix} 0 & E \\ -E & 0 \end{pmatrix}, \tag{3.6}$$

где E - единичная матрица размера $n \times n$. Легко проверить, что $f^2 = -E$. Если теперь потребовать, чтобы карты $\mathbb{C}V_n$ были согласованы голоморфными преобразованиями, то таким образом можно отождествить два многообразия: $\mathbb{C}V_n$ и V_{2n}. При этом можно построить изоморфизм между слоями касательных расслоений: $\tau^{\mathbb{C}}(\mathbb{C}V_n)$ и $\tau^{\mathbb{C}}(V_{2n})$, так же как и между слоями их вещественных реализаций: $\tau^{\mathbb{R}}(\mathbb{C}V_n)$ и $\tau^{\mathbb{R}}(V_{2n})$.

Пусть в том же базисе, в котором комплексная структура имеет канонический вид, наша метрика такая

$$G_{\tilde\Lambda\tilde\Psi} = \begin{pmatrix} E & 0 \\ 0 & -E \end{pmatrix}, \tag{3.7}$$

где E - единичная матрица размера $n \times n$. За соответствие $V_{2n} \longleftrightarrow \mathbb{C}V_n$ отвечают специальные операторы $m_\Lambda{}^{\tilde\Lambda}$, $m^\Lambda{}_{\tilde\Lambda}$, которые по определению должны удовлетворять следующей системе уравнений

$$\begin{cases} m_{\Lambda}{}^{\tilde{\Lambda}} m^{\Psi}{}_{\tilde{\Lambda}} = \delta_{\Lambda}{}^{\Psi}, \\ \\ m_{\Lambda}{}^{\tilde{\Lambda}} \bar{m}^{\Psi'}{}_{\tilde{\Lambda}} = 0, \\ \\ m_{\Lambda}{}^{\tilde{\Lambda}} m^{\Lambda}{}_{\tilde{\Psi}} = \frac{1}{2}(\delta_{\tilde{\Psi}}{}^{\tilde{\Lambda}} + i f_{\tilde{\Psi}}{}^{\tilde{\Lambda}}) := \triangle_{\tilde{\Psi}}{}^{\tilde{\Lambda}}. \end{cases} \tag{3.8}$$

Тогда из среднего уравнения будет следовать

$$m^{\Psi}{}_{\tilde{\Omega}} = m^{\Lambda}{}_{\tilde{\Omega}}(m_{\Lambda}{}^{\tilde{\Lambda}} m^{\Psi}{}_{\tilde{\Lambda}}) = (m^{\Lambda}{}_{\tilde{\Omega}} m_{\Lambda}{}^{\tilde{\Lambda}}) m^{\Psi}{}_{\tilde{\Lambda}} = \frac{1}{2} m^{\Psi}{}_{\tilde{\Omega}} + \frac{i}{2} f_{\tilde{\Omega}}{}^{\tilde{\Lambda}} m^{\Psi}{}_{\tilde{\Lambda}}. \tag{3.9}$$

Откуда

$$m^{\Psi}{}_{\tilde{\Omega}} = i\ f_{\tilde{\Omega}}{}^{\tilde{\Lambda}} m^{\Psi}{}_{\tilde{\Lambda}} \tag{3.10}$$

опять есть условия Коши-Римана. Поэтому в присутствии метрики $G_{\tilde{\Lambda}\tilde{\Psi}}$ на $\mathbb{R}^{2n}_{(n,n)}$ в комплексной реализации индуцируется метрика

$$\begin{gathered} g_{\Lambda\Psi} := G_{\tilde{\Lambda}\tilde{\Psi}} m_{\Lambda}{}^{\tilde{\Lambda}} m_{\Psi}{}^{\tilde{\Psi}}, \quad \bar{g}_{\Lambda'\Psi'} := G_{\tilde{\Lambda}\tilde{\Psi}} \bar{m}_{\Lambda'}{}^{\tilde{\Lambda}} \bar{m}_{\Psi'}{}^{\tilde{\Psi}}, \\ g_{\Lambda'\Psi} := G_{\tilde{\Lambda}\tilde{\Psi}} \bar{m}_{\Lambda'}{}^{\tilde{\Lambda}} m_{\Psi}{}^{\tilde{\Psi}}, \quad \bar{g}_{\Lambda\Psi'} := G_{\tilde{\Lambda}\tilde{\Psi}} \bar{m}_{\Lambda}{}^{\tilde{\Lambda}} \bar{m}_{\Psi'}{}^{\tilde{\Psi}}. \end{gathered} \tag{3.11}$$

В стандартной комплексной реализации (где вместо пространства $\mathbb{R}^{2n}_{(n,n)}$ берется пространство $\mathbb{R}^{2n}$) подразумевается, что $g_{\Lambda'\Psi'} = \bar{g}_{\Lambda'\Psi'} = 0$, и это приводит к эрмитовости метрики $g_{\Lambda'\Psi}$. Однако, в нашем случае следует воспользоваться условиями

$$g_{\Lambda'\Psi} = \bar{g}_{\Lambda\Psi'} = 0. \tag{3.12}$$

Это приведет к некоторым ограничениям на метрический тензор

$$\begin{gathered} 0 = g_{\Lambda'\Psi} \bar{m}^{\Lambda'}{}_{\tilde{\Omega}} m^{\Psi}{}_{\tilde{\Theta}} = (G_{\tilde{\Lambda}\tilde{\Psi}} \bar{m}_{\Lambda'}{}^{\tilde{\Lambda}} m_{\Psi}{}^{\tilde{\Psi}}) \bar{m}^{\Lambda'}{}_{\tilde{\Omega}} m^{\Psi}{}_{\tilde{\Theta}} = G_{\tilde{\Lambda}\tilde{\Psi}} (\bar{m}_{\Lambda'}{}^{\tilde{\Lambda}} \bar{m}^{\Lambda'}{}_{\tilde{\Omega}})(m_{\Psi}{}^{\tilde{\Psi}} m^{\Psi}{}_{\tilde{\Theta}}) = \\ = G_{\tilde{\Lambda}\tilde{\Psi}} \cdot \tfrac{1}{2}(\delta_{\tilde{\Omega}}{}^{\tilde{\Lambda}} - i f_{\tilde{\Omega}}{}^{\tilde{\Lambda}}) \cdot \tfrac{1}{2}(\delta_{\tilde{\Theta}}{}^{\tilde{\Psi}} + i f_{\tilde{\Theta}}{}^{\tilde{\Psi}}) = \\ = \tfrac{1}{4}(\underbrace{G_{\tilde{\Omega}\tilde{\Theta}} + (f_{\tilde{\Omega}}{}^{\tilde{\Lambda}} f_{\tilde{\Theta}}{}^{\tilde{\Psi}} G_{\tilde{\Lambda}\tilde{\Psi}})}_{=0} + i(\underbrace{(f_{\tilde{\Omega}}{}^{\tilde{\Lambda}} G_{\tilde{\Lambda}\tilde{\Theta}}) - (f_{\tilde{\Theta}}{}^{\tilde{\Psi}} G_{\tilde{\Omega}\tilde{\Psi}}))}_{=0}) = 0. \end{gathered} \tag{3.13}$$

Поэтому наша псевдоевклидова метрика с канонической комплексной структурой удовлетворяет уравнению

$$f_{\tilde{\Omega}}{}^{\tilde{\Lambda}} G_{\tilde{\Lambda}\tilde{\Psi}} f_{\tilde{\Theta}}{}^{\tilde{\Psi}} = -G_{\tilde{\Omega}\tilde{\Theta}}, \tag{3.14}$$

$$\begin{pmatrix} 0 & E \\ -E & 0 \end{pmatrix} \begin{pmatrix} E & 0 \\ 0 & -E \end{pmatrix} \begin{pmatrix} 0 & -E \\ E & 0 \end{pmatrix} = - \begin{pmatrix} E & 0 \\ 0 & -E \end{pmatrix}.$$

Кроме того, будет выполнено

$$G_{\tilde{\Lambda}\tilde{\Psi}} = m^{\Lambda}{}_{\tilde{\Lambda}} m^{\Psi}{}_{\tilde{\Psi}} g_{\Lambda\Psi} + \bar{m}^{\Lambda'}{}_{\tilde{\Lambda}} \bar{m}^{\Psi'}{}_{\tilde{\Psi}} \bar{g}_{\Lambda'\Psi'}. \tag{3.15}$$

Example 3.1. *При n=1 для* $\mathbb{R}^2_{(1,1)}$

$$\begin{gathered} {m_\Lambda}^{\tilde\Lambda} = \tfrac{1}{\sqrt{2}}(1, i), \quad {(m^T)_{\tilde\Lambda}}^{\Lambda} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -i \end{pmatrix}, \quad g_{\Lambda\Psi} = 1, \\ G_{\tilde\Lambda\tilde\Psi} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad {f_{\tilde\Lambda}}^{\tilde\Psi} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \end{gathered} \tag{3.16}$$

Теперь операторы (3.8) можно применить к операторам (2.25). Для этого рассмотрим разложение при n=2 (N=1)

$$ {f_{\tilde\Psi}}^{\tilde\Lambda}\gamma_{\tilde\Lambda} = \tilde{F}\gamma_{\tilde\Psi}\tilde{\tilde{F}}, \quad \tilde{F}^4 = \tilde{\tilde{F}}^4 = E. \tag{3.17}$$

Доказательство этой формулы будет дано в следующем параграфе (6.13). Из (3.17) будет следовать

$$\begin{gathered} {\triangle_{\tilde\Psi}}^{\tilde\Lambda}\gamma_{\tilde\Lambda} = \tfrac{1}{2}({\delta_{\tilde\Psi}}^{\tilde\Lambda} + i{f_{\tilde\Psi}}^{\tilde\Lambda})\gamma_{\tilde\Lambda} = \\ = \underbrace{\frac{1}{2}(\delta + \frac{(1+i)}{2}\tilde{F} + \frac{(1-i)}{2}\tilde{F}^3)}_{:=\tilde\triangle_+}\gamma_{\tilde\Psi}\underbrace{\frac{1}{2}(\delta + \frac{(1+i)}{2}\tilde{\tilde{F}} + \frac{(1-i)}{2}\tilde{\tilde{F}}^3)}_{:=\tilde{\tilde\triangle}_+} + \\ + \underbrace{\frac{1}{2}(\delta - \frac{(1+i)}{2}\tilde{F} - \frac{(1-i)}{2}\tilde{F}^3)}_{:=\tilde\triangle_-}\gamma_{\tilde\Psi}\underbrace{\frac{1}{2}(\delta - \frac{(1+i)}{2}\tilde{\tilde{F}} - \frac{(1-i)}{2}\tilde{\tilde{F}}^3)}_{:=\tilde{\tilde\triangle}_-} = \\ = \tilde\triangle_+\gamma_{\tilde\Psi}\tilde{\tilde\triangle}_+ + \tilde\triangle_-\gamma_{\tilde\Psi}\tilde{\tilde\triangle}_-. \end{gathered} \tag{3.18}$$

Это означает, что можно положить при n=2 (N=1)

$$\left\{\begin{aligned} & {\tilde{m}_{\pm\hat A}}^{\tilde B}{\tilde{m}_{\pm}}^{\hat C}{}_{\tilde B} = {\delta_{\hat A}}^{\hat C}, \\ & {\tilde{m}_{\pm\hat A}}^{\tilde B}{\tilde{m}_{\mp}}^{\hat C}{}_{\tilde B} = 0, \\ & {\tilde{m}_{\pm\hat A}}^{\tilde B}{\tilde{m}_{\pm}}^{\hat A}{}_{\tilde C} = \\ & = \underbrace{\tfrac{1}{2}({\delta_{\tilde C}}^{\tilde B} \pm \tfrac{(1+i)}{2}{\tilde{F}_{\tilde C}}^{\tilde B} \pm \tfrac{(1-i)}{2}{(\tilde{F}^3)_{\tilde C}}^{\tilde B})}_{:={\tilde\triangle_{\pm\tilde C}}^{\tilde B}}, \end{aligned}\right. \qquad \left\{\begin{aligned} & {\tilde{\tilde{m}}_{\pm\hat A}}^{\tilde B}{\tilde{\tilde{m}}_{\pm}}^{\hat C}{}_{\tilde B} = {\delta_{\hat A}}^{\hat C}, \\ & {\tilde{\tilde{m}}_{\pm\hat A}}^{\tilde B}{\tilde{\tilde{m}}_{\mp}}^{\hat C}{}_{\tilde B} = 0, \\ & {\tilde{\tilde{m}}_{\pm\hat A}}^{\tilde B}{\tilde{\tilde{m}}_{\pm}}^{\hat A}{}_{\tilde C} = \\ & = \underbrace{\tfrac{1}{2}({\delta_{\tilde C}}^{\tilde B} \pm \tfrac{(1+i)}{2}{\tilde{\tilde{F}}_{\tilde C}}^{\tilde B} \pm \tfrac{(1-i)}{2}{(\tilde{\tilde{F}}^3)_{\tilde C}}^{\tilde B})}_{:={\tilde{\tilde\triangle}_{\pm\tilde C}}^{\tilde B}}. \end{aligned}\right. \tag{3.19}$$

Здесь $\tilde A, \quad \tilde B, \quad \tilde C, \quad \tilde D, ... = \overline{1,4}$, $\hat A, \hat B, \hat C, \hat D, ... = \overline{1,2}$. Заметим, что можно ввести операцию сопряжения (не комплексного!) такую, что операторы $\tilde{m}_\pm$ ($\tilde{\tilde{m}}_\pm$) и $\tilde{m}_\mp$ ($\tilde{\tilde{m}}_\mp$) относительно такой операции будут сопряжены друг другу. Следовательно, в качестве операторов γ_Λ можно взять либо $\gamma_{+\Lambda}$, либо $\gamma_{-\Lambda}$

$$ {\gamma_{\pm\Lambda\hat A}}^{\hat B} := {m_\Lambda}^{\tilde\Lambda}{\gamma_{\tilde\Lambda\tilde C}}^{\tilde D}{\tilde{m}_{\pm\hat A}}^{\tilde C}{\tilde{\tilde{m}}_{\pm}}^{\hat B}{}_{\tilde D}. \tag{3.20}$$

Example 3.2. *При n=2 для* $\mathbb{R}^4_{(2,2)}$

$$
\begin{gathered}
{m_\Lambda}^{\tilde\Lambda} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & i & 0 & 0 \\ 0 & 0 & 1 & i \end{pmatrix}, \quad {(m^T)_{\tilde\Lambda}}^{\Lambda} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ -i & 0 \\ 0 & 1 \\ 0 & -i \end{pmatrix}, \quad g_{\Lambda\Psi} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \\
G_{\tilde\Lambda\tilde\Psi} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad {f_{\tilde\Lambda}}^{\tilde\Psi} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}.
\end{gathered} \tag{3.21}
$$

$\gamma_{\tilde\Lambda}$ *определены согласно (2.26),* γ_Λ *определены согласно (2.28).*

$$
\begin{gathered}
{m_{\hat A}}^{\tilde B} := {\tilde m_{+\hat A}}^{\tilde B} = {\tilde{\tilde m}_{+\hat A}}^{\tilde B} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & i & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}, \\
{(m^T)_{\tilde B}}^{\hat A} = {(\tilde m^T_+)_{\tilde B}}^{\hat A} = {(\tilde{\tilde m}^T_+)_{\tilde B}}^{\hat A} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ -i & 0 \\ 0 & 1 \\ 0 & 1 \end{pmatrix}, \\
{\tilde m_{-\hat A}}^{\tilde B} = {\tilde{\tilde m}_{-\hat A}}^{\tilde B} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & -i & 0 & 0 \\ 0 & 0 & 1 & -1 \end{pmatrix}, \\
{(\tilde m^T_-)_{\tilde B}}^{\hat A} = {(\tilde{\tilde m}^T_-)_{\tilde B}}^{\hat A} = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ i & 0 \\ 0 & 1 \\ 0 & -1 \end{pmatrix}, \\
{F^{\tilde A}}_{\tilde B} := {\tilde F^{\tilde A}}_{\tilde B} = {\tilde{\tilde F}^{\tilde A}}_{\tilde B} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}.
\end{gathered} \tag{3.22}
$$

Кстати, это еще раз говорит о том, что оператор F *является матричным корнем 4 степени из единицы. Квадрат* F^2 *соответствует в пространстве* $\mathbb{R}^4_{(2,2)}$ *тривиальному преобразованию, умноженному на -1.*

В случае произвольного n алгоритм понижения спинорной размерности следующий.

Алгоритм 3.1. *Комплексная структура представима в виде произведения элементарных преобразований размерности 4×4. Каждое I-ое элементарное преобразование порождает свой оператор* ${(m_I)_\Lambda}^{\tilde\Lambda}$*, который отвечает за переход от вещественной реализации* *подпространства* $(R_I)^4_{(2,2)} \subset \mathbb{R}^{2n}_{(2n,2n)}$ *к комплексной реализации* $(C_I)^2 \subset R^{2n-4}_{(2n-2,2n-2)} \oplus (C_I)^2$ *и, следовательно, понижает размерность пространства на 2. Соответственно, в спинорном пространстве генерируются операторы* $\tilde{\tilde{m}}_{I_\pm}$, $\tilde{m}_{I_\pm}$, *способные понизить размерность спинорного пространства в 2 раза. Очевидно, что требуется* $n/2$ *таких шагов* $(I = \overline{1, \frac{n}{2}})$. *Это понизит спинорную размерность в* $2^{\frac{n}{2}} = 2N$ *раз и приведет к комплексному матричному представлению размерности* $2N \times 2N$. *При этом операторы* $\tilde{\tilde{m}}_{I_\pm}$, $\tilde{m}_{I_\pm}$ *будут удовлетворять следующим соотношениям*

$$\left\{ \begin{array}{l} \tilde{m}_{I_\pm \hat{A}}{}^{\tilde{B}} \tilde{m}_{I_\pm}{}^{\hat{C}}{}_{\tilde{B}} = \delta_{\hat{A}}{}^{\hat{C}}, \\ \\ \tilde{m}_{I_\pm \hat{A}}{}^{\tilde{B}} \tilde{m}_{I_\mp}{}^{\hat{C}}{}_{\tilde{B}} = 0, \\ \\ \tilde{m}_{I_\pm \hat{A}}{}^{\tilde{B}} \tilde{m}_{I_\pm}{}^{\hat{A}}{}_{\tilde{C}} = \tilde{\triangle}_{I_\pm \tilde{C}}{}^{\tilde{B}}, \end{array} \right. \qquad \left\{ \begin{array}{l} \tilde{\tilde{m}}_{I_\pm \hat{A}}{}^{\tilde{B}} \tilde{\tilde{m}}_{I_\pm}{}^{\hat{C}}{}_{\tilde{B}} = \delta_{\hat{A}}{}^{\hat{C}}, \\ \\ \tilde{\tilde{m}}_{I_\pm \hat{A}}{}^{\tilde{B}} \tilde{\tilde{m}}_{I_\mp}{}^{\hat{C}}{}_{\tilde{B}} = 0, \\ \\ \tilde{\tilde{m}}_{I_\pm \hat{A}}{}^{\tilde{B}} \tilde{\tilde{m}}_{I_\pm}{}^{\hat{A}}{}_{\tilde{C}} = \tilde{\tilde{\triangle}}_{I_\pm \tilde{C}}{}^{\tilde{B}}. \end{array} \right. \tag{3.23}$$

Здесь для I *шага* $\tilde{A}, \quad \tilde{B}, \quad \tilde{C}, \quad \tilde{D}, ... = \overline{1, \frac{(2N)^2}{2^{I-1}}}$, $\hat{A}, \hat{B}, \hat{C}, \hat{D}, ... = \overline{1, \frac{(2N)^2}{2^I}}$. *Тогда можно построить операторы* $(\tilde{m}_{J_{z_J}} := \tilde{m}_{J_\pm \hat{A}}{}^{\tilde{B}}$, $\tilde{\tilde{m}}^*_{J_{z_J}} := \tilde{\tilde{m}}_{J_\pm}{}^{\hat{A}}{}_{\tilde{B}})$

$$\tilde{M}_K := \tilde{m}_{\frac{n}{2}_{z_{\frac{n}{2}}}} \tilde{m}_{\frac{n}{2}-1_{z_{\frac{n}{2}-1}}} \cdot ... \cdot \tilde{m}_{2_{z_2}} \tilde{m}_{1_{z_1}}, \quad \tilde{\tilde{M}}_K := \tilde{\tilde{m}}^*_{\frac{n}{2}_{z_{\frac{n}{2}}}} \tilde{\tilde{m}}^*_{\frac{n}{2}-1_{z_{\frac{n}{2}-1}}} \cdot ... \cdot \tilde{\tilde{m}}^*_{2_{z_2}} \tilde{\tilde{m}}^*_{1_{z_1}}, \tag{3.24}$$

где z_J $(J = \overline{1, \frac{n}{2}})$ *равно 0 при выборе знака* $\ll - \gg$ *или 1 при выборе знака* $\ll + \gg$, *тогда* $K = \sum\limits_{J=1}^{\frac{n}{2}} z_J \cdot 2^{J-1} + 1$. *С помощью этих операторов можно определить*

$$(\gamma_K)_\Lambda := \tilde{M}_K (m_\Lambda{}^{\tilde{A}} \gamma_{\tilde{A}}) \tilde{\tilde{M}}_K. \tag{3.25}$$

Соответствующий пример будет разобран ниже (алгоритм 8.1).

4 Вещественные вложения. Инволюции

В этом параграфе рассказывается о том, как в комплексную реализацию $\mathbb{C}V_n$ *2n-мерного псевдориманова пространства локально вложить действительное n-мерное (псевдо-)риманово пространство. Для этого рассматривается вещественная поверхность в* $\mathbb{C}V_n$ *с вещественной параметризацией. Это индуцирует в касательном расслоении оператор вложения* $H_i{}^\Lambda$ *[15], с помощью которого можно получить слой вещественного касательного расслоения, снабженный (псевдо-)евклидовой метрикой. Сигнатура такой метрики будет существенно зависеть от вида оператора вложения. Вывод всех результатов этого параграфа сделан на основе [15].*

Вещественное (псевдо-)риманово пространство V_n будем рассматривать как поверхность вещественной размерности n в пространстве $\mathbb{C}V_n$, т.е. определять с помощью параметрического уравнения

$$w^\Lambda = w^\Lambda(u^i), \quad (\Lambda, \Psi, ..., i, j, g, h = \overline{1, n}), \tag{4.1}$$

где w^Λ - комплексные координаты точки x базы, а u^i - параметры: локальные координаты точки пространства V_n. Частные производные $(\partial_i w^\Lambda =: H_i{}^\Lambda)$ определяют локальное вложение вещественного касательного пространства поверхности (4.1) в комплексное касательное пространство $\tau_x^\mathbb{C}$ следующим образом

$$H : \tau_x^\mathbb{R} \longmapsto \tau_x^\mathbb{C}, \tag{4.2}$$

$$\begin{gathered} w^\Lambda = w^\Lambda(u^i(t)), \quad r^\Lambda := \tfrac{dw^\Lambda}{dt} = H_i{}^\Lambda \tfrac{du^i}{dt} =: H_i{}^\Lambda r^i \\ \tfrac{du^i}{dt} \in \tau_x^\mathbb{R} \longmapsto \tfrac{dw^\Lambda}{dt} \in \tau_x^\mathbb{C}, \end{gathered} \tag{4.3}$$

где дифференцирование ведется вдоль вещественной кривой $\gamma(t)$ поверхности (4.1). Так как матрица $\| \, H_i{}^\Lambda \, \|$ есть невырожденная якобиева матрица, то существует оператор $H^i{}_\Lambda$ такой, что

$$\begin{cases} H^i{}_\Lambda H_i{}^\Psi = \delta_\Lambda{}^\Psi, \\ H^i{}_\Lambda H_j{}^\Lambda = \delta_j{}^i. \end{cases} \tag{4.4}$$

Отсюда следует, что оператор $H_i{}^\Lambda$ определяет в комплексном пространстве инволюцию

$$S_\Lambda{}^{\Psi'} = H^i{}_\Lambda \bar{H}_i{}^{\Psi'}, \tag{4.5}$$

где координаты $\bar{H}_i{}^{\Psi'}$ комплексно сопряжены координатам $H_i{}^\Psi$. Поэтому

$$r^i = H^i{}_\Lambda r^\Lambda = \overline{H^i{}_\Lambda r^\Lambda} \quad \Rightarrow \quad S_\Lambda{}^{\Psi'} r^\Lambda = \bar{r}^{\Psi'}. \tag{4.6}$$

Это есть необходимое и достаточное условие того, что вектор $r^\Lambda \in \tau_x^\mathbb{C}$ будет вещественным. При этом

$$S_\Lambda{}^{\Psi'} \bar{S}_{\Psi'}{}^\Phi = \delta_\Lambda{}^\Phi. \tag{4.7}$$

Метрику V_n (вещественного (псевдо-)риманова пространства) определим условием

$$g_{\Lambda\Psi} r^\Lambda r^\Psi = \overline{g_{\Lambda\Psi} r^\Lambda r^\Psi}, \quad \forall \bar{r}^{\Psi'} = S_\Lambda{}^{\Psi'} r^\Lambda. \tag{4.8}$$

Это означает, что вещественный тензор пространства V_n определяется как тензор, самосопряженный относительно указанной эрмитовой инволюции

$$g_{\Lambda\Psi} = S_\Lambda{}^{\Phi'} S_\Psi{}^{\Theta'} \bar{g}_{\Phi'\Theta'}. \tag{4.9}$$

Поэтому тензор

$$g_{ij} := H_i{}^{\Lambda} H_j{}^{\Psi} g_{\Lambda\Psi} = \overline{H_i{}^{\Lambda} H_j{}^{\Psi} g_{\Lambda\Psi}} \tag{4.10}$$

будет метрическим тензором $V_n \subset \mathbb{C}V_n$. Вид метрики g_{ij} существенно зависит от структуры оператора $H_i{}^{\Lambda}$ и, следовательно, от тензора инволюции $S_{\Lambda}{}^{\Psi'}$.

Example 4.1. *Пусть имеется комплексно-евклидово пространство $\mathbb{CR}^4$. Определить вложение действительного пространства в комплексное можно тремя существенно различными способами*

$$H_i{}^{\Lambda} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad H_i{}^{\Lambda} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \end{pmatrix}, \quad H_i{}^{\Lambda} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \end{pmatrix}.$$

$$G_{\Lambda\Psi} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

$$g_{ij} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad g_{ij} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad g_{ij} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}. \tag{4.11}$$

5 Элементарные преобразования ортогональной группы

В этом параграфе рассказывается о том, как привести псевдоортогональное преобразование к блочно-диагональному виду, а затем перейти к комплексной реализации такого преобразования. Вывод всех результатов этого параграфа сделан на основе [10], [20].

Рассмотрим ортогональные преобразования в пространстве $\mathbb{R}^{2n}_{(n,n)}$, которые задаются соотношением

$$G_{\tilde\Lambda\tilde\Psi} S_{\tilde\Omega}{}^{\tilde\Lambda} S_{\tilde\Gamma}{}^{\tilde\Psi} = G_{\tilde\Omega\tilde\Gamma}. \tag{5.1}$$

При этом, базис выберем так, что $\| \, G_{\tilde\Lambda\tilde\Psi} \, \|$ будет иметь в нем диагональный вид.

Теорема 5.1. *Конформные преобразования пространства* $\mathbb{R}^{2n-2}_{(n-1,n-1)}$ *образуют группу* $O(n,n)$, *состоящую из:*

1. *вращений из* $O(n-1,n-1)$,
2. *дилатаций,*
3. *трансляций,*
4. *инверсий,*
5. *суперпозиции 1-4.*

Тогда любое преобразование из $O(n,n)$ *представимо в виде произведения элементарных преобразований*

$$S_{\tilde\Lambda_1}{}^{\tilde\Lambda_{J+1}} = \pm\prod_{I=1}^{J}(\pm(r_I)_{\tilde\Lambda_I}(r_I)^{\tilde\Lambda_{I+1}} - \delta_{\tilde\Lambda_I}{}^{\tilde\Lambda_{I+1}}), \quad (r_I)_{\tilde\Lambda_I}(r_I)^{\tilde\Lambda_I} = \pm 2. \tag{5.2}$$

Доказательство. Элементарные преобразования из (5.2) действительно ортогональны

$$(\pm r_{\tilde\Lambda} r^{\tilde\Psi} - \delta_{\tilde\Lambda}{}^{\tilde\Psi})(\pm r_{\tilde\Psi} r^{\tilde\Theta} - \delta_{\tilde\Psi}{}^{\tilde\Theta}) = (\pm 2) r_{\tilde\Lambda} r^{\tilde\Theta} \mp r_{\tilde\Lambda} r^{\tilde\Theta} \mp r_{\tilde\Lambda} r^{\tilde\Theta} + \delta_{\tilde\Lambda}{}^{\tilde\Theta} = \delta_{\tilde\Lambda}{}^{\tilde\Theta}. \tag{5.3}$$

Обычные вращения из группы $O(2)$ представимы в виде (5.2), при этом, $r^{\tilde\Lambda} = \sqrt{2}(\cos\frac{\alpha}{2},\ \sin\frac{\alpha}{2})$, и имеется два несвязных класса вращений

$$a).\begin{pmatrix} \cos\alpha & \sin\alpha \\ \sin\alpha & -\cos\alpha \end{pmatrix}, \quad b).\begin{pmatrix} \cos\alpha & \sin\alpha \\ -\sin\alpha & \cos\alpha \end{pmatrix}. \tag{5.4}$$

Однако преобразование b). получается суперпозицией двух преобразований из a).

1. Преобразования из группы $O(1,1)$ представляют собой буст одного из 4 видов

$$\begin{aligned} &a). \begin{pmatrix} \operatorname{ch}\theta & \operatorname{sh}\theta \\ \operatorname{sh}\theta & \operatorname{ch}\theta \end{pmatrix}, \quad b). \begin{pmatrix} \operatorname{ch}\theta & -\operatorname{sh}\theta \\ \operatorname{sh}\theta & -\operatorname{ch}\theta \end{pmatrix}, \\ &c). \begin{pmatrix} -\operatorname{ch}\theta & \operatorname{sh}\theta \\ -\operatorname{sh}\theta & \operatorname{ch}\theta \end{pmatrix}, \quad d). \begin{pmatrix} -\operatorname{ch}\theta & -\operatorname{sh}\theta \\ -\operatorname{sh}\theta & -\operatorname{ch}\theta \end{pmatrix}. \end{aligned} \tag{5.5}$$

   Преобразование b). представимо в виде (5.2), где $r^{\tilde\Lambda} = \sqrt{2}(\operatorname{ch}\frac{\theta}{2},\ \ \operatorname{sh}\frac{\theta}{2})$. Преобразование a). есть суперпозиция двух преобразований вида b). Преобразования c). и d). отличаются лишь знаком от преобразований b). и a). соответственно. Этим описываются 4 несвязные компоненты группы $O(1,1)$.

2. Пусть одномерная дилатация в $\mathbb{R}^1$ имеет вид: $\tilde{x} = \lambda x$. Рассмотрим световой конус в $\mathbb{R}^3_{(1,2)}$, заданный как $T^2 - Z^2 - X^2 = 0$. Рассечем его плоскостью $T + Z = 1$, а затем выполним стереографическую проекцию сечения на прямую $T = 1,\ Z = 0$, что индуцирует одномерное пространство $\mathbb{R}^1$ с единственной координатой $x = \frac{X}{T-Z}$. Тогда $\tilde{T}+\tilde{Z} = \lambda(T+Z)$, $\tilde{T}-\tilde{Z} = \lambda^{-1}(T-Z)$, $\tilde{X} = X$. При $\lambda > 0$ данное преобразование будет бустом вида a).; при $\lambda < 0$ данное преобразование будет бустом вида d).

3. Рассмотрим трансляцию $\tilde{x} = x + a$ пространства $\mathbb{R}^1$. Это приведет к соотношениям: $\tilde{T} - \tilde{Z} = T - Z$, $\tilde{T} + \tilde{Z} = T + Z + 2aX + a^2(T - Z)$, $\tilde{X} = X + a(T - Z)$. Неподвижным вектором данного преобразования будет изотропный вектор вида $(b, 0, b)$. Никаким псевдоортогональным преобразованием невозможно перевести изотропный вектор в неизотропный, поскольку такие преобразования сохраняют величину $T^2 - X^2 - Z^2$ неизменной. Это означает, что невозможно подействовать такими преобразованиями на базис так, чтобы изотропный вектор был направлен вдоль неизотропной координатной оси. Однако, это не мешает разложить дилатацию на элементарные преобразования: вращение, буст и еще одно такое же вращение. Для этого в качестве вращения возьмем элементарное преобразование
$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\alpha & -\sin\alpha \\ 0 & \sin\alpha & \cos\alpha \end{pmatrix} \tag{5.6}$$
с $\operatorname{tg}\alpha = \frac{a}{2}$, а в качестве буста вида а). - элементарное преобразование
$$\begin{pmatrix} \operatorname{ch}\theta & \operatorname{sh}\theta & 0 \\ \operatorname{sh}\theta & \operatorname{ch}\theta & 0 \\ 0 & 0 & 1 \end{pmatrix} \tag{5.7}$$
с $\operatorname{ch}\theta = \frac{a^2}{2} + 1$. Тогда указанная композиция будет иметь вид
$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{1+\frac{a^2}{4}}} & -\frac{\frac{a}{2}}{\sqrt{1+\frac{a^2}{4}}} \\ 0 & \frac{\frac{a}{2}}{\sqrt{1+\frac{a^2}{4}}} & \frac{1}{\sqrt{1+\frac{a^2}{4}}} \end{pmatrix} \begin{pmatrix} 1 + \frac{a^2}{2} & a\sqrt{1 + \frac{a^2}{4}} & 0 \\ a\sqrt{1 + \frac{a^2}{4}} & 1 + \frac{a^2}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix} \cdot$$
$$\cdot \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{1+\frac{a^2}{4}}} & -\frac{\frac{a}{2}}{\sqrt{1+\frac{a^2}{4}}} \\ 0 & \frac{\frac{a}{2}}{\sqrt{1+\frac{a^2}{4}}} & \frac{1}{\sqrt{1+\frac{a^2}{4}}} \end{pmatrix} = \begin{pmatrix} 1 + \frac{a^2}{2} & a & -\frac{a^2}{2} \\ a & 1 & -a \\ \frac{a^2}{2} & a & 1 - \frac{a^2}{2} \end{pmatrix}. \tag{5.8}$$

4. Инверсия $\tilde{x} = \frac{1}{x}$ в $\mathbb{R}^1$ индуцирует вращение $\tilde{T} = T$, $\tilde{Z} = -Z$, $\tilde{X} = X$.

5. Суперпозиции всех одномерных преобразований вида 1-4 вместе с вращениями $O(n-1, n-1)$ представляют группу конформных преобразований пространства $\mathbb{R}^{2n-2}_{(n-1,n-1)}$, которая будет изоморфна группе $O(n, n)$.

$\square$

Example 5.1. *Рассмотрим группу псевдоортогональных преобразований $O(1,2)$ пространства $\mathbb{R}^3_{(1,2)}$. Как известно, собственные значения любого такого преобразования являются корнями многочлена третьей степени. Одно из таких значений обязательно должно быть вещественным числом. Поскольку псевдоортогональные преобразования удовлетворяют соотношению (5.1), то квадрат такого вещественного числа есть единица. Поэтому любое преобразование из группы $O(1,2)$ обладает неподвижной осью. Однако, не всегда эту ось можно совместить с координатной осью, используя только псевдоортогональные преобразования базиса.*

1. *Рассмотрим композицию вращения и буста для пространства $\mathbb{R}^3_{(1,2)}$*

$$\begin{pmatrix} \operatorname{ch}\theta & -\operatorname{sh}\theta & 0 \\ \operatorname{sh}\theta & -\operatorname{ch}\theta & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\alpha & \sin\alpha \\ 0 & \sin\alpha & -\cos\alpha \end{pmatrix} = \begin{pmatrix} \operatorname{ch}\theta & -\cos\alpha\operatorname{sh}\theta & -\sin\alpha\operatorname{sh}\theta \\ \operatorname{sh}\theta & -\cos\alpha\operatorname{ch}\theta & -\sin\alpha\operatorname{ch}\theta \\ 0 & \sin\alpha & -\cos\alpha \end{pmatrix}. \tag{5.9}$$

Пусть эта композиция вектор (a,b,c) оставит неподвижным

$$\left\{\begin{array}{rcl} a\operatorname{ch}\theta - b\cos\alpha\operatorname{sh}\theta - c\sin\alpha\operatorname{sh}\theta & = & a \\ a\operatorname{sh}\theta - b\cos\alpha\operatorname{ch}\theta - c\sin\alpha\operatorname{ch}\theta & = & b \\ b\sin\alpha - c\cos\alpha & = & c \end{array}\right. , \qquad \begin{array}{l} \frac{c}{b} = \operatorname{tg}\frac{\alpha}{2}, \quad \frac{a}{b} = \operatorname{cth}\frac{\theta}{2}, \\ \\ r = (b\operatorname{cth}\frac{\theta}{2}, b, b\operatorname{tg}\frac{\alpha}{2}). \end{array} \tag{5.10}$$

Проведем преобразование базиса

$$\begin{pmatrix} \operatorname{ch}\frac{\theta}{2} & -\operatorname{sh}\frac{\theta}{2} & 0 \\ -\operatorname{sh}\frac{\theta}{2} & \operatorname{ch}\frac{\theta}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} b\operatorname{cth}\frac{\theta}{2} \\ b \\ b\operatorname{tg}\frac{\alpha}{2} \end{pmatrix} = \begin{pmatrix} b/\operatorname{sh}\frac{\theta}{2} \\ 0 \\ b\operatorname{tg}\frac{\alpha}{2} \end{pmatrix}. \tag{5.11}$$

Поскольку теперь $|r|^2 = b^2(1/\operatorname{sh}^2\frac{\theta}{2} - \operatorname{tg}^2\frac{\alpha}{2})$, положим $b := \pm\frac{1}{\sqrt{|1/\operatorname{sh}^2\frac{\theta}{2} - \operatorname{tg}^2\frac{\alpha}{2}|}}$. Тогда существует преобразование базиса

$$\begin{pmatrix} \operatorname{ch}\phi & 0 & \operatorname{sh}\phi \\ 0 & 1 & 0 \\ \operatorname{sh}\phi & 0 & \operatorname{ch}\phi \end{pmatrix} \begin{pmatrix} b/\operatorname{sh}\frac{\theta}{2} \\ 0 \\ b\operatorname{tg}\frac{\alpha}{2} \end{pmatrix}, \qquad \begin{array}{l} a).\ \text{либо}\ \operatorname{ch}\phi = \frac{b}{\operatorname{sh}\frac{\theta}{2}}, \quad \operatorname{sh}\phi = -b\operatorname{tg}\frac{\alpha}{2}, \\ b).\ \text{либо}\ \operatorname{sh}\phi = -\frac{b}{\operatorname{sh}\frac{\theta}{2}}, \quad \operatorname{ch}\phi = \ b\operatorname{tg}\frac{\alpha}{2}. \end{array} \tag{5.12}$$

В случае a). неподвижна координатная ось (1,0,0), а в случае b). - ось (0,0,1). Если $1/\operatorname{sh}\frac{\theta}{2} = \pm\operatorname{tg}\frac{\alpha}{2}$, то никаким псевдоортогональным вращением базиса невозможно упростить первоначальную суперпозицию.

2. Рассмотрим композицию двух бустов для пространства $\mathbb{R}^3_{(1,2)}$

$$\begin{pmatrix} \operatorname{ch}\theta & -\operatorname{sh}\theta & 0 \\ \operatorname{sh}\theta & -\operatorname{ch}\theta & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \operatorname{ch}\psi & 0 & -\operatorname{sh}\psi \\ 0 & 1 & 0 \\ \operatorname{sh}\psi & 0 & -\operatorname{ch}\psi \end{pmatrix} = \begin{pmatrix} \operatorname{ch}\theta\operatorname{ch}\psi & -\operatorname{sh}\theta & -\operatorname{sh}\psi\operatorname{ch}\theta \\ \operatorname{sh}\theta\operatorname{ch}\psi & -\operatorname{ch}\theta & -\operatorname{sh}\psi\operatorname{sh}\theta \\ \operatorname{sh}\psi & 0 & -\operatorname{ch}\psi \end{pmatrix}. \tag{5.13}$$

Пусть эта композиция оставит вектор (a, b, c) *неподвижным*

$$\left\{\begin{array}{rcl} a\operatorname{ch}\theta\operatorname{ch}\psi - b\operatorname{sh}\theta - c\operatorname{sh}\psi\operatorname{ch}\theta & = & a \\ a\operatorname{sh}\theta\operatorname{ch}\psi - b\operatorname{ch}\theta - c\operatorname{sh}\psi\operatorname{sh}\theta & = & b \\ a\operatorname{sh}\psi - c\operatorname{ch}\psi & = & c \end{array}\right., \qquad \begin{array}{l} \frac{c}{a} = \operatorname{th}\frac{\psi}{2}, \quad \frac{b}{a} = \operatorname{th}\frac{\theta}{2}, \\ \\ r = (a, a\operatorname{th}\frac{\theta}{2}, a\operatorname{th}\frac{\psi}{2}). \end{array} \tag{5.14}$$

Проведем преобразование базиса

$$\begin{pmatrix} \operatorname{ch}\frac{\theta}{2} & -\operatorname{sh}\frac{\theta}{2} & 0 \\ -\operatorname{sh}\frac{\theta}{2} & \operatorname{ch}\frac{\theta}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a \\ a\operatorname{th}\frac{\theta}{2} \\ a\operatorname{th}\frac{\psi}{2} \end{pmatrix} = \begin{pmatrix} a/\operatorname{ch}\frac{\theta}{2} \\ 0 \\ a\operatorname{th}\frac{\psi}{2} \end{pmatrix}. \tag{5.15}$$

Поскольку теперь $|r|^2 = a^2(1/\operatorname{ch}^2\frac{\theta}{2} - \operatorname{th}^2\frac{\psi}{2})$*, положим* $a := \pm\frac{1}{\sqrt{|1/\operatorname{ch}^2\frac{\theta}{2} - \operatorname{th}^2\frac{\psi}{2}|}}$*. Тогда существует преобразование базиса*

$$\begin{pmatrix} \operatorname{ch}\phi & 0 & \operatorname{sh}\phi \\ 0 & 1 & 0 \\ \operatorname{sh}\phi & 0 & \operatorname{ch}\phi \end{pmatrix} \begin{pmatrix} a/\operatorname{ch}\frac{\theta}{2} \\ 0 \\ a\operatorname{th}\frac{\psi}{2} \end{pmatrix}, \qquad \begin{array}{l} a).\ \textit{либо}\ \operatorname{ch}\phi = \frac{a}{\operatorname{ch}\frac{\theta}{2}}, \quad \operatorname{sh}\phi = -a\operatorname{th}\frac{\psi}{2}, \\ b).\ \textit{либо}\ \operatorname{sh}\phi = -\frac{a}{\operatorname{ch}\frac{\theta}{2}}, \quad \operatorname{ch}\phi = a\operatorname{th}\frac{\psi}{2}. \end{array} \tag{5.16}$$

В случае a). неподвижна координатная ось (1,0,0), а в случае b). - ось (0,0,1). Если $1/\operatorname{ch}^2\frac{\theta}{2} = \pm\operatorname{th}^2\frac{\psi}{2}$*, то никаким псевдоортогональным вращением базиса невозможно упростить первоначальную суперпозицию.*

Example 5.2. *Рассмотрим суперпозицию элементарных преобразований из группы $O(1,2)$*

$$\begin{pmatrix} \operatorname{ch}\phi & 0 & \operatorname{sh}\phi \\ 0 & 1 & 0 \\ \operatorname{sh}\phi & 0 & \operatorname{ch}\phi \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\alpha & \sin\alpha \\ 0 & -\sin\alpha & \cos\alpha \end{pmatrix} \begin{pmatrix} \operatorname{ch}\psi & \operatorname{sh}\psi & 0 \\ \operatorname{sh}\psi & \operatorname{ch}\psi & 0 \\ 0 & 0 & 1 \end{pmatrix} =$$
$$= \begin{pmatrix} \operatorname{ch}\tilde\phi & 0 & \operatorname{sh}\tilde\phi \\ 0 & 1 & 0 \\ \operatorname{sh}\tilde\phi & 0 & \operatorname{ch}\tilde\phi \end{pmatrix} \begin{pmatrix} \operatorname{ch}\tilde\psi & \operatorname{sh}\tilde\psi & 0 \\ \operatorname{sh}\tilde\psi & \operatorname{ch}\tilde\psi & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\tilde\alpha & \sin\tilde\alpha \\ 0 & -\sin\tilde\alpha & \cos\tilde\alpha \end{pmatrix},$$
$$\begin{pmatrix} \operatorname{ch}(\phi-\tilde\phi)\operatorname{ch}\psi-\operatorname{sh}(\phi-\tilde\phi)\operatorname{sh}\psi\sin\alpha & \operatorname{ch}(\phi-\tilde\phi)\operatorname{sh}\psi-\operatorname{sh}(\phi-\tilde\phi)\operatorname{ch}\psi\sin\alpha & \operatorname{sh}(\phi-\tilde\phi)\cos\alpha \\ \cos\alpha\operatorname{sh}\psi & \cos\alpha\operatorname{ch}\psi & \sin\alpha \\ \operatorname{sh}(\phi-\tilde\phi)\operatorname{ch}\psi-\operatorname{ch}(\phi-\tilde\phi)\operatorname{sh}\psi\sin\alpha & \operatorname{sh}(\phi-\tilde\phi)\operatorname{sh}\psi-\operatorname{ch}(\phi-\tilde\phi)\operatorname{ch}\psi\sin\alpha & \operatorname{ch}(\phi-\tilde\phi)\cos\alpha \end{pmatrix} =$$
$$= \begin{pmatrix} \operatorname{ch}\tilde\psi & \operatorname{sh}\tilde\psi\cos\tilde\alpha & \operatorname{sh}\tilde\psi\sin\tilde\alpha \\ \operatorname{sh}\tilde\psi & \operatorname{ch}\tilde\psi\cos\tilde\alpha & \operatorname{ch}\tilde\psi\sin\tilde\alpha \\ 0 & -\sin\tilde\alpha & \cos\tilde\alpha \end{pmatrix} \tag{5.17}$$

Решением этой системы уравнений будет

$$\begin{gathered} \sin\alpha = \operatorname{ch}\tilde\psi\sin\tilde\alpha, \quad \operatorname{sh}\psi = \frac{\operatorname{sh}\tilde\psi}{\sqrt{1-\operatorname{ch}^2\tilde\psi\sin^2\tilde\alpha}}, \quad \operatorname{sh}(\phi-\tilde\phi) = \frac{\operatorname{sh}\tilde\psi\sin\tilde\alpha}{\sqrt{1-\operatorname{ch}^2\tilde\psi\sin^2\tilde\alpha}}. \\ \sin\tilde\alpha = \frac{\sin\alpha}{\sqrt{1+\operatorname{sh}^2\psi\cos^2\alpha}}, \quad \operatorname{sh}\tilde\psi = \operatorname{sh}\psi\cos\alpha, \quad \operatorname{sh}(\phi-\tilde\phi) = \frac{\operatorname{sh}\psi\sin\alpha}{\sqrt{1+\operatorname{sh}^2\psi\cos^2\alpha}}. \end{gathered} \tag{5.18}$$

Опять же исключительным случаем является $\operatorname{ch}^2\tilde\psi\sin^2\tilde\alpha \geq 1$. *Поэтому, не всегда возможно переставить в суперпозиции элементарные преобразования, в отличие от ортогональной группы* $O(3)$.

Example 5.3. *Рассмотрим теперь преобразования из группы $O(2,2)$. Любое преобразование из этой группы согласно теореме Картана-Дьедонне [10, т. 2, с. 33] можно представить с помощью четырех преобразований вида (5.2)*

$$\pm\, r_{\tilde\Lambda} r^{\tilde\Psi} - \delta_{\tilde\Lambda}{}^{\tilde\Psi}, \quad r_{\tilde\Lambda} r^{\tilde\Lambda} = \pm 2. \tag{5.19}$$

Однако, всегда можно подобрать разложение таким образом, чтобы два из таких преобразований имели диагональную матрицу. Элементы главной диагонали такой матрицы $I_{\tilde\Lambda}{}^{\tilde\Psi}$ *по модулю равны 1. Поэтому (5.2) может быть переписано как*

$$\begin{gathered} (I_1)_{\tilde\Phi}{}^{\tilde\Lambda}(\pm(r_1)_{\tilde\Lambda}(r_1)^{\tilde\Psi} - \delta_{\tilde\Lambda}{}^{\tilde\Psi})(I_2)_{\tilde\Psi}{}^{\tilde\Theta}(\pm(r_2)_{\tilde\Theta}(r_2)^{\tilde\Omega} - \delta_{\tilde\Theta}{}^{\tilde\Omega}), \\ (r_1)^{\tilde\Psi}(I_2)_{\tilde\Psi}{}^{\tilde\Theta}(r_2)_{\tilde\Theta} = 0, \quad (r_2)^{\tilde\Psi}(I_1)_{\tilde\Psi}{}^{\tilde\Theta}(r_1)_{\tilde\Theta} = 0. \end{gathered} \tag{5.20}$$

Существует 3 существенно различных взаимоисключающих варианта.

1. *В этом случае псевдоортогональное преобразование может иметь блочно-диагональный вид*

$$I_1 \begin{pmatrix} \operatorname{ch}\theta & -\operatorname{sh}\theta & 0 & 0 \\ \operatorname{sh}\theta & -\operatorname{ch}\theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} I_2 \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \operatorname{ch}\phi & -\operatorname{sh}\phi \\ 0 & 0 & \operatorname{sh}\phi & -\operatorname{ch}\phi \end{pmatrix} =$$

$$I_1 \left(\underbrace{\begin{pmatrix} \sqrt{2}\operatorname{sh}\frac{\theta}{2} \\ \sqrt{2}\operatorname{ch}\frac{\theta}{2} \\ 0 \\ 0 \end{pmatrix}}_{(r_1)_{\tilde{\Theta}}} \underbrace{\begin{pmatrix} \sqrt{2}\operatorname{sh}\frac{\theta}{2} & -\sqrt{2}\operatorname{ch}\frac{\theta}{2} & 0 & 0 \end{pmatrix}}_{(r_1)^{\tilde{\Omega}}} + \underbrace{\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\delta_{\tilde{\Theta}}{}^{\tilde{\Omega}}} \right) \tag{5.21}$$

$$I_2 \left(\underbrace{\begin{pmatrix} 0 \\ 0 \\ \sqrt{2}\operatorname{sh}\frac{\phi}{2} \\ \sqrt{2}\operatorname{ch}\frac{\phi}{2} \end{pmatrix}}_{(r_2)_{\tilde{\Theta}}} \underbrace{\begin{pmatrix} 0 & 0 & \sqrt{2}\operatorname{sh}\frac{\phi}{2} & -\sqrt{2}\operatorname{ch}\frac{\phi}{2} \end{pmatrix}}_{(r_2)^{\tilde{\Omega}}} + \underbrace{\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\delta_{\tilde{\Theta}}{}^{\tilde{\Omega}}} \right),$$

$$(r_1)^{\tilde{\Psi}}(r_1)_{\tilde{\Psi}} = -2, \quad (r_2)^{\tilde{\Psi}}(r_2)_{\tilde{\Psi}} = -2.$$

2. *В этом случае псевдоортогональное преобразование может иметь блочно-диагональный вид*

$$I_1 \begin{pmatrix} \cos\theta & \sin\theta & 0 & 0 \\ \sin\theta & -\cos\theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} I_2 \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -\cos\phi & -\sin\phi \\ 0 & 0 & -\sin\phi & \cos\phi \end{pmatrix} =$$

$$I_1 \left(\underbrace{\begin{pmatrix} \sqrt{2}\cos\frac{\theta}{2} \\ \sqrt{2}\sin\frac{\theta}{2} \\ 0 \\ 0 \end{pmatrix}}_{(r_1)_{\tilde{\Theta}}} \underbrace{\begin{pmatrix} \sqrt{2}\cos\frac{\theta}{2} & \sqrt{2}\sin\frac{\theta}{2} & 0 & 0 \end{pmatrix}}_{(r_1)^{\tilde{\Omega}}} - \underbrace{\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\delta_{\tilde{\Theta}}{}^{\tilde{\Omega}}} \right) \tag{5.22}$$

$$I_2\left(\underbrace{\begin{pmatrix}0\\0\\-\sqrt{2}\cos\frac{\phi}{2}\\-\sqrt{2}\sin\frac{\phi}{2}\end{pmatrix}}_{(r_2)_{\tilde{\Theta}}}\underbrace{\begin{pmatrix}0 & 0 & \sqrt{2}\cos\frac{\phi}{2} & \sqrt{2}\sin\frac{\phi}{2}\end{pmatrix}}_{(r_2)^{\tilde{\Omega}}}+\underbrace{\begin{pmatrix}1&0&0&0\\0&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}}_{\delta_{\tilde{\Theta}}{}^{\tilde{\Omega}}}\right),$$

$$(r_1)^{\tilde{\Psi}}(r_1)_{\tilde{\Psi}}=2,\quad (r_2)^{\tilde{\Psi}}(r_2)_{\tilde{\Psi}}=-2.$$

3. *В этом случае псевдоортогональное преобразование не будет иметь блочно-диагонального вида, но может быть представлено как*

$$\begin{pmatrix}1&0&0&0\\0&1+\frac{a^2}{2}&a&-\frac{a^2}{2}\\0&a&1&-a\\0&\frac{a^2}{2}&a&1-\frac{a^2}{2}\end{pmatrix}\begin{pmatrix}-1&-b&0&b\\b&-1+\frac{b^2}{2}&0&-\frac{b^2}{2}\\0&0&-1&0\\b&\frac{b^2}{2}&0&-1-\frac{b^2}{2}\end{pmatrix}=$$

$$=\underbrace{\begin{pmatrix}1&0&0&0\\0&1&0&0\\0&0&-1&0\\0&0&0&1\end{pmatrix}}_{(I_1)_{\tilde{\Phi}}{}^{\tilde{\Lambda}}}\left(\underbrace{\begin{pmatrix}0\\\frac{a}{\sqrt{2}}\\-\sqrt{2}\\\frac{a}{\sqrt{2}}\end{pmatrix}}_{(r_1)_{\tilde{\Lambda}}}\underbrace{\begin{pmatrix}0&\frac{a}{\sqrt{2}}&\sqrt{2}&-\frac{a}{\sqrt{2}}\end{pmatrix}}_{(r_1)^{\tilde{\Psi}}}+\underbrace{\begin{pmatrix}1&0&0&0\\0&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}}_{\delta_{\tilde{\Lambda}}{}^{\tilde{\Psi}}}\right) \tag{5.23}$$

$$\underbrace{\begin{pmatrix}-1&0&0&0\\0&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}}_{(I_2)_{\tilde{\Psi}}{}^{\tilde{\Theta}}}\left(\underbrace{\begin{pmatrix}\sqrt{2}\\\frac{b}{\sqrt{2}}\\0\\\frac{b}{\sqrt{2}}\end{pmatrix}}_{(r_2)_{\tilde{\Theta}}}\underbrace{\begin{pmatrix}\sqrt{2}&\frac{b}{\sqrt{2}}&0&-\frac{b}{\sqrt{2}}\end{pmatrix}}_{(r_2)^{\tilde{\Omega}}}-\underbrace{\begin{pmatrix}1&0&0&0\\0&1&0&0\\0&0&1&0\\0&0&0&1\end{pmatrix}}_{\delta_{\tilde{\Theta}}{}^{\tilde{\Omega}}}\right),$$

$$(r_1)^{\tilde{\Psi}}(r_1)_{\tilde{\Psi}}=-2,\quad (r_2)^{\tilde{\Psi}}(r_2)_{\tilde{\Psi}}=2.$$

Согласно все той же теореме Картана-Дьедонне любое преобразование из группы $O(2n,2n)$ в некотором базисе будет иметь блочно-диагональный вид, состоящий из блоков $O(2,2)$. Это означает, что выполнено (5.2)

$$S_{\tilde{\Lambda}_1}{}^{\tilde{\Lambda}_{J+1}}=\pm\prod_{I=1}^{J}(\pm(r_I)_{\tilde{\Lambda}_I}(r_I)^{\tilde{\Lambda}_{I+1}}-\delta_{\tilde{\Lambda}_I}{}^{\tilde{\Lambda}_{I+1}}). \tag{5.24}$$

Пусть теперь в пространстве $\mathbb{R}^{2n}_{(n,n)}$ задана каноническая комплексная структура, тогда в разложении (5.2) можно выделить $J/2$ сомножителей так, чтобы было выполнено

$$\begin{gathered} f_{\tilde{\Lambda}_{2I}}{}^{\tilde{\Lambda}_I}((r_I)_{\tilde{\Lambda}_I}(r_I)^{\tilde{\Lambda}_{I+1}}-\delta_{\tilde{\Lambda}_I}{}^{\tilde{\Lambda}_{I+1}})f_{\tilde{\Lambda}_{I+1}}{}^{\tilde{\Lambda}_{2I+1}}=(r_{2I})_{\tilde{\Lambda}_{2I}}(r_{2I})^{\tilde{\Lambda}_{2I+1}}+\delta_{\tilde{\Lambda}_{2I}}{}^{\tilde{\Lambda}_{2I+1}}, \\ (r_I)_{\tilde{\Lambda}}(r_I)^{\tilde{\Lambda}}=2, \quad (r_{2I})_{\tilde{\Lambda}}(r_{2I})^{\tilde{\Lambda}}=-2, \quad \forall I=\overline{1,J/2}. \end{gathered} \tag{5.25}$$

Этим для любого псевдо-ортогонального преобразования $S_{\tilde{\Lambda}}{}^{\tilde{\Psi}}$ определятся соотношения

$$\begin{gathered} S_{\Lambda}{}^{\Psi}:=m_{\Lambda}{}^{\tilde{\Lambda}}S_{\tilde{\Lambda}}{}^{\tilde{\Psi}}m^{\Psi}{}_{\tilde{\Psi}}, \quad S_{\Lambda}{}^{\Psi'}:=m_{\Lambda}{}^{\tilde{\Lambda}}S_{\tilde{\Lambda}}{}^{\tilde{\Psi}}\bar{m}^{\Psi'}{}_{\tilde{\Psi}}=0, \\ \bar{S}_{\Lambda'}{}^{\Psi}:=\bar{m}_{\Lambda'}{}^{\tilde{\Lambda}}S_{\tilde{\Lambda}}{}^{\tilde{\Psi}}m^{\Psi}{}_{\tilde{\Psi}}=0, \quad \bar{S}_{\Lambda'}{}^{\Psi'}:=\bar{m}_{\Lambda'}{}^{\tilde{\Lambda}}S_{\tilde{\Lambda}}{}^{\tilde{\Psi}}\bar{m}^{\Psi'}{}_{\tilde{\Psi}}, \\ -f_{\tilde{\Omega}}{}^{\tilde{\Lambda}}S_{\tilde{\Lambda}}{}^{\tilde{\Psi}}f_{\tilde{\Psi}}{}^{\tilde{\Theta}}=S_{\tilde{\Omega}}{}^{\tilde{\Theta}}. \end{gathered} \tag{5.26}$$

Таким образом определится комплексная симметрия

$$\begin{gathered} m_{\Theta}{}^{\tilde{\Theta}}(r_{\tilde{\Theta}}r^{\tilde{\Sigma}}-\delta_{\tilde{\Theta}}{}^{\tilde{\Sigma}})f_{\tilde{\Sigma}}{}^{\tilde{\Psi}}(r_{\tilde{\Psi}}r^{\tilde{\Omega}}-\delta_{\tilde{\Psi}}{}^{\tilde{\Omega}})f_{\tilde{\Omega}}{}^{\tilde{\Xi}}m^{\Xi}{}_{\tilde{\Xi}}= \\ =m_{\Theta}{}^{\tilde{\Theta}}(-r_{\tilde{\Theta}}r^{\tilde{\Sigma}}f_{\tilde{\Sigma}}{}^{\tilde{\Psi}}f_{\tilde{\Psi}}{}^{\tilde{\Xi}}-f_{\tilde{\Theta}}{}^{\tilde{\Psi}}r_{\tilde{\Psi}}r^{\tilde{\Omega}}f_{\tilde{\Omega}}{}^{\tilde{\Xi}}+f_{\tilde{\Theta}}{}^{\tilde{\Psi}}f_{\tilde{\Psi}}{}^{\tilde{\Xi}})m^{\Xi}{}_{\tilde{\Xi}}= \\ =(2r_{\Theta}r^{\Xi}-\delta_{\Theta}{}^{\Xi}). \end{gathered} \tag{5.27}$$

Кроме того,

$$r_{\Lambda}r^{\Lambda}=m_{\Lambda}{}^{\tilde{\Psi}}r_{\tilde{\Psi}}m^{\Lambda}{}_{\tilde{\Phi}}r^{\tilde{\Phi}}=\frac{1}{2}(\delta_{\tilde{\Phi}}{}^{\tilde{\Psi}}+if_{\tilde{\Phi}}{}^{\tilde{\Psi}})r_{\tilde{\Psi}}r^{\tilde{\Phi}}=\frac{1}{2}r_{\tilde{\Psi}}r^{\tilde{\Psi}}=1, \tag{5.28}$$

что окончательно подтверждает существование разложения

$$S_{\Lambda_1}{}^{\Lambda_{J+1}}=\pm\prod_{I=1}^{J}((r_I)_{\Lambda_I}(r_I)^{\Lambda_{I+1}}-\delta_{\Lambda_I}{}^{\Lambda_{I+1}}), \quad (r_I)_{\Lambda_I}(r_I)^{\Lambda_I}=2. \tag{5.29}$$

Однако, следует отметить, что не всегда $(r_I)_{\Lambda}(r_J)^{\Lambda}=0, \quad (I\neq J)$ ввиду существования изотропных векторов, что и было показано в предыдущем примере. Что касается блочной структуры преобразований из группы $O(n,\mathbb{C})$, то размерность блоков тоже будет равна четырем.

Example 5.4. *Пусть имеется трехмерное псевдоевклидово пространство* $\mathbb{R}^3_{(1,2)}$. *Рассмотрим преобразование*

$$S_i{}^j=\begin{pmatrix} 1+\frac{a^2}{2} & a & -\frac{a^2}{2} \\ a & 1 & -a \\ \frac{a^2}{2} & a & 1-\frac{a^2}{2} \end{pmatrix}. \tag{5.30}$$

Определим оператор вложения как

$$H^i{}_{\Lambda}=\begin{pmatrix} 1 & 0 & 0 \\ 0 & -i & 0 \\ 0 & 0 & -i \end{pmatrix}, \quad H_i{}^{\Lambda}=\begin{pmatrix} 1 & 0 & 0 \\ 0 & i & 0 \\ 0 & 0 & i \end{pmatrix}. \tag{5.31}$$

Это определит комплексификацию нашего преобразования

$$S_{\Lambda}{}^{\Psi} = H^{i}{}_{\Lambda} S_{i}{}^{j} H_{j}{}^{\Psi} = \begin{pmatrix} 1+\frac{a^2}{2} & ia & -i\frac{a^2}{2} \\ -ia & 1 & -a \\ -i\frac{a^2}{2} & a & 1-\frac{a^2}{2} \end{pmatrix} \tag{5.32}$$

с комплексно-евклидовой метрикой. Чтобы перейти к действительной реализации, нам потребуются операторы следующего вида

$$(m^T)_{\tilde{\varLambda}}{}^{\Lambda} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & i & 0 \\ 0 & 0 & i \\ -i & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad m_{\Lambda}{}^{\tilde{\varLambda}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 & i & 0 & 0 \\ 0 & -i & 0 & 0 & 1 & 0 \\ 0 & 0 & -i & 0 & 0 & 1 \end{pmatrix}. \tag{5.33}$$

Соответственно, действительная реализация нашего преобразования будет иметь вид

$$S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} = m^{\Lambda}{}_{\tilde{\varLambda}} S_{\Lambda}{}^{\Psi} m_{\Psi}{}^{\tilde{\varPsi}} + \bar{m}^{\Lambda'}{}_{\tilde{\varLambda}} S_{\Lambda'}{}^{\Psi'} \bar{m}_{\Psi'}{}^{\tilde{\varPsi}} = \begin{pmatrix} 1+\frac{a^2}{2} & a & -\frac{a^2}{2} & 0 & 0 & 0 \\ a & 1 & -a & 0 & 0 & 0 \\ \frac{a^2}{2} & a & 1-\frac{a^2}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1+\frac{a^2}{2} & a & -\frac{a^2}{2} \\ 0 & 0 & 0 & a & 1 & -a \\ 0 & 0 & 0 & \frac{a^2}{2} & a & 1-\frac{a^2}{2} \end{pmatrix}. \tag{5.34}$$

Видно, что такая реализация имеет блочно-диагональную структуру, однако комплексная реализация такой структуры не имеет.

Наряду с преобразованиями (5.26) имеет смысл рассматривать псевдоортогональные преобразования

$$\begin{gathered} S_{\Lambda}{}^{\Psi} := m_{\Lambda}{}^{\tilde{\varLambda}} S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} m^{\Psi}{}_{\tilde{\varPsi}} = 0, \quad S_{\Lambda}{}^{\Psi'} := m_{\Lambda}{}^{\tilde{\varLambda}} S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} \bar{m}^{\Psi'}{}_{\tilde{\varPsi}}, \\ \bar{S}_{\Lambda'}{}^{\Psi} := \bar{m}_{\Lambda'}{}^{\tilde{\varLambda}} S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} m^{\Psi}{}_{\tilde{\varPsi}}, \quad S_{\Lambda'}{}^{\Psi'} := \bar{m}_{\Lambda'}{}^{\tilde{\varLambda}} S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} \bar{m}^{\Psi'}{}_{\tilde{\varPsi}} = 0, \\ f_{\tilde{\varOmega}}{}^{\tilde{\varLambda}} S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} f_{\tilde{\varPsi}}{}^{\tilde{\varTheta}} = S_{\tilde{\varOmega}}{}^{\tilde{\varTheta}}, \quad S_{\tilde{\varLambda}}{}^{\tilde{\varPsi}} S_{\tilde{\varPsi}}{}^{\tilde{\varPhi}} = \delta_{\tilde{\varLambda}}{}^{\tilde{\varPhi}}, \quad S_{\Lambda}{}^{\Psi'} \bar{S}_{\Psi'}{}^{\Phi} = \delta_{\Lambda}{}^{\Phi}, \quad S_{\Lambda}{}^{\Psi'} S_{\Phi}{}^{\Omega'} \bar{g}_{\Psi'\Omega'} = g_{\Lambda\Phi}. \end{gathered} \tag{5.35}$$

Комплексная реализация таких преобразований будет являться инволюциями на пространстве $\mathbb{C}^n$, ответственными за некоторое вещественное вложение. Поэтому в разложении (5.2) можно выделить $J/2$ сомножителей так, чтобы было выполнено

$$\begin{gathered} f_{\tilde{\varLambda}_{2I}}{}^{\tilde{\varPsi}_I} \left((r_I)_{\tilde{\varPsi}_I} (r_I)^{\tilde{\varLambda}_{I+1}} - \delta_{\tilde{\varPsi}_I}{}^{\tilde{\varLambda}_{I+1}} \right) f_{\tilde{\varLambda}_{I+1}}{}^{\tilde{\varLambda}_{2I+1}} = -\left((r_{2I})_{\tilde{\varLambda}_{2I}} (r_{2I})^{\tilde{\varLambda}_{2I+1}} + \delta_{\tilde{\varLambda}_{2I}}{}^{\tilde{\varLambda}_{2I+1}} \right), \\ (r_I)_{\tilde{\varLambda}} (r_I)^{\tilde{\varLambda}} = 2, \quad (r_{2I})_{\tilde{\varLambda}} (r_{2I})^{\tilde{\varLambda}} = -2, \quad \forall I = \overline{1, J/2}. \end{gathered} \tag{5.36}$$

Таким образом собственные значения преобразования $S_\Lambda{}^{\Psi'}$ суть ±1. Это автоматически означает, что такое преобразование ортогонально диагонализируемо в комплексной области, а вещественная реализация в некотором базисе имеет блочно-диагональный вид

$$\begin{pmatrix} \cos\theta & \sin\theta & 0 & 0 & ...... \\ \sin\theta & -\cos\theta & 0 & 0 & ...... \\ 0 & 0 & -\cos\theta & -\sin\theta & ...... \\ 0 & 0 & -\sin\theta & \cos\theta & ...... \\ .... & .... & .... & .... & ...... \end{pmatrix} \tag{5.37}$$

и тоже диагонализируема. За более подробным изложением следует обратиться к [20, с. 79-126], [10, т. 2].

6 Уравнение Клиффорда. Двулистное накрытие

> *В этом параграфе рассказывается о том, как перейти к редуцированным спинорам Картана. Это даст возможность построить взаимно однозначное отображение между векторами пространства $\mathbb{C}^n$ и бивекторами спинорного пространства $\mathbb{C}^N$, что приведет к построению двулистного накрытия $Spin(n,\mathbb{C})/\{\pm1\} \cong SO(n,\mathbb{C})$. Вывод всех результатов этого параграфа сделан на основе [18], [22].*

Пусть имеется некоторое псевдориманово пространство V_{2n}, где n-число четное. Оно является вещественной реализацией некоторого комплексного пространства $\mathbb{C}V_n$. Пусть в каждой точке V_{2n} задан набор невырожденных операторов $\gamma_{\tilde{\varLambda}}$. $\tilde{\varLambda},\ \tilde{\varPsi},\ ... = \overline{1,2n}$. При этом будем считать, что каждый оператор $\gamma_{\tilde{\varLambda}} := m_{\tilde{\varLambda}}{}^{\Lambda}\gamma_\Lambda + \bar{m}_{\tilde{\varLambda}}{}^{\Lambda'}\gamma_{\Lambda'}$ может быть представлен вещественной матрицей $(2N)^2 \times (2N)^2$, где $N = 2^{n/2-1}$, и удовлетворяют условию (2.3)

$$\gamma_{\tilde{\varLambda}}\gamma_{\tilde{\varPsi}} + \gamma_{\tilde{\varPsi}}\gamma_{\tilde{\varLambda}} = G_{\tilde{\varLambda}\tilde{\varPsi}}, \tag{6.1}$$

где $G_{\tilde{\varLambda}\tilde{\varPsi}} = G_{\tilde{\varPsi}\tilde{\varLambda}}$ - невырожденный метрический тензор вида (3.11)-(3.12), заданный в вещественной реализации касательного расслоения к V_{2n}. Назовем комплексной реализацией операторов $\gamma_{\tilde{\varLambda}}$ операторы (3.25) (если не важно, какой из K операторов будет выбран, то номер K будем опускать)

$$\gamma_\Lambda := m_\Lambda{}^{\tilde{\varLambda}}(\tilde{M}\gamma_{\tilde{\varLambda}}\tilde{\tilde{M}}). \tag{6.2}$$

Они будут удовлетворять уравнению

$$\gamma_\Lambda\gamma_\Psi + \gamma_\Psi\gamma_\Lambda = g_{\Lambda\Psi}, \quad \bar{\gamma}_{\Lambda'}\bar{\gamma}_{\Psi'} + \bar{\gamma}_{\Psi'}\bar{\gamma}_{\Lambda'} = g_{\Lambda'\Psi'}. \tag{6.3}$$

Пусть r^Λ - вектор комплексного касательного расслоения $\tau^\mathbb{C}$. В качестве базы можно рассматривать как комплексную $\mathbb{C}V_n$, так и вещественную реализацию V_{2n}. Слой этого расслоения изоморфен пространству $\mathbb{C}^n$, вещественной реализацией которого будет псевдоевклидово пространство $R^{2n}_{(n,n)}$, где n - индекс псевдоевклидовой метрики (количество "$+$" на главной диагонали). Таким образом, вектору из пространства $\mathbb{C}^n(\mathbb{R}^{2n}_{(n,n)})$ можно однозначно сопоставить некоторый оператор из пространства $\mathbb{C}^{2N}$ по правилу

$$R := r^\Phi \gamma_\Phi. \tag{6.4}$$

Поскольку из уравнения Клиффорда следует, что

$$tr(\gamma_\Lambda \gamma_\Psi) = N g_{\Lambda\Psi}, \tag{6.5}$$

то любому такому оператору R можно однозначно сопоставить вектор пространства $\mathbb{C}^n(R^{2n}_{(n,n)})$, взяв след от обеих частей соответствия (6.4) с $\gamma^\Lambda := g^{\Lambda\Psi}\gamma_\Psi$

$$g^{\Lambda\Psi} tr(\gamma_\Psi R) = r^\Phi g^{\Lambda\Psi} tr(\gamma_\Psi \gamma_\Phi), \tag{6.6}$$

и тогда

$$\frac{1}{N} tr(\gamma^\Psi R) = r^\Psi. \tag{6.7}$$

Таким образом, получено взаимно однозначное соответствие между векторами r^Λ пространства $\mathbb{C}^n(\mathbb{R}^{2n}_{(n,n)})$ и операторами R (специального вида, конечно же), действующими на пространстве $\mathbb{C}^{2N}$. Используя уравнение Клиффорда (6.3), можно написать тождество

$$R\gamma_\Psi R = r^\Lambda \gamma_\Lambda \gamma_\Psi \gamma_\Phi r^\Phi = r^\Lambda (g_{\Lambda\Psi} - \gamma_\Psi \gamma_\Lambda)\gamma_\Phi r^\Phi = (r_\Psi r^\Phi - \frac{1}{2}(r_\Lambda r^\Lambda)\delta_\Psi{}^\Phi)\gamma_\Phi. \tag{6.8}$$

Пусть $r_\Lambda r^\Lambda := \pm 2$. Известно, что любое (псевдо-)ортогональное преобразование $S_\Psi{}^\Phi$ может быть представлено как конечное произведение элементарных преобразований вида $S_I := (r_I)_\Psi (r_I)^\Phi - \frac{1}{2}((r_I)_\Lambda (r_I)^\Lambda)\delta_\Psi{}^\Phi$ (I пробегает конечные значения от 1 до какого-нибудь конечного J, например). Доказательству этого факта посвящен предыдущий параграф. Определим

$$R_I := (r_I)^\Lambda \gamma_\Lambda, \quad S = \prod_{I=1}^{J} S_I = S_1 S_2 \dots S_J, \tag{6.9}$$

$$\tilde{S} = \prod_{I=1}^{J} R_I = R_1 R_2 \dots R_J, \quad \tilde{\tilde{S}} := \prod_{I=J}^{1} R_I = R_J \dots R_2 R_1. \tag{6.10}$$

Здесь $\tilde{S}$ есть правое произведение операторов R_I, а $\tilde{\tilde{S}}$ есть левое произведение операторов R_I. Поэтому матрицы операторов $\tilde{S}$ и $\tilde{\tilde{S}}$ в общем случае различны и никак друг с другом не связаны. Это позволит переписать соответствие (6.8) для ортогонального преобразования ${S_\Lambda}^\Psi$ как

$$ {S_\Lambda}^\Psi \gamma_\Psi = \tilde{S}\gamma_\Lambda \tilde{\tilde{S}}, \quad \tilde{\tilde{S}} = \tilde{S}^{-1}. \tag{6.11}$$

Уравнение (6.8) справедливо и для вещественной реализации, что даст

$$ {S_{\tilde{\Lambda}}}^{\tilde{\Psi}} \gamma_{\tilde{\Psi}} = \tilde{S}\gamma_{\tilde{\Lambda}} \tilde{\tilde{S}}, \quad {f_{\tilde{\Theta}}}^{\tilde{\Lambda}} {S_{\tilde{\Lambda}}}^{\tilde{\Psi}} {f_{\tilde{\Psi}}}^{\tilde{\Phi}} = -{S_{\tilde{\Theta}}}^{\tilde{\Phi}}. \tag{6.12}$$

Соответственно, уравнение (6.12) для инволюции перепишется как

$$ {S_{\tilde{\Lambda}}}^{\tilde{\Psi}} \gamma_{\tilde{\Psi}} = \tilde{S}\gamma_{\tilde{\Lambda}} \tilde{\tilde{S}}, \quad {f_{\tilde{\Theta}}}^{\tilde{\Lambda}} {S_{\tilde{\Lambda}}}^{\tilde{\Psi}} {f_{\tilde{\Psi}}}^{\tilde{\Phi}} = {S_{\tilde{\Theta}}}^{\tilde{\Phi}}, \quad {S_{\tilde{\Lambda}}}^{\tilde{\Psi}} {S_{\tilde{\Psi}}}^{\tilde{\Phi}} = {\delta_{\tilde{\Lambda}}}^{\tilde{\Phi}}. \tag{6.13}$$

Тогда для комплексной реализации получим тождество

$$ {S_\Lambda}^{\Psi'} \bar{\gamma}_{\Psi'} = \bar{\tilde{S}}\gamma_\Lambda \tilde{\tilde{S}}, \quad {S_\Lambda}^{\Psi'} {\bar{S}_{\Psi'}}{}^{\Phi} = {\delta_\Lambda}^\Phi, \quad \tilde{S}\bar{\tilde{S}} = \pm E, \quad \tilde{\tilde{S}}\bar{\tilde{\tilde{S}}} = \pm E. \tag{6.14}$$

Теперь возможно комплексифицировать уже и вещественную реализацию $\mathbb{R}^{2n}_{(n,n)}$, рассматривая вектор $r^{\tilde{\Lambda}}$ в том же базисе, но с комплексными коэффициентами. При этом $\gamma_{\tilde{\Lambda}}$ останутся прежними. В этом случае ${S_{\tilde{\Lambda}}}^{\tilde{\Psi}}$ из (6.12) будет уже комплексным преобразованием. В частности, в качестве такого преобразования можно рассмотреть преобразование $i{f_{\tilde{\Psi}}}^{\tilde{\Phi}}$ ввиду (3.14). Это означает, что комплексная структура разложима как

$$ i{f_{\tilde{\Lambda}}}^{\tilde{\Psi}} \gamma_{\tilde{\Psi}} = \tilde{I}\tilde{F}\gamma_{\tilde{\Lambda}} \tilde{\tilde{F}}\tilde{\tilde{I}}. \tag{6.15}$$

Для дальнейшего исследования этого соответствия необходимо изучить структуру операторов $\gamma_{\tilde{\Lambda}}$. Не ограничивая общности, можно рассматривать базис, в котором метрический тензор $G_{\tilde{\Lambda}\tilde{\Psi}}$ на главной диагонали имеет ±1, а остальные компоненты равны 0

$$ \gamma_{\tilde{\Lambda}}\gamma_{\tilde{\Psi}} = -\gamma_{\tilde{\Psi}}\gamma_{\tilde{\Lambda}}, \quad \tilde{\Lambda} \neq \tilde{\Psi}. \tag{6.16}$$

Построим оператор

$$ \gamma_0 := \prod_{\tilde{\Lambda}=1}^{2n} \gamma_{\tilde{\Lambda}}. \tag{6.17}$$

Тогда из (6.16) будет следовать

$$ \gamma_0\gamma_{\tilde{\Lambda}} = -\gamma_{\tilde{\Lambda}}\gamma_0, \tag{6.18}$$

$$ (\gamma_0 - \lambda E)\gamma_{\tilde{\Lambda}} = -\gamma_{\tilde{\Lambda}}(\gamma_0 + \lambda E).$$

Поэтому

$$ det(\gamma_0 - \lambda E) = 0 = det(\gamma_0 + \lambda E) \tag{6.19}$$

будет уравнением на собственные значения оператора γ_0. Видно, что указанные значения парны: $\pm\lambda$. Поэтому положительным и отрицательным собственным значениям будут соответствовать 2 различных собственных подпространства. Тогда существует базис в котором возможно блочное представление оператора γ_0 (блоки имеют размер $N^2 \times N^2$)

$$\gamma_0 = \begin{pmatrix} \zeta_0 & 0 \\ 0 & \xi_0 \end{pmatrix}, \tag{6.20}$$

где ζ_0 отвечают за положительные собственные значения, а ξ_0 - за отрицательные. Определим

$$\gamma_{\tilde{\varLambda}} = \begin{pmatrix} \zeta_{\tilde{\varLambda}} & \sigma_{\tilde{\varLambda}} \\ \eta_{\tilde{\varLambda}} & \xi_{\tilde{\varLambda}} \end{pmatrix}. \tag{6.21}$$

С учетом этого определения условие (6.16) приведет к соотношениям

$$\zeta_0 \zeta_{\tilde{\varLambda}} = -\zeta_{\tilde{\varLambda}} \zeta_0, \quad \xi_0 \xi_{\tilde{\varLambda}} = -\xi_{\tilde{\varLambda}} \xi_0. \tag{6.22}$$

Это означает, что ζ_0 и ξ_0 должны иметь парные положительные и отрицательные значения (даже если $\zeta_{\tilde{\varLambda}}$ и $\xi_{\tilde{\varLambda}}$ вырождены, но не равны 0). Поэтому $\zeta_{\tilde{\varLambda}} = \xi_{\tilde{\varLambda}} \equiv 0$. Таким образом

$$\gamma_{\tilde{\varLambda}} = \begin{pmatrix} 0 & \sigma_{\tilde{\varLambda}} \\ \eta_{\tilde{\varLambda}} & 0 \end{pmatrix}, \tag{6.23}$$

и уравнение (6.1) равносильно системе

$$\begin{cases} \eta_{\tilde{\varLambda}} \sigma_{\tilde{\varPsi}} + \eta_{\tilde{\varPsi}} \sigma_{\tilde{\varLambda}} = G_{\tilde{\varLambda} \tilde{\varPsi}}, \\ \sigma_{\tilde{\varLambda}} \eta_{\tilde{\varPsi}} + \sigma_{\tilde{\varPsi}} \eta_{\tilde{\varLambda}} = G_{\tilde{\varLambda} \tilde{\varPsi}}. \end{cases} \tag{6.24}$$

Отметим, что если перейти к другому базису в пространстве $\mathbb{R}^{(2N)^2}$ с помощью невырожденного оператора S из общей линейной группы $\tilde{\gamma}_{\tilde{\varLambda}} := S \gamma_{\tilde{\varLambda}} S^{-1}$, то уравнение Клиффорда (6.18), вообще говоря, не сохранит свой вид.

Вернемся теперь к комплексной структуре. Ее квадрат равен $-E$ - эта матрица представляет некоторое ортогональное преобразование. Согласно (6.24) при n=2, N=1 такое преобразование представимо по формуле (6.12) как

$$\tilde{F}^2 = \tilde{\tilde{F}}^2 = \begin{pmatrix} -E & 0 \\ 0 & E \end{pmatrix}. \tag{6.25}$$

Если комплексная структура $f_{\tilde{\varLambda}}{}^{\tilde{\varPsi}}$ имеет в данном базисе канонический вид, то в разложении (6.15) $F := \tilde{F} = \tilde{\tilde{F}}$, $I := \tilde{I} = \tilde{\tilde{I}}$ тоже имеет блочный вид (из-за блочной структуры $\gamma_{\tilde{\varLambda}}$ и соответствия (6.8), написанного для вещественной реализации)

$$F = \begin{pmatrix} \hat{F} & 0 \\ 0 & \hat{\hat{F}}^T \end{pmatrix}, \quad \hat{F}^2 = -E, \quad \hat{\hat{F}}^2 = E, \quad I = \begin{pmatrix} iE & 0 \\ 0 & E \end{pmatrix}. \tag{6.26}$$

Что касается преобразования I, то оно определяется из соотношения

$$\begin{gathered} i\gamma_\Lambda = \tfrac{1}{2}(i(\gamma_\Lambda) + (-i)(-\gamma_\Lambda)) = \\ = \underbrace{\frac{1}{2}((1+i)E + (1-i)\tilde{F}^2)}_{:=\tilde{I}}\,\gamma_\Lambda\,\underbrace{\frac{1}{2}((1+i)E + (1-i)\tilde{\tilde{F}}^2)}_{:=\tilde{\tilde{I}}}. \end{gathered} \tag{6.27}$$

Ввиду того, что количество элементарных преобразований, составляющих комплексную структуру $f_{\tilde{\Lambda}}{}^{\tilde{\Psi}}$ всегда кратно двум (поскольку размерность вещественной реализации 2n при четном n кратна 4), то блочная структура будет именно по главной диагонали. Следовательно, для блочных составляющих $\gamma_{\tilde{\Lambda}}$ будет верно

$$f_{\tilde{\Lambda}}{}^{\tilde{\Psi}}\sigma_{\tilde{\Psi}} = \hat{F}\sigma_{\tilde{\Lambda}}\hat{\hat{F}}^T, \quad f_{\tilde{\Lambda}}{}^{\tilde{\Psi}}\eta_{\tilde{\Psi}} = \hat{\hat{F}}^T\eta_{\tilde{\Lambda}}\hat{F}, \tag{6.28}$$

что определит положительную реализацию (3.19) для верхнего знака и отрицательную реализацию (3.19) для нижнего знака. Если теперь положить ($\tilde{A}, \tilde{B}, \tilde{C}, \tilde{D}, ... = \overline{1,4}$, $\hat{A}, \hat{B}, \hat{C}, \hat{D}, ... = \overline{1,2}$, $\tilde{\tilde{A}}, \tilde{\tilde{B}}, \tilde{\tilde{C}}, \tilde{\tilde{D}}, ... = \overline{1,2}$, $A, B, C, D... = \overline{1,1}$)

$$\tilde{m}_{\pm\hat{A}}{}^{\tilde{B}} := \begin{pmatrix} \tilde{m}_{\pm A_1}{}^{\tilde{\tilde{B}}_1} & 0 \\ 0 & \tilde{\tilde{m}}_\pm{}^{A_2}{}_{\tilde{\tilde{B}}_2} \end{pmatrix}, \quad \tilde{\tilde{m}}_{\pm\hat{A}}{}^{\tilde{B}} := \begin{pmatrix} \tilde{m}_{\pm A_1}{}^{\tilde{\tilde{B}}_1} & 0 \\ 0 & \tilde{\tilde{m}}_\pm{}^{A_2}{}_{\tilde{\tilde{B}}_2} \end{pmatrix}, \tag{6.29}$$

то условие (3.19) примет вид

$$\left\{\begin{aligned} &(\tilde{m}_\pm)_A{}^{\tilde{\tilde{B}}}(\tilde{m}_\pm)^C{}_{\tilde{\tilde{B}}} = \delta_A{}^C, \\ &(\tilde{m}_\pm)_A{}^{\tilde{\tilde{B}}}(\tilde{m}_\mp)^C{}_{\tilde{\tilde{B}}} = 0, \\ &(\tilde{m}_\pm)_A{}^{\tilde{\tilde{B}}}(\tilde{m}_\pm)^A{}_{\tilde{\tilde{C}}} = \\ &\qquad = \underbrace{\frac{1}{2}(\delta_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} \pm i\hat{F}_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}})}_{:=(\tilde{\triangle}_\pm)_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}}, \end{aligned}\right. \qquad \left\{\begin{aligned} &(\tilde{\tilde{m}}_\pm)_A{}^{\tilde{\tilde{B}}}(\tilde{\tilde{m}}_\pm)^C{}_{\tilde{\tilde{B}}} = \delta_A{}^C, \\ &(\tilde{\tilde{m}}_\pm)_A{}^{\tilde{\tilde{B}}}(\tilde{\tilde{m}}_\mp)^C{}_{\tilde{\tilde{B}}} = 0, \\ &(\tilde{\tilde{m}}_\pm)_A{}^{\tilde{\tilde{B}}}(\tilde{\tilde{m}}_\pm)_\pm{}^A{}_{\tilde{\tilde{C}}} = \\ &\qquad = \underbrace{\frac{1}{2}(\delta_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} \pm \hat{\hat{F}}_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}})}_{:=(\tilde{\tilde{\triangle}}_\pm)_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}}. \end{aligned}\right. \tag{6.30}$$

Чтобы построить пошаговые операторы (6.30) для произвольного четного n, воспользуемся алгоритмом 3.1 ($\tilde{A}, \tilde{B}, \tilde{C}, \tilde{D}, ... = \overline{1, \frac{(2N)^2}{2^{I-1}}}$, $\hat{A}, \hat{B}, \hat{C}, \hat{D}, ... = \overline{1, \frac{(2N)^2}{2^I}}$, $\tilde{\tilde{A}}, \tilde{\tilde{B}}, \tilde{\tilde{C}}, \tilde{\tilde{D}}, ... = \overline{1, \frac{2N^2}{2^{I-1}}}$, $A, B, C, D... = \overline{1, \frac{2N^2}{2^I}}$, $N = 2^{n/2-1}$, $I = \overline{1, \frac{n}{2}}$)

$$\left\{\begin{aligned} &(\tilde{m}_{I_\pm})_A{}^{\tilde{\tilde{B}}}(\tilde{m}_{I_\pm})^C{}_{\tilde{\tilde{B}}} = \delta_A{}^C, \\ &(\tilde{m}_{I_\pm})_A{}^{\tilde{\tilde{B}}}(\tilde{m}_{I_\mp})^C{}_{\tilde{\tilde{B}}} = 0, \\ &(\tilde{m}_{I_\pm})_A{}^{\tilde{\tilde{B}}}(\tilde{m}_{I_\pm})^A{}_{\tilde{\tilde{C}}} = (\tilde{\triangle}_{I_\pm})_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}, \end{aligned}\right. \qquad \left\{\begin{aligned} &(\tilde{\tilde{m}}_{I_\pm})_A{}^{\tilde{\tilde{B}}}(\tilde{\tilde{m}}_{I_\pm})^C{}_{\tilde{\tilde{B}}} = \delta_A{}^C, \\ &(\tilde{\tilde{m}}_{I_\pm})_A{}^{\tilde{\tilde{B}}}(\tilde{\tilde{m}}_{I_\mp})^C{}_{\tilde{\tilde{B}}} = 0, \\ &(\tilde{\tilde{m}}_{I_\pm})_A{}^{\tilde{\tilde{B}}}(\tilde{\tilde{m}}_{I_\pm})_\pm{}^A{}_{\tilde{\tilde{C}}} = (\tilde{\tilde{\triangle}}_{I_\pm})_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}}. \end{aligned}\right. \tag{6.31}$$

При этом

$$(\tilde{\triangle}_{I_\pm})_{\tilde{\tilde{C}}}{}^{\tilde{\hat{A}}} := \frac{1}{2}(\delta_{\tilde{\tilde{C}}}{}^{\tilde{\hat{A}}} \pm \hat{I}_{\tilde{\tilde{C}}}{}^{\tilde{\hat{K}}} \hat{F}_{\tilde{\tilde{K}}}{}^{\tilde{\hat{A}}}), \quad (\tilde{\tilde{\triangle}}_{I_\pm})_{\tilde{\tilde{C}}}{}^{\tilde{\hat{A}}} := \frac{1}{2}(\delta_{\tilde{\tilde{C}}}{}^{\tilde{\hat{A}}} \pm \hat{\hat{I}}_{\tilde{\tilde{C}}}{}^{\tilde{\hat{K}}} \hat{\hat{F}}_{\tilde{\tilde{K}}}{}^{\tilde{\hat{A}}}). \tag{6.32}$$

Теперь возможно определить операторы

$$\tilde{M}_K := \tilde{m}_{\frac{n}{2}_{z_{\frac{n}{2}}}} \tilde{m}_{\frac{n}{2}-1_{z_{\frac{n}{2}-1}}} \cdot \ldots \cdot \tilde{m}_{2_{z_2}} \tilde{m}_{1_{z_1}}, \qquad \tilde{\tilde{M}}_K := \tilde{\tilde{m}}_{\frac{n}{2}_{z_{\frac{n}{2}}}} \tilde{\tilde{m}}_{\frac{n}{2}-1_{z_{\frac{n}{2}-1}}} \cdot \ldots \cdot \tilde{\tilde{m}}_{2_{z_2}} \tilde{\tilde{m}}_{1_{z_1}}, \tag{6.33}$$

где z_J $(J = \overline{1, \frac{n}{2}})$ равно 0 при выборе знака $\ll - \gg$ или 1 при выборе знака $\ll + \gg$, тогда $K = \sum\limits_{J=1}^{\frac{n}{2}} z_J \cdot 2^{J-1} + 1$. Если не важно, какой из K операторов будет выбран, то номер K будем опускать. Используя эти операторы, можно определить ($\tilde{M} := \tilde{M}_C{}^{\tilde{\hat{A}}}$, $\tilde{M}^* := \tilde{M}^{*C}{}_{\tilde{\hat{A}}}$, $\tilde{\tilde{M}} := \tilde{\tilde{M}}_C{}^{\tilde{\hat{A}}}$, $\tilde{\tilde{M}}^* := \tilde{\tilde{M}}^{*C}{}_{\tilde{\hat{A}}}$)

$$\sigma_\Lambda := \tilde{M}(m_\Lambda{}^{\tilde{A}} \sigma_{\tilde{A}}) \tilde{\tilde{M}}^T, \quad \eta_\Lambda := \tilde{\tilde{M}}^*(m_\Lambda{}^{\tilde{A}} \eta_{\tilde{A}})(\tilde{M}^*)^T. \tag{6.34}$$

Для дальнейших выкладок необходимо наложить условие

$$tr(\eta_{\tilde{\Psi}} \sigma_{\tilde{\Phi}}) = \frac{N}{2} g_{\tilde{\Psi}\tilde{\Phi}}. \tag{6.35}$$

Переходя в (6.24) к комплексной реализации с помощью операторов (6.33), можно получить

$$\begin{cases} \eta_\Lambda \sigma_\Psi + \eta_\Psi \sigma_\Lambda = g_{\Lambda\Psi}, \\ \sigma_\Lambda \eta_\Psi + \sigma_\Psi \eta_\Lambda = g_{\Lambda\Psi}, \\ tr(\eta_\Psi \sigma_\Lambda) = \frac{N}{2} g_{\Psi\Lambda}. \end{cases} \tag{6.36}$$

Это приведет согласно (6.4) к

$$R_\eta := r^\Lambda \eta_\Lambda, \quad R_\sigma := r^\Lambda \sigma_\Lambda, \quad \frac{2}{N} tr((R_\sigma)\eta_\Lambda) = r_\Lambda, \quad \frac{2}{N} tr((R_\eta)\sigma_\Lambda) = r_\Lambda, \tag{6.37}$$

$$R = \begin{pmatrix} 0 & R_\sigma \\ R_\eta & 0 \end{pmatrix}, \tag{6.38}$$

где тензоры R_η, R_σ имеют размерность уже $N \times N$. Следовательно, (6.8) разобьется на пару уравнений

$$R_\sigma \eta_\Psi R_\sigma = (r_\Psi r^\Phi - \frac{1}{2}(r_\Omega r^\Omega)\delta_\Psi{}^\Phi)\sigma_\Phi, \quad R_\eta \sigma_\Psi R_\eta = (r_\Psi r^\Phi - \frac{1}{2}(r_\Omega r^\Omega)\delta_\Psi{}^\Phi)\eta_\Phi. \tag{6.39}$$

Поскольку

$$R_\sigma R_\eta = \frac{1}{2}(r_\Omega r^\Omega) E, \tag{6.40}$$

то есть возможность определить формализацию операторов η_Λ ,σ_Λ следующим образом

$$\eta_\Lambda := \eta_\Lambda{}^{AB}, \quad \sigma_\Lambda := \sigma_{\Lambda AB} = \eta_{\Lambda BA}, \quad (A, B, \ldots = \overline{1, N}). \tag{6.41}$$

Тогда (6.11) можно переписать как

$$I).\ S_{\Psi}{}^{\Lambda}\eta_{\Lambda} = \tilde{S}^T\eta_{\Psi}\tilde{\tilde{S}},\quad II).\ S_{\Psi}{}^{\Lambda}\eta_{\Lambda} = \tilde{\tilde{S}}\sigma_{\Psi}\tilde{S}^T, \tag{6.42}$$

но уже здесь операторы $\tilde{\tilde{S}}$ и $\tilde{S}$ будут иметь размерность $N\times N$. Однако, теперь разложение этих операторов для собственного $S_{\Lambda}{}^{\Psi}$ ($\det \parallel S_{\Lambda}{}^{\Psi} \parallel = 1$) (случай I) будет иметь вид

$$\tilde{S}^T = \prod_{I=1}^{J}(R_\eta)_{2I-1}(R_\sigma)_{2I} = (R_\eta)_1(R_\sigma)_2(R_\eta)_3(R_\sigma)_4\dots(R_\eta)_{2J-1}(R_\sigma)_{2J}, \tag{6.43}$$

$$\tilde{\tilde{S}} = \prod_{I=J}^{1}(R_\sigma)_{2I}(R_\eta)_{2I-1} = (R_\sigma)_{2J}(R_\eta)_{2J-1}\dots(R_\sigma)_4(R_\eta)_3(R_\sigma)_2(R_\eta)_1, \tag{6.44}$$

а для несобственного преобразования $S_{\Lambda}{}^{\Psi}$ ($\det \parallel S_{\Lambda}{}^{\Psi} \parallel = -1$) (случай II) соответственно

$$\tilde{S}^T = (R_\eta)_1\prod_{I=1}^{J}(R_\sigma)_{2I}(R_\eta)_{2I+1} = (R_\eta)_1(R_\sigma)_2(R_\eta)_3\dots(R_\sigma)_{2J}(R_\eta)_{2J+1}, \tag{6.45}$$

$$\tilde{\tilde{S}} = (\prod_{I=J}^{1}(R_\eta)_{2I+1}(R_\sigma)_{2I})(R_\eta)_1 = (R_\eta)_{2J+1}(R_\sigma)_{2J}\dots(R_\eta)_3(R_\sigma)_2(R_\eta)_1. \tag{6.46}$$

Таким образом, уравнение (6.42) определяет алгебраическую реализацию двулистного накрытия $Spin(n,\mathbb{C})/\{\pm1\} \cong SO(n,\mathbb{C})$. Соответственно, для инволюции (6.14) определит

$$\begin{array}{ll} I).\quad S_{\Psi}{}^{\Lambda'}\eta_{\Lambda'} = \tilde{S}^T\eta_{\Psi}\tilde{\tilde{S}}, & II).\quad S_{\Psi}{}^{\Lambda'}\eta_{\Lambda'} = \tilde{\tilde{S}}^T\sigma_{\Psi}\tilde{S}, \\ \tilde{S}\bar{\tilde{S}} = \pm E,\quad \tilde{\tilde{S}}\bar{\tilde{\tilde{S}}} = \pm E. & \tilde{S} = \pm\bar{\tilde{S}},\quad \tilde{\tilde{S}} = \pm\bar{\tilde{\tilde{S}}}. \end{array} \tag{6.47}$$

Нижняя строка соотношений наследуется вещественной реализацией инволюции как в касательном, так и в спинорном пространстве. А вещественная реализация инволюции в касательном пространстве есть ортогональное преобразование, квадрат которого имеет единичную матрицу. Соответственно, в вещественной реализации спинорного пространства наследуется это свойство, и квадрат спинорных инволюций (случай I) в вещественной реализации с точностью до знака тоже является тождественным преобразованием. Осталось перейти к комплексной реализации по правилу (3.20) .

Example 6.1. *2n=2, N=1.*

Если $g_{\tilde{\Lambda}\tilde{\Psi}} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, *то*

$$\gamma_1 = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},\quad \gamma_2 = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}.$$

$$\eta_1 = \tfrac{1}{\sqrt{2}},\quad \eta_2 = -\tfrac{i}{\sqrt{2}},\quad \sigma_1 = \tfrac{1}{\sqrt{2}},\quad \sigma_2 = \tfrac{i}{\sqrt{2}}. \tag{6.48}$$

$$tr(\eta_{\tilde{\Lambda}}\sigma_{\tilde{\Psi}}) = \tfrac{1}{2}\begin{pmatrix} 1 & i \\ -i & 1 \end{pmatrix},\quad \text{- \textit{тензор вырожден.}}$$

Таким образом, тождество $tr(\eta_{\tilde\Lambda}\sigma_{\tilde\Psi}) = \frac{1}{2}g_{\tilde\Lambda\tilde\Psi}$ *неверно. Тем не менее можно воспользоваться (6.11) и (6.12). Положим*

$$\begin{gathered}(r_1)_{\tilde\Lambda} = (r_1)^{\tilde\Lambda} := \sqrt{2}(\cos\phi\ ,\ \sin\phi)\\ (S_1)_{\tilde\Lambda}{}^{\tilde\Psi} = \begin{pmatrix} 2\cos^2\phi-1 & 2\cos\phi\sin\phi \\ 2\cos\phi\sin\phi & -1+2\sin^2\phi\end{pmatrix} = \begin{pmatrix}\cos 2\phi & \sin 2\phi\\ \sin 2\phi & -\cos 2\phi\end{pmatrix} = (r_1)^{\tilde\Psi}(r_1)_{\tilde\Lambda} - \delta_{\tilde\Lambda}{}^{\tilde\Psi},\\ R_1 = (r_1)^{\tilde\Lambda}\gamma_{\tilde\Lambda} = \cos\phi\begin{pmatrix}0&1\\1&0\end{pmatrix} + \sin\phi\begin{pmatrix}0&i\\-i&0\end{pmatrix}.\end{gathered} \tag{6.49}$$

Тогда для любого вектора $x^{\tilde\Lambda} = (T, X)$ *соответствия* $X = x^{\tilde\Lambda}\gamma_{\tilde\Lambda}$, $(S_1)_{\tilde\Lambda}{}^{\tilde\Psi}x_{\tilde\Psi} \longleftrightarrow R_1XR_1$ *примут вид*

$$\begin{pmatrix}\cos 2\phi & \sin 2\phi\\ \sin 2\phi & -\cos 2\phi\end{pmatrix}\begin{pmatrix}T\\X\end{pmatrix} \longleftrightarrow \begin{pmatrix}0 & e^{i\phi}\\ e^{-i\phi} & 0\end{pmatrix}\begin{pmatrix}0 & T+iX\\ T-iX & 0\end{pmatrix}\begin{pmatrix}0 & e^{i\phi}\\ e^{-i\phi} & 0\end{pmatrix}. \tag{6.50}$$

Положим далее

$$\begin{gathered}(r_2)_{\tilde\Lambda} = (r_2)^{\tilde\Lambda} := \sqrt{2}(\cos\phi,\quad -\sin\phi).\\ (S_2)_{\tilde\Lambda}{}^{\tilde\Psi} = \begin{pmatrix}\cos 2\phi & -\sin 2\phi\\ -\sin 2\phi & -\cos 2\phi\end{pmatrix} = (r_2)^{\tilde\Psi}(r_2)_{\tilde\Lambda} - \delta_{\tilde\Lambda}{}^{\tilde\Psi},\\ R_2 = (r_2)^{\tilde\Lambda}\gamma_{\tilde\Lambda} = \cos\phi\begin{pmatrix}0&1\\1&0\end{pmatrix} - \sin\phi\begin{pmatrix}0&i\\-i&0\end{pmatrix}.\end{gathered} \tag{6.51}$$

Тогда для любого вектора $x^{\tilde\Lambda} = (T, X)$ *соответствия* $X = x^{\tilde\Lambda}\gamma_{\tilde\Lambda}$, $(S_2)_{\tilde\Lambda}{}^{\tilde\Psi}x_{\tilde\Psi} \longleftrightarrow R_2XR_2$ *примут вид*

$$\begin{pmatrix}\cos 2\phi & -\sin 2\phi\\ -\sin 2\phi & -\cos 2\phi\end{pmatrix}\begin{pmatrix}T\\X\end{pmatrix} \longleftrightarrow \begin{pmatrix}0 & e^{-i\phi}\\ e^{i\phi} & 0\end{pmatrix}\begin{pmatrix}0 & T+iX\\ T-iX & 0\end{pmatrix}\begin{pmatrix}0 & e^{-i\phi}\\ e^{i\phi} & 0\end{pmatrix}. \tag{6.52}$$

Теперь возможно построить собственное преобразование $S := S_1S_2$, *для которого* $S_{\tilde\Lambda}{}^{\tilde\Psi}x_{\tilde\Psi} \longleftrightarrow R_1R_2XR_2R_1$ *и* $\tilde S = R_1R_2$, $\tilde{\tilde S} = R_2R_1$. *Явная запись соответствия будет иметь вид*

$$\begin{pmatrix}\cos 4\phi & -\sin 4\phi\\ \sin 4\phi & \cos 4\phi\end{pmatrix}\begin{pmatrix}T\\X\end{pmatrix} \longleftrightarrow \begin{pmatrix}e^{2i\phi} & 0\\ 0 & e^{-2i\phi}\end{pmatrix}\begin{pmatrix}0 & T+iX\\ T-iX & 0\end{pmatrix}\begin{pmatrix}e^{-2i\phi} & 0\\ 0 & e^{2i\phi}\end{pmatrix}. \tag{6.53}$$

Очевидно, что SO(2) исчерпывется такими преобразованиями $S_{\tilde\Lambda}{}^{\tilde\Psi}$, *тогда* $\begin{pmatrix}e^{2i\phi} & 0\\ 0 & e^{-2i\phi}\end{pmatrix}$ *представляет группу, двулистно накрывающую группу* $\begin{pmatrix}e^{4i\phi} & 0\\ 0 & e^{-4i\phi}\end{pmatrix} \cong U(1)$. *При этом* $\tilde{\tilde S} = \overline{\tilde S}$ *(комплексносопряжены). Этим доказано, что* $SO(2) \cong U(1)$.

Example 6.2. *2n=2, N=1.*

Если $g_{\tilde\Lambda\tilde\Psi} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, *то*

$$\begin{aligned}
\gamma_1 &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma_2 = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \\
\eta_1 &= \tfrac{1}{\sqrt{2}}, \quad \eta_2 = -\tfrac{1}{\sqrt{2}}, \quad \sigma_1 = \tfrac{1}{\sqrt{2}}, \quad \sigma_2 = \tfrac{1}{\sqrt{2}}. \\
tr(\eta_{\tilde\Lambda}\sigma_{\tilde\Psi}) &= \tfrac{1}{2} \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix}, \quad \text{- тензор вырожден.}
\end{aligned} \tag{6.54}$$

Таким образом тождество $tr(\eta_{\tilde\Lambda}\sigma_{\tilde\Psi}) = \frac{1}{2}g_{ij}$ *неверно. Явная запись соответствия* $S_{\tilde\Lambda}{}^{\tilde\Psi}x_{\tilde\Psi} \longleftrightarrow \tilde{S}X\tilde{\tilde{S}}$ *будет иметь вид*

$$\begin{pmatrix} \text{ch } 4\phi & \text{sh } 4\phi \\ \text{sh } 4\phi & \text{ch } 4\phi \end{pmatrix} \begin{pmatrix} T \\ X \end{pmatrix} \longleftrightarrow \begin{pmatrix} e^{2\phi} & 0 \\ 0 & e^{-2\phi} \end{pmatrix} \begin{pmatrix} 0 & T+X \\ T-X & 0 \end{pmatrix} \begin{pmatrix} e^{-2\phi} & 0 \\ 0 & e^{2\phi} \end{pmatrix}, \tag{6.55}$$

что определит соответствие $SO^{+}(1,1) \cong R$ *(аддитивная), где* $SO^{+}(1,1)$ *- связная компонента единицы с* $S_1{}^1 > 0$.

Example 6.3. *2n=4, N=2. Если* $g_{ij} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$, *то можно построить*

$$\begin{aligned}
\eta_1 &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \eta_2 = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \\
\eta_3 &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \eta_4 = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}, \\
\sigma_1 &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma_2 = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \\
\sigma_3 &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma_4 = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.
\end{aligned} \tag{6.56}$$

Операторы γ_i *строятся согласно* $\gamma_i = \begin{pmatrix} 0 & \sigma_i \\ \eta_i & 0 \end{pmatrix}$. *Легко проверить, что* $tr(\eta_i\sigma_j) = tr(\sigma_i\eta_j) = g_{ij}$. *Вычисления в этом случае более громоздки, но аналогичны случаю n=2* *(и частично будут проведены ниже). Просто отметим тот факт, что здесь действует накрытие* $SO^+(2,2) \cong SL(2) \times SL(2)/\pm 1$. *Однако теперь хотелось бы знать, как перейти от случая n=4 к случаю n=2.*

Example 6.4. $n = 2 \subset n = 4,\ N = 2.$

$$
\begin{gathered}
m_\Lambda{}^{\tilde{\Lambda}} = \tfrac{1}{2}\begin{pmatrix} 1 & -1 & -i & i \\ i & i & 1 & 1 \end{pmatrix}, \quad m^\Lambda{}_{\tilde{\Lambda}} = \tfrac{1}{2}\begin{pmatrix} 1 & -1 & i & -i \\ -i & -i & 1 & 1 \end{pmatrix}, \\
g_{\Lambda\Psi} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad G_{\tilde{\Lambda}\tilde{\Psi}} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad f_{\tilde{\Lambda}}{}^{\tilde{\Psi}} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix},
\end{gathered}
\tag{6.57}
$$

$$
x^{\tilde{\Lambda}} = (T, Y, Z, X), \quad x^\Lambda = m^\Lambda{}_{\tilde{\Lambda}} x^{\tilde{\Lambda}} = \tfrac{1}{2}(T - Y + i(Z - X), -i(T + Y) + Z + X),
$$

тогда любое собственное ортогональное преобразование $S_{\tilde{\Lambda}}{}^{\tilde{\Psi}} \in SO^+(2,2)(\det \parallel S_\Lambda{}^\Psi \parallel = 1$ *и определитель левого верхнего минора* 2×2 *- положителен) представимо в виде конечного произведения элементарных преобразований. Поскольку* $fSf = -S$ *из-за наличия комплексной структуры, поэтому такие преобразования будут двух видов*

$$
\begin{gathered}
(S_1)_{\tilde{\Lambda}}{}^{\tilde{\Psi}} = \begin{pmatrix} \cos 2\varphi & \sin 2\varphi & 0 & 0 \\ \sin 2\varphi & -\cos 2\varphi & 0 & 0 \\ 0 & 0 & \cos 2\varphi & \sin 2\varphi \\ 0 & 0 & \sin 2\varphi & -\cos 2\varphi \end{pmatrix}, \\
(S_2)_{\tilde{\Lambda}}{}^{\tilde{\Psi}} = \begin{pmatrix} \mathrm{ch}\ 2\psi & 0 & 0 & -\mathrm{sh}\ 2\psi \\ 0 & -\mathrm{ch}\ 2\psi & -\mathrm{sh}\ 2\psi & 0 \\ 0 & \mathrm{sh}\ 2\psi & \mathrm{ch}\ 2\psi & 0 \\ \mathrm{sh}\ 2\psi & 0 & 0 & -\mathrm{ch}\ 2\psi \end{pmatrix}.
\end{gathered}
\tag{6.58}
$$

Комплексную реализацию $S_{\tilde{\Lambda}}{}^{\tilde{\Psi}}$ *можно определить как* $S_\Lambda{}^\Psi = m_\Lambda{}^{\tilde{\Lambda}} S_{\tilde{\Lambda}}{}^{\tilde{\Psi}} m^\Psi{}_{\tilde{\Psi}}$

$$
(S_1)_\Lambda{}^\Psi = \begin{pmatrix} -\sin 2\varphi & -i\cos 2\varphi \\ i\cos 2\varphi & \sin 2\varphi \end{pmatrix}, \quad (S_2)_\Lambda{}^\Psi = \begin{pmatrix} i\,\mathrm{sh}\ 2\psi & -i\,\mathrm{ch}\ 2\psi \\ i\,\mathrm{ch}\ 2\psi & -i\,\mathrm{sh}\ 2\psi \end{pmatrix}. \tag{6.59}
$$

Поэтому $S = S_1 S_2$ *описывает группу* $SO(2, \mathbb{C})$ *полностью. Пусть теперь*

$$
\begin{aligned}
(r_1)^{\tilde{\Lambda}} &= \sqrt{2}(\cos\phi, \sin\phi, 0, 0) \quad , \quad (r_2)^{\tilde{\Lambda}} = \sqrt{2}(0, 0, \cos\phi, \sin\phi), \\
(r_1)_{\tilde{\Lambda}} &= \sqrt{2}(\cos\phi, \sin\phi, 0, 0) \quad , \quad (r_2)_{\tilde{\Lambda}} = \sqrt{2}(0, 0, -\cos\phi, -\sin\phi).
\end{aligned}
\tag{6.60}
$$

Построим $(S_1)_{\tilde\Lambda}{}^{\tilde\Theta} = ((r_1)_{\tilde\Lambda}(r_1)^{\tilde\Psi} - \delta_{\tilde\Lambda}{}^{\tilde\Psi})((r_2)_{\tilde\Psi}(r_2)^{\tilde\Theta} + \delta_{\tilde\Psi}{}^{\tilde\Theta})$. *При этом* $(r_1)_{\tilde\Lambda}(r_1)^{\tilde\Lambda} = -(r_2)_{\tilde\Lambda}(r_2)^{\tilde\Lambda} = 2$. *Пусть* $\eta_{\tilde\Lambda}$ *и* $\sigma_{\tilde\Lambda}$ *такие же как в предыдущем примере, то*

$$R_1 = (r_1)^{\tilde\Lambda}\eta_{\tilde\Lambda} = \begin{pmatrix} \cos\phi & \sin\phi \\ -\sin\phi & \cos\phi \end{pmatrix} \quad , \quad R_2 = (r_2)^{\tilde\Lambda}\sigma_{\tilde\Lambda} = \begin{pmatrix} -\cos\phi & \sin\phi \\ \sin\phi & \cos\phi \end{pmatrix},$$

$$R_1R_2 = \begin{pmatrix} -\cos 2\phi & \sin 2\phi \\ \sin 2\phi & \cos 2\phi \end{pmatrix} \quad , \qquad R_2R_1 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \tag{6.61}$$

$$(r_3)^{\tilde\Lambda} = \sqrt{2}(\mathrm{ch}\ \psi, 0, 0, -\mathrm{sh}\ \psi) \quad , \quad (r_4)^{\tilde\Lambda} = \sqrt{2}(0, -\mathrm{sh}\ \psi, -\mathrm{ch}\ \psi, 0),$$

$$(r_3)_{\tilde\Lambda} = \sqrt{2}(\mathrm{ch}\ \psi, 0, 0, \mathrm{sh}\ \psi) \quad , \quad (r_4)_{\tilde\Lambda} = \sqrt{2}(0, -\mathrm{sh}\ \psi, \mathrm{ch}\ \psi, 0).$$

Построим $(S_2)_{\tilde\Lambda}{}^{\tilde\Theta} = ((r_3)_{\tilde\Lambda}(r_3)^{\tilde\Psi} - \delta_{\tilde\Lambda}{}^{\tilde\Psi})((r_4)_{\tilde\Psi}(r_4)^{\tilde\Theta} + \delta_{\tilde\Psi}{}^{\tilde\Theta})$. *При этом* $(r_3)_{\tilde\Lambda}(r_3)^{\tilde\Lambda} = -(r_4)_{\tilde\Lambda}(r_4)^{\tilde\Lambda} = 2$.

$$R_3 = (r_3)^{\tilde\Lambda}\eta_{\tilde\Lambda} = \begin{pmatrix} \mathrm{ch}\ \psi & \mathrm{sh}\ \psi \\ \mathrm{sh}\ \psi & \mathrm{ch}\ \psi \end{pmatrix} \quad , \quad R_4 = (r_4)^{\tilde\Lambda}\sigma_{\tilde\Lambda} = \begin{pmatrix} \mathrm{ch}\ \psi & \mathrm{sh}\ \psi \\ -\mathrm{sh}\ \psi & -\mathrm{ch}\ \psi \end{pmatrix},$$

$$\begin{gathered} R_3R_4 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad , \qquad R_4R_3 = \begin{pmatrix} \mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ -\mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix}, \\ \tilde S^T = R_1R_2R_3R_4 = \begin{pmatrix} -\cos 2\phi & -\sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix}, \\ \tilde{\tilde S} = R_4R_3R_2R_1 = \begin{pmatrix} -\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ \mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix}. \end{gathered} \tag{6.62}$$

Поскольку $x^{\tilde\Lambda} = (T, Y, Z, X)$, $\quad x^\Lambda = \frac{1}{2}(T - Y + i(Z - X), -i(T + Y) + Z + X)$, *то*

$$\begin{gathered} x^\Lambda S_\Lambda{}^\Psi \longleftrightarrow \tilde S^T X_\eta \tilde{\tilde S} = \\ = \begin{pmatrix} -\cos 2\phi & -\sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix} \begin{pmatrix} T + Z & -X + Y \\ -X - Y & T - Z \end{pmatrix} \begin{pmatrix} -\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ \mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix}. \end{gathered} \tag{6.63}$$

Таким образом, $SO(2, \mathbb{C}) \cong U(1) \times R(additive)$, *где* $\begin{pmatrix} -\cos 2\phi & -\sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix}$ *представляет группу U(1), а* $\begin{pmatrix} -\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ \mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix}$ *представляет аддитивную группу R. Если в соответствующих уравнениях сделать замену* η *на* σ, *то*

$$\begin{gathered} x^\Lambda S_\Lambda{}^\Psi \longleftrightarrow \tilde{\tilde S}'^T X_\sigma \tilde S' = \\ = \begin{pmatrix} -\mathrm{ch}\ 2\psi & -\mathrm{sh}\ 2\psi \\ -\mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix} \begin{pmatrix} T - Z & X - Y \\ X + Y & T + Z \end{pmatrix} \begin{pmatrix} -\cos 2\phi & \sin 2\phi \\ -\sin 2\phi & \cos 2\phi \end{pmatrix}. \end{gathered} \tag{6.64}$$

Example 6.5. $n = 2 \subset 2n = 4,\ N = 2.$

Пусть тензор инволюции в касательном слое для вещественной реализации V_4 *представляется тензором (слева вложение для метрики индекса 2, справа - для индекса 1)*

$$S_{\tilde\Lambda}{}^{\tilde\Psi} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad S_{\tilde\Lambda}{}^{\tilde\Psi} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \tag{6.65}$$

$$G_{\tilde\Lambda\tilde\Psi} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix},$$

и выполнены все тождества предыдущих примеров. Тогда для комплексной реализации $\mathbb{C}V_2$ *он будет иметь вид* $S_\Lambda{}^{\Psi'} = m_\Lambda{}^{\tilde\Lambda} S_{\tilde\Lambda}{}^{\tilde\Psi} \bar m^{\Lambda'}{}_{\tilde\Psi}$

$$S_\Lambda{}^{\Psi'} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad S_\Lambda{}^{\Psi'} = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \tag{6.66}$$

С другой стороны, у инволюции существует образ (буковка T *означает транспонирование) и в некотором базисе* $S_{\tilde\Lambda}{}^{\tilde\Psi}\eta_{\tilde\Psi} = \tilde S^T \eta_{\tilde\Psi} \tilde S$

$$\tilde S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \tilde S = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \tag{6.67}$$

В первом случае получается (индекс 2)

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} -\cos 2\phi & -\sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} -\cos 2\phi & -\sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix},$$
$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} -\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi \\ \mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} -\mathrm{ch}\ 2\psi & -\mathrm{sh}\ 2\psi \\ -\mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi \end{pmatrix},$$
$$H_i{}^\Lambda = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -i \\ -1 & -i \end{pmatrix}, \quad H^i{}_\Lambda = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & i \\ -1 & i \end{pmatrix}, \quad g_{ij} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad g_{\Lambda\Psi} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \tag{6.68}$$

Поэтому $\psi = 0$

$$(S_1)_i{}^j = \begin{pmatrix} \cos 2\phi & \sin 2\phi \\ \sin 2\phi & -\cos 2\phi \end{pmatrix}, \quad (S_2)_i{}^j = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \tag{6.69}$$

и мы опять приходим к изоморфизму $SO(2) \cong U(1)$

$$\begin{gathered}\begin{pmatrix} T, & Y \end{pmatrix}\begin{pmatrix}\cos 2\phi & \sin 2\phi\\ -\sin 2\phi & \cos 2\phi\end{pmatrix}\longleftrightarrow\\ \longleftrightarrow \left(\cos 2\phi \begin{pmatrix}1 & 0\\ 0 & 1\end{pmatrix} + \sin 2\phi \begin{pmatrix}0 & 1\\ -1 & 0\end{pmatrix}\right)\begin{pmatrix}T & Y\\ -Y & T\end{pmatrix}.\end{gathered}\tag{6.70}$$

Во втором случае (индекс 1)

$$\begin{gathered}\begin{pmatrix}0 & 1\\ 1 & 0\end{pmatrix}\begin{pmatrix}-\cos 2\phi & -\sin 2\phi\\ \sin 2\phi & -\cos 2\phi\end{pmatrix}\begin{pmatrix}0 & 1\\ 1 & 0\end{pmatrix} = \begin{pmatrix}-\cos 2\phi & \sin 2\phi\\ -\sin 2\phi & -\cos 2\phi\end{pmatrix},\\ \begin{pmatrix}0 & 1\\ 1 & 0\end{pmatrix}\begin{pmatrix}-\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi\\ \mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi\end{pmatrix}\begin{pmatrix}0 & 1\\ 1 & 0\end{pmatrix} = \begin{pmatrix}-\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi\\ \mathrm{sh}\ 2\psi & -\mathrm{ch}\ 2\psi\end{pmatrix},\\ H_i{}^\Lambda = \tfrac{1}{\sqrt{2}}\begin{pmatrix}1 & i\\ i & 1\end{pmatrix}, \quad H^i{}_\Lambda = \tfrac{1}{\sqrt{2}}\begin{pmatrix}1 & -i\\ -i & 1\end{pmatrix},\\ g_{ij} = \begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}, \quad g_{\Lambda\Psi} = \begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}.\end{gathered}\tag{6.71}$$

Поэтому $2\phi = 0, \pi$

$$(S_1)_i{}^j = \pm\begin{pmatrix}-1 & 0\\ 0 & 1\end{pmatrix}, \quad (S_2)_i{}^j = \begin{pmatrix}-\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi\\ -\mathrm{sh}\ 2\psi & \mathrm{ch}\ 2\psi\end{pmatrix},\tag{6.72}$$

и мы опять приходим к изоморфизму $SO^+(1,1) \cong R$ *(аддитивная)*

$$\begin{gathered}\begin{pmatrix} T, & X \end{pmatrix}\begin{pmatrix}\mathrm{ch}\ 2\psi & \mathrm{sh}\ 2\psi\\ \mathrm{sh}\ 2\phi & \mathrm{ch}\ 2\phi\end{pmatrix}\longleftrightarrow\\ \longleftrightarrow \begin{pmatrix}T & -X\\ -X & T\end{pmatrix}\left(\mathrm{ch}\ 2\psi\begin{pmatrix}1 & 0\\ 0 & 1\end{pmatrix} + \mathrm{sh}\ 2\psi\begin{pmatrix}0 & -1\\ -1 & 0\end{pmatrix}\right).\end{gathered}\tag{6.73}$$

Следующий пример достоин отдельного параграфа.

7 Кватернионы

В этом параграфе рассказывается о том, как связана алгебра кватернионов с клиффордовой алгеброй и о том, как построить в явном виде двулистные накрытия, соответствующие этому случаю. Вывод всех результатов этого параграфа сделан на основе [18].

Известно, что алгебру Клиффорда $CL(g^n)$ можно представить как

$$\begin{aligned} CL(g^n) &: AI + B^{\Lambda}e_{\Lambda} + C^{\Lambda\Psi}e_{\Lambda}e_{\Psi} + D^{\Lambda\Psi\Phi}e_{\Lambda}e_{\Psi}e_{\Phi} + ... = \\ &= \begin{cases} CL^0(g^n)(\text{четная}) : AI + C^{\Lambda\Psi}e_{\Lambda}e_{\Psi} + ..., \\ CL^1(g^n)(\text{нечетная}) : B^{\Lambda}e_{\Lambda} + D^{\Lambda\Psi\Phi}e_{\Lambda}e_{\Psi}e_{\Phi} + ..., \end{cases} \end{aligned} \tag{7.1}$$

что и называется Z_2 градуировкой. При этом $CL^0(g^n)$ есть подалгебра $CL(g^n)$. Если рассматривать n=4 с 8-мерной алгеброй Клиффорда $CL^0(g^n)$, то базис указанной алгебры будет иметь вид

$$1, \;\; e_1e_2, \;\; e_1e_3, \;\; e_2e_3, \;\; e_1e_4, \;\; e_2e_4, \;\; e_3e_4, \;\; e_1e_2e_3e_4. \tag{7.2}$$

Из уравнения Клиффорда (2.8) будет следовать

$$e_{\Lambda}e_{\Psi} = 1, \;\; \Lambda = \Psi, \;\; e_{\Lambda}e_{\Psi} = -e_{\Psi}e_{\Lambda}, \;\; \Lambda \neq \Psi \tag{7.3}$$

в некотором базисе

$$\begin{aligned} &I = 1, && i := e_1e_2, \\ &j =: e_1e_3, && k := e_2e_3, \\ &e := e_3e_4, && f := ie = e_1e_2e_3e_4, \\ &g := je = e_1e_3e_3e_4 = e_1e_4, && h := ke = e_2e_3e_3e_4 = e_2e_4, \end{aligned} \tag{7.4}$$

который будет представлять собой базис, подобный базису алгебры октав, но с другой нормировкой. Для этого необходимо рассмотреть вложение $\mathbb{R}^4 \subset \mathbb{C}^4$. Затем можно найти представление указанного базиса с помощью построенных выше операторов $\gamma_i = H_i{}^{\Lambda}\gamma_{\Lambda}$. Но такой базис, в свою очередь, получается прямой суммой двух кватернионных базисов, за что отвечает изоморфизм $CL^0(g^n_{(n,0)}) \cong \mathbb{H} \oplus \mathbb{H}$. Для того, чтобы построить соответствующее представление указанного базиса, необходимо потребовать для частей операторов γ_i выполнение условия $tr(\eta_i\sigma_j) = \frac{N}{2}g_{ij}$. И тогда, действительно, возможно ограничиться кватернионным базисом (как раз имеющим 2 генератора i и j (k=ij) и кватернионную единицу I)

$$\begin{gathered} \tfrac{1}{\sqrt{2}}I := \eta_1 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \;\; \tfrac{1}{\sqrt{2}}i := \eta_2 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \\ \tfrac{1}{\sqrt{2}}j := \eta_3 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \;\; \tfrac{1}{\sqrt{2}}k := \eta_4 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \\ \sigma_1 = \tfrac{1}{2}\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \;\; \sigma_2 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix}, \\ \sigma_3 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \;\; \sigma_4 = \tfrac{1}{\sqrt{2}}\begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}. \end{gathered} \tag{7.5}$$

Пусть теперь на пространстве $\mathbb{R}^4 \subset \mathbb{C}^4$ заданы генерирующие элементарные ортогональные преобразования векторы

$$\begin{gathered}
r_1 = (\cos\phi_1, \sin\phi_1, 0, 0), \quad r_2 = (\cos\phi_2, 0, \sin\phi_2, 0), \quad r_3 = (0, \cos\phi_3, \sin\phi_3, 0), \\
r_4 = (\cos\phi_4, 0, 0, \sin\phi_4), \quad r_5 = (0, \cos\phi_5, 0, \sin\phi_5), \quad r_6 = (0, 0, \cos\phi_6, \sin\phi_6).
\end{gathered} \tag{7.6}$$

При этом, как и прежде, преобразования $(r_I)^i (r_I)_j - \delta_j{}^i$ $(I = \overline{1,6})$ генерируют все вращения из группы $\mathrm{O}(4)$. Соответственно, на пространстве $\mathbb{C}^2$ $(N = 2^{2/2-1} = 2)$ будут индуцированы их образы

$$\begin{gathered}
R_1 := r_1{}^i \eta_i = \begin{pmatrix} \cos\phi_1 & i\sin\phi_1 \\ i\sin\phi_1 & \cos\phi_1 \end{pmatrix}, \quad R_2 := r_2{}^i \sigma_i = \begin{pmatrix} \cos\phi_2 & \sin\phi_2 \\ -\sin\phi_2 & \cos\phi_2 \end{pmatrix}, \\
R_3 := r_3{}^i \eta_i = \begin{pmatrix} 0 & i\cos\phi_3 - \sin\phi_3 \\ i\cos\phi_3 + \sin\phi_3 & 0 \end{pmatrix}, \\
R_4 := r_4{}^i \sigma_i = \begin{pmatrix} \cos\phi_4 - i\sin\phi_4 & 0 \\ 0 & \cos\phi_4 + i\sin\phi_4 \end{pmatrix}, \\
R_5 := r_5{}^i \eta_i = \begin{pmatrix} i\sin\phi_5 & i\cos\phi_5 \\ i\cos\phi_5 & -i\sin\phi_5 \end{pmatrix}, \quad R_6 := r_6{}^i \sigma_i = \begin{pmatrix} -i\sin\phi_6 & \cos\phi_6 \\ -\cos\phi_6 & i\sin\phi_6 \end{pmatrix}.
\end{gathered} \tag{7.7}$$

Чтобы упростить выкладки, ограничимся случаем, когда $\phi_1 = \phi_2 = \phi_4 = \frac{\pi}{2}$. Тогда

$$\begin{gathered}
R_1 R_2 R_3 R_4 R_5 R_6 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 0 & ie^{i\phi_3} \\ ie^{-i\phi_3} & 0 \end{pmatrix} \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} \\
\begin{pmatrix} i\sin\phi_5 & i\cos\phi_5 \\ i\cos\phi_5 & -i\sin\phi_5 \end{pmatrix} \begin{pmatrix} -i\sin\phi_6 & \cos\phi_6 \\ -\cos\phi_6 & i\sin\phi_6 \end{pmatrix} = \begin{pmatrix} 0 & ie^{i\phi_3} \\ ie^{-i\phi_3} & 0 \end{pmatrix} \\
\begin{pmatrix} \overbrace{\sin\phi_5\sin\phi_6 - i\cos\phi_5\cos\phi_6}^{=:\bar{a}} & \overbrace{i\sin\phi_5\cos\phi_6 - \cos\phi_5\sin\phi_6}^{=:-\bar{b}} \\ \underbrace{i\sin\phi_5\cos\phi_6 + \cos\phi_5\sin\phi_6}_{=:b} & \underbrace{\sin\phi_5\sin\phi_6 + i\cos\phi_5\cos\phi_6}_{=:a} \end{pmatrix} = \\
= \begin{pmatrix} ie^{i\phi_3} b & ie^{i\phi_3} a \\ ie^{-i\phi_3} \bar{a} & -ie^{-i\phi_3} \bar{b} \end{pmatrix} =: A_1.
\end{gathered} \tag{7.8}$$

$$R_6R_5R_4R_3R_2R_1 = \begin{pmatrix} -i\sin\phi_6 & \cos\phi_6 \\ -\cos\phi_6 & i\sin\phi_6 \end{pmatrix} \begin{pmatrix} i\sin\phi_5 & i\cos\phi_5 \\ i\cos\phi_5 & -i\sin\phi_5 \end{pmatrix}$$
$$\begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} \begin{pmatrix} 0 & ie^{i\phi_3} \\ ie^{-i\phi_3} & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} =$$
$$= \begin{pmatrix} \overbrace{\sin\phi_5\sin\phi_6 + i\cos\phi_5\cos\phi_6}^{=:a} & \overbrace{-i\sin\phi_5\cos\phi_6 + \cos\phi_5\sin\phi_6}^{=:\bar{b}} \\ \underbrace{-i\sin\phi_5\cos\phi_6 - \cos\phi_5\sin\phi_6}_{=:-b} & \underbrace{\sin\phi_5\sin\phi_6 - i\cos\phi_5\cos\phi_6}_{=:\bar{a}} \end{pmatrix} \tag{7.9}$$
$$\begin{pmatrix} 0 & -ie^{i\phi_3} \\ -ie^{-i\phi_3} & 0 \end{pmatrix} = \begin{pmatrix} -ie^{-i\phi_3}\bar{b} & -ie^{i\phi_3}a \\ -ie^{-i\phi_3}\bar{a} & ie^{-i\phi_3}b \end{pmatrix} =: \bar{A_1}^T = A_1{}^*.$$

Этим построено двулистное накрытие $SO(3) \cong SU(2)/\{\pm 1\}$ и поэтому $Spin(3) \cong SU(2)$

$$S_{1i}{}^j\eta_j = A_1{}^T\eta_i A_1^*. \tag{7.10}$$

Если теперь положить $\phi_3 = \phi_5 = \phi_6 = \frac{\pi}{2}$ и провести аналогичные выкладки, то получим

$$R_1R_2R_3R_4R_5R_6 =$$
$$\begin{pmatrix} \cos\phi_1 & i\sin\phi_1 \\ i\sin\phi_1 & \cos\phi_1 \end{pmatrix} \begin{pmatrix} \cos\phi_2 & \sin\phi_2 \\ -\sin\phi_2 & \cos\phi_2 \end{pmatrix}$$
$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} e^{-i\phi_4} & \\ 0 & e^{i\phi_4} \end{pmatrix} \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} =$$
$$= \begin{pmatrix} \overbrace{\cos\phi_1\cos\phi_2 - i\sin\phi_1\sin\phi_2}^{:=\bar{a}} & \overbrace{\cos\phi_1\sin\phi_2 + i\sin\phi_1\cos\phi_2}^{=:b} \\ \underbrace{i\sin\phi_1\cos\phi_2 - \cos\phi_1\sin\phi_2}_{:=-\bar{b}} & \underbrace{i\sin\phi_1\sin\phi_1 + \cos\phi_1\cos\phi_2}_{=:a} \end{pmatrix} \tag{7.11}$$
$$\begin{pmatrix} 0 & -e^{i\phi_4} \\ e^{-i\phi_4} & 0 \end{pmatrix} = \begin{pmatrix} e^{-i\phi_4}b & -e^{i\phi_4}\bar{a} \\ e^{-i\phi_4}a & e^{i\phi_4}\bar{b} \end{pmatrix} =: A_2{}^T.$$

$$R_6R_5R_4R_3R_2R_1 =$$
$$\begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \begin{pmatrix} e^{-i\phi_4} & 0 \\ 0 & e^{i\phi_4} \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$
$$\begin{pmatrix} \cos\phi_2 & \sin\phi_2 \\ -\sin\phi_2 & \cos\phi_2 \end{pmatrix} \begin{pmatrix} \cos\phi_1 & i\sin\phi_1 \\ i\sin\phi_1 & \cos\phi_1 \end{pmatrix} = \begin{pmatrix} 0 & -e^{-i\phi_4} \\ e^{i\phi_4} & 0 \end{pmatrix}$$
$$\begin{pmatrix} \overbrace{\cos\phi_1\cos\phi_2 + i\sin\phi_1\sin\phi_2}^{:=a} & \overbrace{\cos\phi_1\sin\phi_2 + i\sin\phi_1\cos\phi_2}^{:=b} \\ \underbrace{i\sin\phi_1\cos\phi_2 - \cos\phi_1\sin\phi_2}_{:=-\bar{b}} & \underbrace{-i\sin\phi_1\sin\phi_1 + \cos\phi_1\cos\phi_2}_{=:\bar{a}} \end{pmatrix} = \tag{7.12}$$
$$= \begin{pmatrix} e^{-i\phi_4}\bar{b} & -e^{-i\phi_4}\bar{a} \\ e^{i\phi_4}a & e^{i\phi_4}b \end{pmatrix} =: A_2{}^\sharp.$$

Поэтому

$$S_{2i}{}^{j}\eta_j = A_2{}^T\eta_i A_2^\sharp. \tag{7.13}$$

Преобразования A_1, A_2 - представители группы SU(2), которую полностью и описывают. Собирая два уравнения вместе, получим, что

$$S_i{}^j\eta_j := (S_1S_2)_i{}^j\eta_j = A_1{}^T A_2{}^T \eta_j A_2^\sharp A_1^* := \tilde{S}^T \eta_j \tilde{\tilde{S}}. \tag{7.14}$$

Вообще говоря, зная преобразование $A_1{}^T A_2{}^T$, ничего нельзя сказать про $A_2^\sharp A_1^*$. Поэтому преобразования $\tilde{S}^T$ и $\tilde{\tilde{S}}$ суть различны. Это означает, что существует двулистное накрытие $SO(4) \cong SU(2) \times SU(2)/\{\pm 1\}$. И поэтому $Spin(4) \cong SU(2) \times SU(2)$.

8 Частные решения уравнения Клиффорда

В этом параграфе рассказывается о том, как построить частные решения уравнения Клиффорда, которые позволят однозначно построить двулистное накрытие $Spin(n, \mathbb{C})/\{\pm 1\} \cong SO(n, \mathbb{C})$. Такие решения дадут возможность в дальнейшем однозначно продолжить риманову связность без кручения в спинорное расслоение. Вывод всех результатов этого параграфа сделан на основе предыдущих выкладок.

Для того, чтобы найти некоторые частные решения уравнения Клиффорда (6.36), перепишем его как

$$\eta_\Lambda{}^{AB}\eta_{\Psi CB} + \eta_\Psi{}^{AB}\eta_{\Lambda CB} = g_{\Lambda\Psi}\delta_C{}^A. \tag{8.1}$$

Example 8.1. *n=4, N=2.*

В этом случае воспользуемся изоморфизмом $\mathbb{C}^4 \cong \mathbb{C}(2)$, который строится с помощью связующих операторов $\eta^\Lambda{}_{AB}$ (6.37)

$$r^\Lambda = \frac{1}{2}\eta^\Lambda{}_{AB}R^{AB}. \tag{8.2}$$

Пусть метрика в $\mathbb{C}^4$ имеет вид

$$g_{\Lambda\Psi} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \tag{8.3}$$

Очевидно, что любой тензор R^{AB} разлагается на 3 симметричных и 1 кососимметричную части

$$\begin{aligned}
R^{AB} &= R_1 \underbrace{\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}}_{(\varepsilon_1)^{AB}} + R_2 \underbrace{\begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix}}_{(\varepsilon_2)^{AB}} + R_3 \underbrace{\begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}}_{(\varepsilon_3)^{AB}} + R_{\tilde{1}} \underbrace{\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}}_{(\varepsilon_{\tilde{1}})^{AB}}, \\
R_{AB} &= R_1 \underbrace{\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}}_{(\varepsilon_1)_{AB}} + R_2 \underbrace{\begin{pmatrix} -i & 0 \\ 0 & -i \end{pmatrix}}_{(\varepsilon_2)_{AB}} + R_3 \underbrace{\begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}}_{(\varepsilon_3)_{AB}} + R_{\tilde{1}} \underbrace{\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}}_{(\varepsilon_{\tilde{1}})_{AB}}
\end{aligned} \tag{8.4}$$

с метрическим спин-тензором $\varepsilon_{AB} = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$.

Example 8.2. *n=6, N=4.*

Построим теперь $(A, B, ... = \overline{1, N},\ a, b, ... = \overline{1, N/2},\ I = \overline{1, 2, 3, \tilde{1}})$

$$\begin{gathered}
(\varepsilon_I)^{AB} = \begin{pmatrix} (\varepsilon_I)^{ab} & 0 \\ 0 & -(\varepsilon_I)^T_{ab} \end{pmatrix}, \\
(\varepsilon_4)^{AB} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad (\varepsilon_{\tilde{2}})_{AB} = \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \\ -i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{pmatrix},
\end{gathered} \tag{8.5}$$

где $(\varepsilon_I)^{ab}$ *берутся из предыдущего примера, там как раз n=4, N=2. Это один из возможных вариантов (4 симметричные и 2 кососимметрические части). Если теперь домножить справа все полученные* $(\varepsilon_I)^{AB}$ $(I = \overline{1, 2, 3, 4, \tilde{1}, \tilde{2}})$ *на*

$$\begin{pmatrix} 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \end{pmatrix}, \tag{8.6}$$

то возможно построить новые тензоры (6 кососимметричных частей)

$$\begin{gathered}
(\varepsilon_{\tilde{1}})^{AB} = \begin{pmatrix} 0 & 0 & 0 & i \\ 0 & 0 & i & 0 \\ 0 & -i & 0 & 0 \\ -i & 0 & 0 & 0 \end{pmatrix}, \quad (\varepsilon_{\tilde{2}})^{AB} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \\
(\varepsilon_{\tilde{3}})^{AB} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}, \quad (\varepsilon_{\tilde{4}})^{AB} = \begin{pmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & i \\ i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{pmatrix}, \\
(\varepsilon_{\tilde{5}})^{AB} = \begin{pmatrix} 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \end{pmatrix}, \quad (\varepsilon_{\tilde{6}})^{AB} = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix},
\end{gathered} \tag{8.7}$$

что даст возможность определить

$$R^{AB} = -R^{BA} = R_{\tilde{1}}(\varepsilon_{\tilde{1}})^{AB} + R_{\tilde{2}}(\varepsilon_{\tilde{2}})^{AB} + R_{\tilde{3}}(\varepsilon_{\tilde{3}})^{AB} + R_{\tilde{4}}(\varepsilon_{\tilde{4}}^{AB}) + R_{\tilde{5}}(\varepsilon_{\tilde{5}})^{AB} + R_{\tilde{6}}(\varepsilon_{\tilde{6}}^{AB}). \tag{8.8}$$

Для того, чтобы получить подобную конструкцию для любого четного n, дадим следующие определения.

Определение 8.1. *Пусть в пространстве $\mathbb{C}^n$ $(n \geq 4)$ определено n ортогональных векторов вида $(\eta_I)^\Lambda$*

$$(\eta_I)^\Lambda (\eta_J)_\Lambda = 2\delta_{IJ}, \quad \Lambda,\ \Psi,\ ... = \overline{1, n}, \tag{8.9}$$

которым в спинорном пространстве поставим в соответствие спин-тензоры

$$(\varepsilon_I)_{AB} := (\eta_I)^\Lambda \eta_{\Lambda AB}. \tag{8.10}$$

Тогда среди n векторов $(\eta_I)^\Lambda$ найдутся $\frac{n}{2} + q$ векторов таких, что

$$\eta^\Psi{}_{(AB)} = \frac{1}{2} \sum_{Q=1}^{\frac{n}{2}+q} (\eta_Q)^\Psi (\varepsilon_Q)_{AB}, \quad (\varepsilon_Q)_{AB} = (\varepsilon_Q)_{BA} \tag{8.11}$$

и $\frac{n}{2} + \tilde{q}$ векторов таких, что

$$\eta^\Psi{}_{[AB]} = \frac{1}{2} \sum_{\tilde{Q}=1}^{\frac{n}{2}+\tilde{q}} (\eta_{\tilde{Q}})^\Psi (\varepsilon_{\tilde{Q}})_{AB}, \quad (\varepsilon_{\tilde{Q}})_{AB} = -(\varepsilon_{\tilde{Q}})_{BA}, \quad q + \tilde{q} = 0. \tag{8.12}$$

Кроме того, потребуем, чтобы вещественная реализация таких операторов удовлетворяла аналогичным соотношениям.

Следствие 8.1. *Из соотношения*

$$\tilde{T}_A{}^B(\varepsilon_I)_{BC} + \tilde{\tilde{T}}_C{}^B(\varepsilon_I)_{AB} = 0, \tag{8.13}$$

выполненного для всех $I = \overline{1,n}$*, следует, что* $\tilde{T}_A{}^B = \tilde{\tilde{T}}_A{}^B = 0$.

Доказательство. Воспользуемся тождеством

$$\begin{gathered}\eta_{[\Gamma}{}^{CN}\eta_{\Omega]MN}\varepsilon_{CD}{}^{ML} = \tfrac{1}{2}\sum_I \eta_{[\Gamma}{}^{CN}\eta_{\Omega]MN}(\varepsilon_I)_{CD}(\varepsilon_I)^{ML} = \\ = \tfrac{1}{2}\sum_I(-\eta_{[\Gamma|CD}(\varepsilon_{I|})^{CN} + (\eta_I)_{[\Gamma}\delta_{|D|}{}^N)\eta_{\Omega]MN}(\varepsilon_I)^{ML} = \\ = \tfrac{1}{2}\sum_I(-\eta_{[\Gamma|CD|})(-\eta_{\Omega]}{}^{CL} + (\eta_{|I|})_{\Omega]}(\varepsilon_I)^{CL}) + (\eta_I)_{[\Gamma}\eta_{\Omega]MN}(\varepsilon_I)^{ML} = \tfrac{n-4}{2}\eta_{[\Gamma|CD|}\eta_{\Omega]}{}^{CL}.\end{gathered} \tag{8.14}$$

Тогда

$$\tilde{T}_A{}^B(\varepsilon_I)_{BC} + \tilde{\tilde{T}}_C{}^B(\varepsilon_I)_{AB} = 0,$$

$$\begin{cases} \tfrac{n}{2}\tilde{T}_A{}^D + \tilde{\tilde{T}}_C{}^B\varepsilon_{AB}{}^{DC} = 0, \\ \tfrac{n}{2}\tilde{\tilde{T}}_A{}^D + \tilde{T}_C{}^B\varepsilon_{AB}{}^{DC} = 0, \end{cases} \quad \Rightarrow \tilde{\tilde{T}}_A{}^A = \tilde{T}_A{}^A = 0,$$

$$\begin{cases} \tfrac{n}{2}\tilde{T}_A{}^D\eta_{\tilde{\Lambda}}{}^{AK}\eta_{\tilde{\Psi}DK} - \tfrac{n-4}{2}\tilde{\tilde{T}}_A{}^D\eta_{\tilde{\Lambda}}{}^{KA}\eta_{\tilde{\Psi}KD} = 0, \\ \tfrac{n}{2}\tilde{\tilde{T}}_A{}^D\eta_{\tilde{\Lambda}}{}^{AK}\eta_{\tilde{\Psi}DK} - \tfrac{n-4}{2}\tilde{T}_A{}^D\eta_{\tilde{\Lambda}}{}^{KA}\eta_{\tilde{\Psi}KD} = 0, \end{cases}$$

$$\begin{cases} \tfrac{n}{2}\tilde{T}_A{}^D\varepsilon_{PL}{}^{AK}\varepsilon^{QL}{}_{DK} - \tfrac{n-4}{2}\tilde{\tilde{T}}_A{}^D\varepsilon_{PL}{}^{KA}\varepsilon^{QL}{}_{KD} = 0, \\ \tfrac{n}{2}\tilde{\tilde{T}}_A{}^D\varepsilon_{LP}{}^{AK}\varepsilon^{LQ}{}_{DK} - \tfrac{n-4}{2}\tilde{T}_A{}^D\varepsilon_{LP}{}^{KA}\varepsilon^{LQ}{}_{KD} = 0, \end{cases} \Rightarrow \begin{cases} \tilde{T}_A{}^D\varepsilon_{PL}{}^{AK}\varepsilon^{QL}{}_{DK} = 0, \\ \tilde{\tilde{T}}_A{}^D\varepsilon_{PL}{}^{AK}\varepsilon^{QL}{}_{DK} = 0. \end{cases} \tag{8.15}$$

Следовательно, $\tilde{\tilde{T}}_A{}^D = \tilde{T}_A{}^D = 0$. □

Таким образом, на образующие тензоры $(\varepsilon_I)_{AB}$ уравнением Клиффорда (8.1) накладывается условие

$$\begin{gathered}(\varepsilon_J)_{AB} = -(\varepsilon_I)_{CB}(\varepsilon_J)^{CD}(\varepsilon_I)_{AD}, \quad \eta_\Lambda{}^{AB}(\varepsilon_I)_{AD}(\varepsilon_I)_{CB} = -\eta_{\Lambda CD} + (\eta_I)_\Lambda(\varepsilon_I)_{CD}, \\ (\eta_I)^\Lambda = \tfrac{2}{N}\eta^{\Lambda AB}(\varepsilon_I)_{AB}, \quad (\varepsilon_I)_{AB} = (\eta_I)^\Lambda\eta_{\Lambda AB}, \quad (\varepsilon_I)_{AB}(\varepsilon_I)^{AC} = \delta_B{}^C.\end{gathered} \tag{8.16}$$

Определение 8.2. *На пространстве бивекторов определен 4-валентный тензор, с помощью которого можно поднимать и опускать парные индексы*

$$\begin{aligned} &\varepsilon_{ABCD} := \eta_{\Lambda AB}\eta^\Lambda{}_{CD}, && \bar{\varepsilon}_{A'B'C'D'} := \bar{\eta}_{\Lambda' A'B'}\bar{\eta}^{\Lambda'}{}_{C'D'}, \\ &g_{\Lambda\Psi} = \left(\tfrac{2}{N}\right)^2\eta^{\Lambda AB}\eta^{\Psi CD}\varepsilon_{ABCD}, && \bar{g}_{\Lambda'\Psi'} = \left(\tfrac{2}{N}\right)^2\bar{\eta}^{\Lambda' A'B'}\bar{\eta}^{\Psi' C'D'}\bar{\varepsilon}_{A'B'C'D'}, \\ &\varepsilon_{ADCB}\eta_\Lambda{}^{AB} = \tfrac{2-n}{2}\eta_{\Lambda CD}, && \bar{\varepsilon}_{A'D'C'B'}\bar{\eta}_{\Lambda'}{}^{A'B'} = \tfrac{2-n}{2}\bar{\eta}_{\Lambda' C'D'}. \end{aligned} \tag{8.17}$$

Замечание 8.1. *Следует отметить, что для действительных вложений условие* $(\eta_I)^i(\eta_J)_i = 2\delta_{IJ}$ *не всегда выполнимо. Его следует модифицировать до* $(\eta_I)^i(\eta_J)_i = 2g_{IJ}$*, где* $g_{IJ} = 0, \quad I \neq J$ *и* $g_{II} = \pm 1$*, при этом знак выбирается в зависимости от индекса метрики и номера вектора. Это буквально означает, что метрика в некотором базисе имеет диагональный вид с* ± 1 *на главной диагонали. Тогда уравнение (8.16) примет вид*

$$\begin{gathered}(\varepsilon_J)_{AB} = -g_{II}(\varepsilon_I)_{CB}(\varepsilon_J)^{CD}(\varepsilon_I)_{AD}, \quad g_{II}\eta_i{}^{AB}(\varepsilon_I)_{AD}(\varepsilon_I)_{CB} = -\eta_{iCD} + (\eta_I)_i(\varepsilon_I)_{CD}, \\ (\eta_I)^i = \tfrac{2}{N}\eta^{iAB}(\varepsilon_I)_{AB}, \quad g_{II}(\varepsilon_I)_{AB} = (\eta_I)^i\eta_{iAB}, \quad g_{II}(\varepsilon_I)_{AB}(\varepsilon_I)^{AC} = \delta_B{}^C.\end{gathered} \tag{8.16'}$$

В случае комплексного пространства всегда можно ограничиться только $+1$ *с евклидовой метрикой* δ_{IJ}*. Этого будет достаточно для доказательства основных тождеств. К действительным же пространствам можно перейти, используя соответствующий оператор вложения.*

Поэтому возможно два существенно различных способа в зависимости от кратности n

$$I). \quad n \; mod \; 4 = 0, \qquad II). \quad n \; mod \; 4 = 2 \tag{8.18}$$

определения либо пары метрических тензоров $\varepsilon_{AB} = -\varepsilon_{BA}$, $\tilde{\varepsilon}_{AB} = \tilde{\varepsilon}_{BA}$ в случае I)., либо пары инволюций $E_A{}^B$, $\tilde{E}_A$ в случае II). (доказательство в приложении)

$$\begin{gathered} I). \quad \varepsilon_{A_1 A_{(n/2+q+1)}} := (\varepsilon_{(n/2+q)})_{A_{(n/2+q)}A_{(n/2+q+1)}} \prod_{Q=1}^{(n/2+q-1)/2} (\varepsilon_{2Q-1})_{A_{2Q-1}A_{2Q}}(\varepsilon_{2Q})^{A_{2Q+1}A_{2Q}}, \\ \varepsilon_{AB}\varepsilon^{CB} = \delta_A{}^C, \quad \left[\begin{array}{ll} \varepsilon_{AB} = -\varepsilon_{BA}, & n/2+q > 2, \\ \varepsilon_{AB} = \varepsilon_{BA}, & n/2+q = 1, \end{array}\right], \quad \eta_{\Lambda AB} = \eta_\Lambda{}^{CD}\varepsilon_{CA}\varepsilon_{DB}, \\ \\ \tilde{\varepsilon}_{A_1 A_{(n/2+\tilde{q})}} := (\varepsilon_{\widetilde{(n/2+\tilde{q})}})_{A_{(n/2+\tilde{q})}A_{(n/2+\tilde{q}+1)}} \prod_{\tilde{Q}=1}^{(n/2+\tilde{q}-1)/2} (\varepsilon_{2\tilde{Q}-1})_{A_{2\tilde{Q}-1}A_{2\tilde{Q}}}(\varepsilon_{2\tilde{Q}})^{A_{2\tilde{Q}}A_{2\tilde{Q}+1}}, \\ \tilde{\varepsilon}_{AB}\tilde{\varepsilon}^{CB} = \delta_A{}^C, \quad \left[\begin{array}{ll} \tilde{\varepsilon}_{AB} = \tilde{\varepsilon}_{BA}, & n/2+q > 2, \\ \tilde{\varepsilon}_{AB} = -\tilde{\varepsilon}_{BA}, & n/2+q = 1, \end{array}\right], \quad \eta_{\Lambda AB} = \eta_\Lambda{}^{CD}\tilde{\varepsilon}_{CA}\tilde{\varepsilon}_{DB}, \end{gathered} \tag{8.19}$$

$$\begin{gathered} II). \quad E_{A_1}{}^{A_{(n/2+q+1)}} := \prod_{Q=1}^{(n/2+q)/2} (\varepsilon_{2Q-1})_{A_{2Q-1}A_{2Q}}(\varepsilon_{2Q})^{A_{2Q+1}A_{2Q}}, \\ E_A{}^B E_B{}^C = -\delta_A{}^C, \quad \eta_\Lambda{}^{AB} = -\eta_\Lambda{}^{CD}E_D{}^A E_C{}^B, \\ \\ \tilde{E}_{A_1}{}^{A_{(n/2+\tilde{q}+1)}} := \prod_{\tilde{Q}=1}^{(n/2+\tilde{q})/2} (\varepsilon_{2\tilde{Q}-1})_{A_{2\tilde{Q}-1}A_{2\tilde{Q}}}(\varepsilon_{2\tilde{Q}})^{A_{2\tilde{Q}+1}A_{2\tilde{Q}}}, \\ \tilde{E}_A{}^B \tilde{E}_B{}^C = -\delta_A{}^C, \quad \eta_\Lambda{}^{AB} = \eta_\Lambda{}^{CD}\tilde{E}_D{}^A \tilde{E}_C{}^B, \end{gathered} \tag{8.19'}$$

Замечание 8.2. *Для действительных вложений тензоры из (8.19) домножаются на* $g := (\prod_{P=1}^{(n/2+q)} \sqrt{g_{PP}})$*, где в качестве корня* $\sqrt{g_{\tilde{P}\tilde{P}}}$ *при* $g_{\tilde{P}\tilde{P}} = -1$ *выбирается любой из двух вариантов* $\pm i$.

Покажем, что такие операторы существуют. Для n=4 возможно представление, описанное в примере 8.1. Пусть построено некоторое представление ${\eta_\alpha}^{ab}$ $(\alpha\ , \beta\ , ... = \overline{1, n-2}; a\ , b\ , ... = \overline{1, 2^{(n-2)/2-1}})$. Операторы ${\eta_\Lambda}^{AB}$ $(\Lambda\ , \Psi\ , ... = \overline{1, n}; A\ , B\ , ... = \overline{1, 2^{n/2-1}})$ строятся следующим образом. Пусть тензор $g_{\alpha\beta}$ содержит только $+1$ на главной диагонали, тогда

$$
\begin{aligned}
{\eta_\alpha}^{AB} &= \begin{pmatrix} {\eta_\alpha}^{ab} & 0 \\ 0 & -(\eta^T)_{\alpha cd} \end{pmatrix}, \quad & {\eta^\alpha}_{AB} &= \begin{pmatrix} {\eta^\alpha}_{pk} & 0 \\ 0 & -(\eta^T)^{\alpha lm} \end{pmatrix}, \\
{\eta_{n-1}}^{AB} &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & i{\delta^a}_d \\ -i{\delta_c}^b & 0 \end{pmatrix}, \quad & {\eta^{n-1}}_{AB} &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & -i{\delta_p}^m \\ i{\delta^l}_k & 0 \end{pmatrix}, \\
{\eta_n}^{AB} &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & {\delta^a}_d \\ {\delta_c}^b & 0 \end{pmatrix}, \quad & {\eta^n}_{AB} &= \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & {\delta_p}^m \\ {\delta^l}_k & 0 \end{pmatrix},
\end{aligned}
\tag{8.20}
$$

При общей схеме построения (8.20) уравнение

$$
{\tilde{T}_A}{}^B(\varepsilon_I)_{BC} + {\tilde{\tilde{T}}_C}{}^B(\varepsilon_I)_{AB} = 0 \tag{8.21}
$$

для n=4 приводит к ${\tilde{T}_A}{}^B = {\tilde{\tilde{T}}_A}{}^B = 0$, что проверяется непосредственно. Пусть данное уравнение выполнено для некоторого четного n-2. Тогда для n уравнение распадется на части такого вида:

1. $$
{\tilde{T}_a}{}^b(\varepsilon_I)_{bc} + {\tilde{\tilde{T}}_c}{}^b(\varepsilon_I)_{ab} = 0, \quad {\tilde{T}^a}{}_b(\varepsilon_I)^{bc} + {\tilde{\tilde{T}}^c}{}_b(\varepsilon_I)^{ab} = 0, \quad I = \overline{1, n-2}, \tag{8.22}
$$

   что немедленно дает ${\tilde{T}_a}{}^b = {\tilde{\tilde{T}}_a}{}^b = 0, \quad {\tilde{T}^a}{}_b = {\tilde{\tilde{T}}^a}{}_b = 0$.

2. $$
\begin{aligned}
\tilde{T}_{ab}{(\varepsilon_{n+1})^b}_c + \tilde{\tilde{T}}_{cb}{(\varepsilon_{n+1})_a}^b &= \tilde{T}_{ac} - \tilde{\tilde{T}}_{ca} = 0, \\
\tilde{T}^{ab}{(\varepsilon_{n+1})_b}^c + \tilde{\tilde{T}}^{cb}{(\varepsilon_{n+1})^a}_b &= -\tilde{T}^{ac} + \tilde{\tilde{T}}^{ca} = 0.
\end{aligned}
\tag{8.23}
$$

3. $$
\begin{aligned}
\tilde{T}_{ab}{(\varepsilon_{n+2})^b}_c + \tilde{\tilde{T}}_{cb}{(\varepsilon_{n+2})_a}^b &= \tilde{T}_{ac} + \tilde{\tilde{T}}_{ca} = 0, \\
\tilde{T}^{ab}{(\varepsilon_{n+2})_b}^c + \tilde{\tilde{T}}^{cb}{(\varepsilon_{n+2})^a}_b &= \tilde{T}^{ac} + \tilde{\tilde{T}}^{ca} = 0,
\end{aligned}
\tag{8.24}
$$

   что немедленно дает $\tilde{T}_{ab} = \tilde{\tilde{T}}_{ab} = 0, \quad \tilde{T}^{ab} = \tilde{\tilde{T}}^{ab} = 0$. Таким образом опять ${\tilde{T}_A}{}^B = {\tilde{\tilde{T}}_A}{}^B = 0$.

Пусть теперь имеются вещественные связующие операторы $\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}$. При этом существует набор ортогональных векторов такой, что часть $(\varepsilon_Q)^{\tilde{\tilde{A}}\tilde{\tilde{B}}}$ симметрична, а оставшаяся часть $(\varepsilon_{\tilde{Q}})^{\tilde{\tilde{A}}\tilde{\tilde{B}}}$ кососимметрична. Покажем, что возможно перейти к комплексной реализации таких операторов, сохраняющих указанное свойство и, следовательно, удовлетворяющих определению 8.1.

Алгоритм 8.1.

1. На первом шаге используем в качестве операторов (6.24) и (6.30) операторы

$$\begin{gathered}
\underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} \delta^a{}_{\tilde{\tilde{a}}} & 0 & 0 & i\delta^a{}_{\tilde{\tilde{q}}} \\ 0 & \delta_c{}^{\tilde{\tilde{c}}} & \delta_c{}^{\tilde{\tilde{p}}} & 0 \end{pmatrix}}_{:=(\tilde{\tilde{m}}_{1_+})^A{}_{\tilde{\tilde{A}}}} \cdot \\
\cdot \underbrace{\begin{pmatrix} \eta_\alpha{}^{\tilde{\tilde{a}}\tilde{\tilde{b}}} & \alpha\delta^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{d}}} & \gamma\delta^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{k}}} & 0 \\ \beta\delta_{\tilde{\tilde{c}}}{}^{\tilde{\tilde{b}}} & -(\eta^T)_{\alpha\tilde{\tilde{c}}\tilde{\tilde{d}}} & 0 & \gamma\delta_{\tilde{\tilde{c}}}{}^{\tilde{\tilde{l}}} \\ \delta\delta_{\tilde{\tilde{p}}}{}^{\tilde{\tilde{b}}} & 0 & -(\eta^T)_{\alpha\tilde{\tilde{p}}\tilde{\tilde{k}}} & -\alpha\delta_{\tilde{\tilde{p}}}{}^{\tilde{\tilde{l}}} \\ 0 & \delta\delta^{\tilde{\tilde{q}}}{}_{\tilde{\tilde{d}}} & -\beta\delta^{\tilde{\tilde{q}}}{}_{\tilde{\tilde{k}}} & \eta_\alpha{}^{\tilde{\tilde{q}}\tilde{\tilde{l}}} \end{pmatrix}}_{\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}} \underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} \delta_{\tilde{\tilde{b}}}{}^b & 0 \\ 0 & \delta^{\tilde{\tilde{d}}}{}_d \\ 0 & \delta^{\tilde{\tilde{k}}}{}_d \\ -i\delta_{\tilde{\tilde{l}}}{}^b & 0 \end{pmatrix}}_{:=(\tilde{m}^T_{1_+})_{\tilde{\tilde{B}}}{}^B} = \\
= \underbrace{\begin{pmatrix} \eta_\alpha{}^{ab} & \underbrace{\frac{1}{2}((\alpha - i\beta) + (\gamma + i\delta))}_{:=\phi_1}\delta^a{}_d \\ \underbrace{\frac{i}{2}((\alpha - i\beta) - (\gamma + i\delta))}_{:=\psi_1}\delta_c{}^b & -(\eta^T)_{\alpha cd} \end{pmatrix}}_{:=\eta_{1_+\tilde{\Lambda}}{}^{AB}}.
\end{gathered} \tag{8.25}$$

В качестве операторов $(\tilde{m}_{1_-})^A{}_{\tilde{\tilde{A}}}$ *можно взять оператор* $\begin{pmatrix} \delta^a{}_{\tilde{\tilde{a}}} & 0 & 0 & -i\delta^a{}_{\tilde{\tilde{q}}} \\ 0 & \delta_c{}^{\tilde{\tilde{c}}} & -\delta_c{}^{\tilde{\tilde{p}}} & 0 \end{pmatrix}$. *Если теперь воспользоваться операторами* $m_\Lambda{}^{\tilde{\Lambda}}$, *переводящими подпространство* $\mathbb{R}^4_{(2,2)}$ *в подпространство* $\mathbb{C}^2$, *а пространство* $\mathbb{R}^{2n-4}_{(n-2,n-2)}$ *оставляющее без изменений, то можно построить связующие операторы* $\eta_\Lambda{}^{ab}$, *соответствующие пространству* $\mathbb{C}^2 \oplus \mathbb{R}^{2n-4}_{(n-2,n-2)}$, *при этом размерность спинорного пространства будет понижена в 2 раза.*

2. На втором шаге воспользуемся операторами

$$\underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} \delta^{a}{}_{\tilde{\tilde{a}}} & 0 & 0 & i\delta^{a}{}_{\tilde{\tilde{q}}} & 0 & 0 & 0 & 0 \\ 0 & \delta_{c}{}^{\tilde{\tilde{c}}} & \delta_{c}{}^{\tilde{\tilde{p}}} & 0 & 0 & 0 & 0 & \\ 0 & 0 & 0 & 0 & \delta_{a_1}{}^{\tilde{\tilde{a}}_1} & 0 & 0 & i\delta_{a_1}{}^{\tilde{\tilde{p}}_1} \\ 0 & 0 & 0 & 0 & 0 & \delta^{c_1}{}_{\tilde{\tilde{c}}_1} & \delta^{c_1}{}_{\tilde{\tilde{q}}_1} & 0 \end{pmatrix}}_{:=(\tilde{\tilde{m}}_2)^{A}{}_{\tilde{\tilde{A}}}} \cdot$$

$$\underbrace{\begin{pmatrix} \eta_\alpha{}^{\tilde{\tilde{a}}\tilde{\tilde{b}}} & \alpha\delta^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{d}}} & \gamma\delta^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{k}}} & 0 & \phi_1\delta^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{b}}_1} & 0 & 0 & 0 \\ \beta\delta_{\tilde{\tilde{c}}}{}^{\tilde{\tilde{b}}} & -(\eta^T)_{\alpha\tilde{\tilde{c}}\tilde{\tilde{d}}} & 0 & \gamma\delta_{\tilde{\tilde{c}}}{}^{\tilde{\tilde{l}}} & 0 & \phi_1\delta_{\tilde{\tilde{c}}}{}^{\tilde{\tilde{d}}_1} & 0 & 0 \\ \delta\delta_{\tilde{\tilde{p}}}{}^{\tilde{\tilde{b}}} & 0 & -(\eta^T)_{\alpha\tilde{\tilde{p}}\tilde{\tilde{k}}} & -\alpha\delta_{\tilde{\tilde{p}}}{}^{\tilde{\tilde{l}}} & 0 & 0 & \phi_1\delta_{\tilde{\tilde{p}}}{}^{\tilde{\tilde{k}}_1} & 0 \\ 0 & \delta\delta^{\tilde{\tilde{q}}}{}_{\tilde{\tilde{d}}} & -\beta\delta^{\tilde{\tilde{q}}}{}_{\tilde{\tilde{k}}} & \eta_\alpha{}^{\tilde{\tilde{q}}\tilde{\tilde{l}}} & 0 & 0 & 0 & \phi_1\delta^{\tilde{\tilde{p}}}{}_{\tilde{\tilde{l}}_1} \\ \psi_1\delta_{\tilde{\tilde{a}}_1}{}^{\tilde{\tilde{b}}} & 0 & 0 & 0 & -(\eta^T)_{\alpha\tilde{\tilde{a}}_1\tilde{\tilde{b}}_1} & -\alpha\delta_{\tilde{\tilde{a}}_1}{}^{\tilde{\tilde{d}}_1} & -\gamma\delta_{\tilde{\tilde{a}}_1}{}^{\tilde{\tilde{k}}_1} & 0 \\ 0 & \psi_1\delta^{\tilde{\tilde{c}}_1}{}_{\tilde{\tilde{d}}} & 0 & 0 & -\beta\delta^{\tilde{\tilde{c}}_1}{}_{\tilde{\tilde{b}}_1} & \eta_\alpha{}^{\tilde{\tilde{c}}_1\tilde{\tilde{d}}_1} & 0 & -\gamma\delta^{\tilde{\tilde{a}}_1}{}_{\tilde{\tilde{l}}_1} \\ 0 & 0 & \psi_1\delta^{\tilde{\tilde{q}}_1}{}_{\tilde{\tilde{k}}} & 0 & -\delta\delta^{\tilde{\tilde{q}}_1}{}_{\tilde{\tilde{b}}_1} & 0 & \eta_\alpha{}^{\tilde{\tilde{q}}_1\tilde{\tilde{k}}_1} & \alpha\delta^{\tilde{\tilde{q}}_1}{}_{\tilde{\tilde{l}}_1} \\ 0 & 0 & 0 & \psi_1\delta_{\tilde{\tilde{p}}_1}{}^{\tilde{\tilde{l}}} & 0 & -\delta\delta_{\tilde{\tilde{p}}_1}{}^{\tilde{\tilde{d}}_1} & \beta\delta_{\tilde{\tilde{p}}_1}{}^{\tilde{\tilde{l}}_1} & -(\eta^T)_{\alpha\tilde{\tilde{p}}_1\tilde{\tilde{l}}_1} \end{pmatrix}}_{\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}}$$

$$\cdot\underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} \delta_{\tilde{\tilde{b}}}{}^{b} & 0 & 0 & 0 \\ 0 & \delta^{\tilde{\tilde{d}}}{}_{d} & 0 & 0 \\ 0 & \delta^{\tilde{\tilde{k}}}{}_{d} & 0 & 0 \\ -i\delta_{\tilde{\tilde{l}}}{}^{b} & 0 & 0 & 0 \\ 0 & 0 & \delta^{\tilde{\tilde{b}}_1}{}_{d_2} & 0 \\ 0 & 0 & 0 & \delta^{\tilde{\tilde{d}}_1}{}_{b_2} \\ 0 & 0 & 0 & \delta^{\tilde{\tilde{k}}_1}{}_{b_2} \\ 0 & 0 & -i\delta^{\tilde{\tilde{l}}_1}{}_{d_2} & 0 \end{pmatrix}}_{:=(\tilde{m}_2^T)_{\tilde{\tilde{B}}}{}^{B}} . \tag{8.26}$$

Если теперь воспользоваться операторами $m_\Lambda{}^{\tilde{\Lambda}}$, *переводящими подпространство* $\mathbb{R}^4_{(2,2)}$ *в подпространство* $\mathbb{C}^2$, *а пространство* $\mathbb{C}^2 \oplus \mathbb{R}^{2n-8}_{(n-4,n-4)}$ *оставляющее без изменений, то можно построить связующие операторы* $\eta_\Lambda{}^{ab}$, *соответствующие пространству* $\mathbb{C}^4 \oplus \mathbb{R}^{2n-8}_{(n-4,n-4)}$, *при этом размерность спинорного пространства будет понижена в 2 раза.*

3. Поступая аналогично, можно получить пространство $\mathbb{C}^{n-2} \oplus \mathbb{R}^4_{(2,2)}$ *и соответствующие связующие операторы. Очевидно, что на каждом шаге у связующих операторов модифицируются только компоненты, отвечающие за переход от действительной реализации подпространства* $\mathbb{R}^4_{(2,2)}$ *к его комплексной реализации* $\mathbb{C}^2$.

4. Заключительный шаг осуществляется с помощью операторов из примера 3.2.

Следствие 8.2. *Связующие операторы из определения 8.1 удовлетворяют тождеству (доказательство в приложении):*

$$\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}\eta^\Theta{}_{CB} = \frac{N}{4}(g_{\Lambda\Psi}g^{\Omega\Theta} + \delta_\Lambda{}^\Theta\delta_\Psi{}^\Omega - \delta_\Psi{}^\Theta\delta_\Lambda{}^\Omega). \tag{8.27}$$

Следствие 8.3. *Для ортогональных преобразований (6.42) существует два варианта представлений для каждого случая:*

I). Собственные вращения.

$$S_\Lambda{}^\Psi\eta_\Psi{}^{AB} = \eta_\Lambda{}^{CD}\tilde{S}_C{}^A\tilde{\tilde{S}}_D{}^B. \tag{8.28}$$

1. n mod 4 =2

$$\tilde{\tilde{S}}_A{}^B = -E_A{}^C\tilde{S}_C{}^D E_D{}^B, \quad \tilde{\tilde{S}}_A{}^B = -\tilde{E}_A{}^C\tilde{S}_C{}^D\tilde{E}_D{}^B. \tag{8.29}$$

2. n mod 4 =0

$$\begin{aligned} &\tilde{\tilde{S}}_A{}^B\tilde{\tilde{S}}_C{}^D\varepsilon_{BD} = \varepsilon_{AC}, \quad \tilde{S}_A{}^B\tilde{S}_C{}^D\varepsilon_{BD} = \varepsilon_{AC}, \\ &\tilde{\tilde{S}}_A{}^B\tilde{\tilde{S}}_C{}^D\tilde{\varepsilon}_{BD} = \tilde{\varepsilon}_{AC}, \quad \tilde{S}_A{}^B\tilde{S}_C{}^D\tilde{\varepsilon}_{BD} = \tilde{\varepsilon}_{AC}. \end{aligned} \tag{8.30}$$

II). Несобственные вращения.

$$S_\Lambda{}^\Psi\eta_\Psi{}^{BA} = \eta_{\Lambda CD}\tilde{S}^{CA}\tilde{\tilde{S}}^{DB}. \tag{8.31}$$

1. n mod 4 =2

$$\tilde{\tilde{S}}^{CA} = -E_K{}^A\tilde{S}^{KL}E_L{}^C, \quad \tilde{\tilde{S}}^{CA} = \tilde{E}_K{}^A\tilde{S}^{KL}\tilde{E}_L{}^C. \tag{8.32}$$

2. n mod 4 =0

$$\begin{aligned} &\tilde{\tilde{S}}^{AB}\tilde{\tilde{S}}^{CD}\varepsilon_{BD} = \varepsilon^{AC}, \quad \tilde{S}^{AB}\tilde{S}^{CD}\varepsilon_{BD} = \varepsilon^{AC}, \\ &\tilde{\tilde{S}}^{AB}\tilde{\tilde{S}}^{CD}\tilde{\varepsilon}_{BD} = \tilde{\varepsilon}^{AC}, \quad \tilde{S}^{AB}\tilde{S}^{CD}\tilde{\varepsilon}_{BD} = \tilde{\varepsilon}^{AC}. \end{aligned} \tag{8.33}$$

Следствие 8.4. *Для инволюций (6.47) существует два варианта разложения:*

I). Аналог собственных вращений.

$$S_\Lambda{}^{\Psi'}\bar{\eta}_{\Psi'}{}^{A'B'} = \eta_\Lambda{}^{CD}\tilde{S}_C{}^{A'}\tilde{\tilde{S}}_D{}^{B'}. \tag{8.34}$$

$$\tilde{\tilde{S}}_A{}^{B'}\bar{\tilde{\tilde{S}}}_{B'}{}^D = \pm\delta_A{}^D, \quad \tilde{S}_A{}^{B'}\bar{\tilde{S}}_{B'}{}^D = \pm\delta_A{}^D. \tag{8.35}$$

II). Аналог несобственных вращений.

$$S_\Lambda{}^{\Psi'}\bar{\eta}_{\Psi'}{}^{B'A'} = \eta_{\Lambda CD}\tilde{S}^{CA'}\tilde{\tilde{S}}^{DB'}. \tag{8.36}$$

$$\tilde{\tilde{S}}^{AB'} = \pm\bar{\tilde{\tilde{S}}}^{B'A}, \quad \tilde{S}^{AB'} = \pm\bar{\tilde{S}}^{B'A}. \tag{8.37}$$

Будем считать, что операторы $\eta_\Lambda{}^{AB}(\sigma_{\Lambda AB})$ строятся либо по схеме (8.20)$(\alpha = \overline{1, n-2})$

$$a). \quad \begin{aligned} \eta_\Lambda{}^{AB} &= \begin{pmatrix} \eta_\alpha{}^{ab} & \frac{1}{2}(i\eta_{n-1} + \eta_n)\delta^a{}_d \\ \frac{1}{2}(-i\eta_{n-1} + \eta_n)\delta_c{}^b & -(\eta^T)_{\alpha cd} \end{pmatrix}, \\ \sigma_{\Lambda AB} &= \begin{pmatrix} (\eta^T)_{\alpha ab} & \frac{1}{2}(i\eta_{n-1} + \eta_n)\delta_a{}^d \\ \frac{1}{2}(-i\eta_{n-1} + \eta_n)\delta^c{}_b & -\eta_\alpha{}^{cd} \end{pmatrix}. \end{aligned} \tag{8.38}$$

либо по схеме

$$b). \quad \begin{aligned} \eta_\Lambda{}^{AB} &= \begin{pmatrix} \eta_\alpha{}^{ab} & \frac{1}{2}(i\eta_{n-1} + \eta_n)\delta^a{}_d \\ \frac{1}{2}(i\eta_{n-1} - \eta_n)\delta_c{}^b & -(\eta^T)_{\alpha cd} \end{pmatrix}, \\ \sigma_{\Lambda AB} &= \begin{pmatrix} (\eta^T)_{\alpha ab} & \frac{1}{2}(i\eta_{n-1} + \eta_n)\delta_a{}^d \\ \frac{1}{2}(i\eta_{n-1} - \eta_n)\delta^c{}_b & -\eta_\alpha{}^{cd} \end{pmatrix}. \end{aligned} \tag{8.38$'$}$$

Это означает, что из предыдущей размерности наследуются спиноры сопряжения (6.47)

$$\begin{aligned} &I). \quad a). \quad S_A{}^{A'} := \begin{pmatrix} 0 & iS_{aa'} \\ -iS^{bb'} & 0 \end{pmatrix}, \quad b). \quad S_A{}^{A'} := \begin{pmatrix} 0 & iS_{aa'} \\ iS^{bb'} & 0 \end{pmatrix}, \\ &II). \quad a). \quad S_{AA'} := \begin{pmatrix} 0 & iS_a{}^{a'} \\ -iS^b{}_{b'} & 0 \end{pmatrix}, \quad b). \quad S_{AA'} := \begin{pmatrix} 0 & iS_a{}^{a'} \\ iS^b{}_{b'} & 0 \end{pmatrix}, \end{aligned} \tag{8.39}$$

которые определяются с точностью до знака.

Example 8.3. *В таблице 1 приведены различные варианты действительных вложений для n=2 (см. примеры 6.1 и 6.2).*

Example 8.4. *В таблице 2 приведены различные варианты действительных вложений для n=4 (см. пример 6.3).*

Example 8.5. *В таблице 3 приведены различные варианты действительных вложений для n=6. В случае, когда $\tilde{\eta}_\Lambda{}^{AC} := \eta_\Lambda{}^{AB}\varepsilon_B{}^C$ кососимметричен (см. пример 8.2), должны быть выполнены соотношения:*

$$\begin{aligned} &I). \quad \tilde{\eta}_\Lambda{}^{AB}S_A{}^{A'}S_B{}^{B'} = S_\Lambda{}^{\Lambda'}\bar{\tilde{\eta}}_{\Lambda'}{}^{A'B'}, &&II). \quad \tilde{\eta}_\Lambda{}^{AB}S_{AA'}S_{BB'} = S_\Lambda{}^{\Lambda'}\bar{\tilde{\eta}}_{\Lambda'B'A'} = S_\Lambda{}^{\Lambda'}\bar{\tilde{\sigma}}_{\Lambda'A'B'}, \\ & \eta_\Lambda{}^{AB}S_A{}^{A'}\tilde{S}_B{}^{B'} = S_\Lambda{}^{\Lambda'}\bar{\eta}_{\Lambda'}{}^{A'B'}, && \eta_\Lambda{}^{AB}S_{AA'}\tilde{S}_{BB'} = S_\Lambda{}^{\Lambda'}\bar{\eta}_{\Lambda'B'A'} = S_\Lambda{}^{\Lambda'}\bar{\sigma}_{\Lambda'A'B'}, \end{aligned}$$

$$I). \quad \tilde{S}_A{}^{A'} = \pm\varepsilon_A{}^B S_B{}^{B'}\bar{\varepsilon}^{A'}{}_{B'}, \qquad II). \quad \tilde{S}_{AA'} = \pm\varepsilon_A{}^B S_{BB'}\varepsilon_{A'}{}^{B'}. \tag{8.40}$$

Напомним, что $\eta_i := H_i{}^\Lambda\eta_\Lambda$ согласно (4.4).

Таблица 1: Вид матрицы тензора S для соответствующих действительных вложений 2-мерных пространств в комплексное для $r^i = (T, Z)$.

П-п	Пространство	Вид S	S	Изоморфизм	Вид $\sqrt{2}r^i\eta_i$	Вид $\sqrt{2}r^i\sigma_i$
1a	R^2 $\ll ++ \gg$	$S_{AA'}$	$\begin{pmatrix} 1 \end{pmatrix}$	$U(1) \cong SO^+(2)$	$T + iZ$	$T - iZ$
1b	R^2 $\ll -- \gg$	$S_{AA'}$	$\begin{pmatrix} i \end{pmatrix}$	$U(1) \cong SO^+(2)$	$-iT - Z$	$-iT + Z$
2a	$R^2_{(1,1)}$ $\ll +- \gg$	$S_A{}^{A'}$	$\begin{pmatrix} 1 \end{pmatrix}$	$R_+ \cong SO^+(1,1)$	$T + Z$	$T - Z$
2b	$R^2_{(1,1)}$ $\ll -+ \gg$	$S_A{}^{A'}$	$\begin{pmatrix} 1 \end{pmatrix}$	$R_+ \cong SO^+(1,1)$	$-T - Z$	$T - Z$

Таблица 2: Вид матрицы тензора S для соответствующих действительных вложений 4-мерных пространств в комплексное для $r^i = (T, X, Y, Z)$.

П-п	Пространство	S	Вид $\sqrt{2}r^i\eta_i/\sqrt{2}r^i\sigma_i$
1a	R^4 $\ll ---- \gg$ $SU(2)\times SU(2)/\pm 1 \cong SO^+(4)$	$\tilde{S}_A{}^{A'} = \tilde{\tilde{S}}_A{}^{A'}$ $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$	$\begin{pmatrix} -iT-Z & iX+Y \\ iX-Y & iT-Z \end{pmatrix}$ $\begin{pmatrix} -iT+Z & iX+Y \\ iX-Y & iT+Z \end{pmatrix}$
1б	R^4 $\ll ++++ \gg$ $SU(2)\times SU(2)/\pm 1 \cong SO^+(4)$	$\tilde{S}_A{}^{A'} = \tilde{\tilde{S}}_A{}^{A'}$ $\begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$	$\begin{pmatrix} T+iZ & iY+X \\ -iY+X & -T+iZ \end{pmatrix}$ $\begin{pmatrix} T-iZ & iY+X \\ -iY+X & -T-iZ \end{pmatrix}$
2a	$R^4_{(3,1)}$ $\ll +-++ \gg$ $SL(2,\mathbb{C})/\pm 1 \cong SO^+(3,1)$	$\tilde{S}_{AA'} = \tilde{\tilde{S}}_{AA'}$ $\begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$	$\begin{pmatrix} T+Z & iY+X \\ -iY+X & -T+Z \end{pmatrix}$ $\begin{pmatrix} T-Z & iY+X \\ -iY+X & -T-Z \end{pmatrix}$
2b	$R^4_{(1,3)}$ $\ll +--- \gg$ $SL(2,\mathbb{C})/\pm 1 \cong SO^+(1,3)$	$\tilde{S}_{AA'} = \tilde{\tilde{S}}_{AA'}$ $\begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$	$\begin{pmatrix} T+Z & iY+X \\ iY-X & -T+Z \end{pmatrix}$ $\begin{pmatrix} T-Z & iY+X \\ iY-X & -T-Z \end{pmatrix}$
3a	$R^4_{(2,2)}$ $\ll ++-- \gg$ $SL(2,\mathbb{R})\times SL(2,\mathbb{R})/\pm 1 \cong SO^+(2,2)$	$\tilde{S}_A{}^{A'} = \tilde{\tilde{S}}_A{}^{A'}$ $\begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$	$\begin{pmatrix} T+iZ & iY+X \\ iY-X & -T+iZ \end{pmatrix}$ $\begin{pmatrix} T-iZ & iY+X \\ iY-X & -T-iZ \end{pmatrix}$
3b	$R^4_{(2,2)}$ $\ll --++ \gg$ $SL(2,\mathbb{R})\times SL(2,\mathbb{R})/\pm 1 \cong SO^+(2,2)$	$\tilde{S}_A{}^{A'} = \tilde{\tilde{S}}_A{}^{A'}$ $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$	$\begin{pmatrix} -iT-Z & iX+Y \\ -iX+Y & iT-Z \end{pmatrix}$ $\begin{pmatrix} -iT+Z & iX+Y \\ -iX+Y & iT+Z \end{pmatrix}$

Таблица 3: Вид матрицы тензора S для соответствующих действительных вложений 6-мерных пространств в комплексное для $r^i = (T, X, V, W, Y, Z)$.

П-п	Пр-во	Вид S	Изоморфизм
			Вид $\sqrt{2} r^i \eta_i$
			Вид $\sqrt{2} r^i \sigma_i$
1	R^6	$S_{AA'}$	$SU(4)/\pm 1 \cong SO^+(6)$
		$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} T+iZ & X+iY & V+iW & 0 \\ X-iY & -T+iZ & 0 & V+iW \\ V-iW & 0 & -T+iZ & -X-iY \\ 0 & V-iW & -X+iY & T+iZ \end{pmatrix}$
			$\begin{pmatrix} T-iZ & X+iY & V+iW & 0 \\ X-iY & -T-iZ & 0 & V+iW \\ V-iW & 0 & -T-iZ & -X-iY \\ 0 & V-iW & -X+iY & T-iZ \end{pmatrix}$
2	$R^6_{(1,5)}$	${S_A}^{A'}$	$SL(2, \mathbb{H})/\pm 1 \cong SO^+(1,5)$
		$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} T+Z & X+iY & V+iW & 0 \\ -X+iY & -T+Z & 0 & V+iW \\ -V+iW & 0 & -T+Z & -X-iY \\ 0 & -V+iW & X-iY & T+Z \end{pmatrix}$
			$\begin{pmatrix} T-Z & X+iY & V+iW & 0 \\ -X+iY & -T-Z & 0 & V+iW \\ -V+iW & 0 & -T-Z & -X-iY \\ 0 & -V+iW & X-iY & T-Z \end{pmatrix}$
3	$R^6_{(2,4)}$	$S_{AA'}$	$SU(2,2)/\pm 1 \cong SO^+(2,4)$
		$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} T+iZ & X+iY & V+iW & 0 \\ -X+iY & -T+iZ & 0 & V+iW \\ -V+iW & 0 & -T+iZ & -X-iY \\ 0 & -V+iW & X-iY & T+iZ \end{pmatrix}$
			$\begin{pmatrix} T-iZ & X+iY & V+iW & 0 \\ -X+iY & -T-iZ & 0 & V+iW \\ -V+iW & 0 & -T-iZ & -X-iY \\ 0 & -V+iW & X-iY & T-iZ \end{pmatrix}$
4	$R^6_{(3,3)}$	${S_A}^{A'}$	$SL(4, \mathbb{R})/\pm 1 \cong SO^+(3,3)$
		$\begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} T+Z & X+iY & V+iW & 0 \\ X-iY & -T+Z & 0 & V+iW \\ -V+iW & 0 & -T+Z & -X-iY \\ 0 & -V+iW & -X+iY & T+Z \end{pmatrix}$
			$\begin{pmatrix} T-Z & X+iY & V+iW & 0 \\ X-iY & -T-Z & 0 & V+iW \\ -V+iW & 0 & -T-Z & -X-iY \\ 0 & -V+iW & -X+iY & T-Z \end{pmatrix}$

9 О структурных константах алгебр без деления. Седенионы

В этом параграфе рассказывается о том, как по связующим операторам, удовлетворяющим уравнению Клиффорда, построить структурные константы алгебры седенионов. Такие алгебры строятся индуктивным переходом для n mod 8=0, исходя из периодичности Ботта. Рассматривается аксиоматика таких алгебр. Вывод всех результатов этого параграфа сделан на основе [8], [27].

Воспользуемся тем фактом, что для n mod 8=0 существуют такие связующие операторы ${\eta_\Lambda}^{AB}$, индуктивное построение которых не является удвоением алгебр, а связано с периодичностью Ботта. Согласно определению 8.1 это означает, что

$$ {\eta_\Lambda}^{AB} + {\eta_\Lambda}^{BA} = \eta_\Lambda \varepsilon^{AB}, \quad \eta_\Lambda := (\eta_n)_\Lambda, \tag{9.1}$$

где ε^{AB} - симметрический метрический спин-тензор на спинорном пространстве. Это даст возможность положить

$$ I). \quad {P_\Lambda}^A := {\eta_\Lambda}^{BA} X_B, \quad {P_\Lambda}^A P_{\Psi A} = g_{\Lambda\Psi}, \qquad II). \quad {P_\Lambda}^A := {\eta_\Lambda}^{AB} X_B, \quad {P_\Lambda}^A P_{\Psi A} = g_{\Lambda\Psi} \tag{9.2}$$

для некоторого X_B $(X^A X_A = 2)$. Тогда можно определить некоторые структурные константы следующим образом

$$ I). \quad {\eta_{\Lambda\Psi}}^\Theta := \sqrt{2} {\eta_\Lambda}^{AB} P_{\Psi A} {P^\Theta}_B, \quad II). \quad {\eta_{\Lambda\Psi}}^\Theta := \sqrt{2} {\eta_\Lambda}^{BA} P_{\Psi A} {P^\Theta}_B. \tag{9.3}$$

Оба варианта равноправны, поэтому в дальнейшем будем рассматривать I). вариант. Будут ли они определять некоторую групповую алгебру? Ответ совсем неоднозначен, и вот почему.

1. По любым двум элементам $(r_1)^\Lambda$, $(r_2)^\Psi$ можно всегда единственным образом построить их произведение
$$ (r_1, r_2)^\Theta := (r_1)^\Lambda (r_2)^\Psi {\eta_{\Lambda\Psi}}^\Theta. \tag{9.4}$$
Поэтому первая аксиома умножения выполнена.

2. Существует единичный элемент по умножению $e := \frac{1}{\sqrt{2}} \eta^\Lambda$.

Доказательство.

$$\begin{aligned} \frac{1}{\sqrt{2}} \eta^\Lambda {\eta_{\Lambda\Psi}}^\Theta &= \varepsilon^{AB} P_{\Psi A} {P^\Theta}_B = {\delta_\Psi}^\Theta, \\ \frac{1}{\sqrt{2}} \eta^\Psi {\eta_{\Lambda\Psi}}^\Theta &= {\eta_\Lambda}^{AB} \varepsilon_{XA} X^X {P^\Theta}_B = {P_\Lambda}^B {P^\Theta}_B = {\delta_\Lambda}^\Theta. \end{aligned} \tag{9.5}$$

□

Поэтому вторая аксиома умножения выполнена.

3. Для любого неизотропного элемента существует обратный элемент по умножению

$$r^{-1} := \frac{1}{<r,r>}(2<r,e>e-r), \quad <r_1,r_2>:= g_{\Lambda\Psi}(r_1)^{\Lambda}(r_2)^{\Psi}. \tag{9.6}$$

Доказательство.

$$\begin{gathered}
(r^{-1})^{\Psi} = \tfrac{1}{r^{\Phi}r_{\Phi}} r^{\Omega}(\eta_{\Omega}\eta^{\Psi} - \delta_{\Omega}{}^{\Psi}),\\
r^{\Lambda}\tfrac{1}{r^{\Phi}r_{\Phi}} r^{\Omega}(\eta_{\Omega}\eta^{\Psi} - \delta_{\Omega}{}^{\Psi})\eta_{\Lambda\Psi}{}^{\Theta} = \tfrac{\sqrt{2}}{r^{\Phi}r_{\Phi}} R^{AB} r^{\Omega}(\eta_{\Omega}\eta^{\Psi} - \delta_{\Omega}{}^{\Psi})\eta_{\Psi XA} X^X P^{\Theta}{}_B =\\
= \tfrac{\sqrt{2}}{r^{\Phi}r_{\Phi}} R^{AB} R_{AX} X^X P^{\Theta}{}_B = \tfrac{1}{\sqrt{2}} X^B P^{\Theta}{}_B = \tfrac{1}{\sqrt{2}}\eta^{\Theta},\\
\tfrac{1}{r^{\Phi}r_{\Phi}} r^{\Omega}(\eta_{\Omega}\eta^{\Lambda} - \delta_{\Omega}{}^{\Lambda}) r^{\Psi}\eta_{\Lambda\Psi}{}^{\Theta} = \tfrac{\sqrt{2}}{r^{\Phi}r_{\Phi}} R^{BA} R_{XA} X^X P^{\Theta}{}_B = \tfrac{1}{\sqrt{2}}\eta^{\Theta}.
\end{gathered} \tag{9.7}$$

Эти тождества, верные для произвольного неизотропного r, равносильны соотношению

$$\eta_{\Lambda\Psi}{}^{\Omega} + \eta_{\Lambda}{}^{\Omega}{}_{\Psi} = \left(\frac{1}{\sqrt{2}}\eta_{\Lambda}\right)\delta_{\Psi}{}^{\Omega}. \tag{9.8}$$

□

Поэтому третья аксиома умножения выполнена частично. Для вложения $\mathbb{R}^n \subset \mathbb{C}^n$ $(n \ mod \ 8 = 0)$ эта аксиома выполнена для любого ненулевого элемента. Соответственно, с помощью операторов вложения $H_i{}^{\Lambda}$ определены структурные константы $\eta_{ij}{}^{k}$, которые, однако, не для всех спиноров X^A будут действительными.

4. Коммутативный закон не выполняется. Однако существует тождество

$$\tfrac{1}{2}((r_1,r_2)+(r_2,r_1)) = -<r_1,r_2>e+<r_1,e>r_2+<r_2,e>r_1. \tag{9.9}$$

Доказательство.

$$\begin{gathered}
\eta_{(\Lambda\Psi)}{}^{\Theta} = \sqrt{2}\eta_{(\Lambda}{}^{AB}P_{\Psi)A}P^{\Theta}{}_B = \sqrt{2}(-\eta_{(\Lambda}{}^{AB}\eta_{\Psi)AX} + 2\eta_{(\Lambda}{}^{AB}\eta_{\Psi)(AX)})X^X P^{\Theta}{}_B =\\
= -g_{\Lambda\Psi}\big(\tfrac{1}{\sqrt{2}}\eta^{\Theta}\big) + 2\big(\tfrac{1}{\sqrt{2}}\eta_{(\Psi}\big)\delta_{\Lambda)}{}^{\Theta}.
\end{gathered} \tag{9.10}$$

□

Частный случай этого тождества

$$r^2 = -<r,r>e + 2<r,e>r \tag{9.9'}$$

приводит к (9.6) и означает выполнение степенной ассоциативности. Поэтому четвертая аксиома умножения не выполнена. Но ей можно найти замену.

5. Ассоциативный закон не выполняется. Однако существует альтернативно-эластичное тождество
$$\begin{gathered}((r_1,r_1),r_2)-(r_1,(r_1,r_2))-(r_1,(r_2,r_1))= \\ =((r_2,(r_1,r_1))-((r_2,r_1),r_1)-((r_1,r_2),r_1).\end{gathered} \tag{9.11}$$

Доказательство.
$$\begin{gathered}((r_1,r_1),r_2)-(r_1,(r_1,r_2))-(r_1,(r_2,r_1))= \\ =-<r_1,r_1>r_2+2<r_1,e>(r_1,r_2)- \\ -2(r_1,-<r_1,r_2>e+<r_1,e>r_2+<r_2,e>r_1)= \\ =-<r_1,r_1>r_2+2<r_1,r_2>r_1- \\ -2<r_2,e>(-<r_1,r_1>e+2<r_1,e>r_1)= \\ =2<r_2,e><r_1,r_1>e+(2<r_1,r_2>-4<r_2,e><r_1,e>)r_1-<r_1,r_1>r_2.\end{gathered} \tag{9.12}$$

□

Поэтому пятая аксиома умножения не выполнена. Но ей можно найти замену.

Согласно общей теории такая алгебра имеет делители 0 при $n > 8$. Действительно,
$$\begin{gathered}(r_1,r_2)=0, \\ (r_1)^{\Lambda}(r_2)^{\Psi}\eta_{\Lambda\Psi}{}^{\Omega}=0, \\ (R_1)^{AB}(R_2)_{DA}X^{D}P^{\Omega}{}_{B}P_{\Omega}{}^{C}=0.\end{gathered} \tag{9.13}$$

Произведение $P^{\Omega}{}_{B}P_{\Omega}{}^{C}$ есть вырожденный оператор. Если принять, что известен вектор r_2, спинор X^A, то неизвестными будут координаты вектора r_1. Но, поскольку $P^{\Omega}{}_{B}P_{\Omega}{}^{C}$ вырожден, указанная система должна иметь нетривиальное решение при некоторых значениях вектора r_2. Однако, существует такой базис алгебры, для которого
$$\begin{gathered}(r_i,e)=(e,r_i),\quad e^2=e, \\ (r_i,r_j)=-(r_j,r_i),\quad r_i\neq e,\quad r_j\neq e,\quad i\neq j, \\ (r_i)^2=-e,\quad r_i\neq e.\end{gathered} \tag{9.14}$$

Наиболее интересными для изучения представляются алгебры, для которых можно построить такой базис (9.14), элементы которого не являются делителями нуля и удовлетворяют тождествам альтернативности (такую алгебру назовем алгеброй седенионов)
$$(r_i(r_i,r_j))=((r_i,r_i),r_j),\quad ((r_j,r_i),r_i)=(r_j,(r_i,r_i)). \tag{9.15}$$

Конечно, не для всех алгебр такой базис существует. Поэтому гиперкомплексные алгебры будут классифицироваться по максимальному количеству базисных элементов, не являющихся делителями нуля. Индуктивное построение метрической групповой альтернативно-эластичной алгебры более высокой размерности выглядит следующим образом.

Алгоритм 9.1. *1. Пусть нам известны кососимметричные связующие операторы $\eta_\alpha{}^{ab}$ для n mod 8 =6. Тогда связующие операторы $\eta_\Lambda{}^{AB}$ для n mod 8 =0 строятся по схеме*

$$\begin{aligned}
\eta_\Lambda{}^{AB} &= \begin{pmatrix} \eta_\alpha{}^{ab} & \frac{1}{2}(i\eta_{n-1}+\eta_n)\delta^a{}_d \\ \frac{1}{2}(-i\eta_{n-1}+\eta_n)\delta_c{}^b & -(\eta^T)_{\alpha cd} \end{pmatrix}, \\
\sigma_{\Lambda AB} &= \begin{pmatrix} (\eta^T)_{\alpha ab} & \frac{1}{2}(i\eta_{n-1}+\eta_n)\delta_a{}^d \\ \frac{1}{2}(-i\eta_{n-1}+\eta_n)\delta^c{}_b & -\eta_\alpha{}^{cd} \end{pmatrix},
\end{aligned} \tag{9.16}$$

что обеспечит наличие симметрического метрического спинора $\tilde{\varepsilon}_{AB} \equiv \varepsilon_{AB}$ для n mod 8 = 0.

2. Пусть нам известны связующие операторы $\eta_\alpha{}^{ab}$ для n mod 8 =0. Тогда связующие операторы $\eta_\Lambda{}^{AB}$ для n mod 8 =2 строятся по схеме

$$\begin{aligned}
\eta_\Lambda{}^{AB} &= \begin{pmatrix} \eta_\alpha{}^{ab} & \frac{1}{2}(i\eta_{n-1}+\eta_n)\delta^a{}_d \\ \frac{1}{2}(-i\eta_{n-1}+\eta_n)\delta_c{}^b & -(\eta^T)_{\alpha cd} \end{pmatrix}, \\
\sigma_{\Lambda AB} &= \begin{pmatrix} (\eta^T)_{\alpha ab} & \frac{1}{2}(i\eta_{n-1}+\eta_n)\delta_a{}^d \\ \frac{1}{2}(-i\eta_{n-1}+\eta_n)\delta^c{}_b & -\eta_\alpha{}^{cd} \end{pmatrix}.
\end{aligned} \tag{9.17}$$

3. Пусть нам известны связующие операторы $\eta_\alpha{}^{ab}$ для n mod 8 =2. Тогда связующие операторы $\eta_\Lambda{}^{AB}$ для n mod 8 =4 строятся по схеме

$$\begin{aligned}
\eta_\Lambda{}^{AB} &= \begin{pmatrix} \eta_\alpha{}^{ab} & \frac{1}{2}(\eta_{n-1}+i\eta_n)\delta^a{}_d \\ \frac{1}{2}(-\eta_{n-1}+i\eta_n)\delta_c{}^b & (\eta^T)_{\alpha cd} \end{pmatrix}, \\
\sigma_{\Lambda AB} &= \begin{pmatrix} (\eta^T)_{\alpha ab} & -\frac{1}{2}(\eta_{n-1}+i\eta_n)\delta_a{}^d \\ -\frac{1}{2}(-\eta_{n-1}+i\eta_n)\delta^c{}_b & \eta_\alpha{}^{cd} \end{pmatrix},
\end{aligned} \tag{9.18}$$

обеспечивая для n mod 8 = 4 наличие метрического кососимметрического спинора

$$\varepsilon_{AB} := (\varepsilon_1)_{AC}(\varepsilon_2)^{DK}(\varepsilon_3)_{BK} = \begin{pmatrix} 0 & 0 & 0 & i(\varepsilon_1)_{ab} \\ 0 & 0 & -i(\varepsilon_1)^{cd} & 0 \\ 0 & i(\varepsilon_1)^{kl} & 0 & 0 \\ -i(\varepsilon_1)_{mn} & 0 & 0 & 0 \end{pmatrix}, \tag{9.19}$$

где $(\varepsilon_1)_{ab}$ - метрический симметричный спинор для n mod 8=0.

4. Пусть нам известны связующие операторы $\eta_\alpha{}^{ab}$ для n mod 8 =4. Тогда связующие операторы $\eta_\Lambda{}^{AB}$ для n mod 8 =6 строятся по схеме

$$\begin{aligned}
\eta_\Lambda{}^{AB} &= \begin{pmatrix} \eta_\alpha{}^{ab} & \frac{1}{2}(\eta_{n-1}+i\eta_n)\delta^a{}_d \\ \frac{1}{2}(-\eta_{n-1}+i\eta_n)\delta_c{}^b & (\eta^T)_{\alpha cd} \end{pmatrix}, \\
\sigma_{\Lambda AB} &= \begin{pmatrix} (\eta^T)_{\alpha ab} & -\frac{1}{2}(\eta_{n-1}+i\eta_n)\delta_a{}^d \\ -\frac{1}{2}(-\eta_{n-1}+i\eta_n)\delta^c{}_b & \eta_\alpha{}^{cd} \end{pmatrix}.
\end{aligned} \tag{9.20}$$

Кроме того, дополнительно проведем преобразование связующих операторов по схеме

$$\tilde{\eta}_\Lambda{}^{AC} := \eta_\Lambda{}^{AB}\varepsilon_B{}^C, \quad \varepsilon_B{}^C := \begin{pmatrix} 0 & -i\varepsilon_{bm} \\ i\varepsilon^{dk} & 0 \end{pmatrix}, \tag{9.20$'$}$$

где ε_{ab} $(n/2+q=3)$ для n mod 8 = 4 будет метрическим кососимметрическим спинором. Тогда для следующей реализации n mod 8 = 6 операторы $\tilde{\eta}_\Lambda{}^{AC} = -\tilde{\eta}_\Lambda{}^{CA}$ будут полностью кососимметричны $(n/2+q)=0$. Легко проверить выполнение уравнения Клиффорда (8.1) для таких операторов. Если дальнейшее построение для n mod 8 = 0 вести на основании уже таких операторов, тогда будет существовать симметричный метрический спинор $\varepsilon_{AB} \equiv (\varepsilon_1)_{AB}$, $n/2+q=1$. Для n mod 8 = 2 уже будет существовать инволюция при $n/2+q=2$. И, снова, для n mod 8 = 4 получаем $n/2+q=3$. Круг замкнулся. Начало же положено в примерах 8.1-8.2.

5. *Пусть имеется алгебра над $\mathbb{R}^n$ со структурными константами, сгенерированными от связующих операторов $\eta_\Lambda{}^{AB}$ с метрическим спинором ε^{XZ} и оператором вложения $H_i{}^\Lambda$. Будем считать, что метрический тензор $g_{\Lambda\Psi}$ на главной диагонали содержит одни $\ll + \gg$. Тогда можно построить кососимметричные связующие операторы для пространства $\mathbb{C}^{n+6}$*

$$\eta_\alpha{}^{ab} = -\eta_\alpha{}^{ba} :=$$
$$\begin{pmatrix}
0 & 0 & 0 & \xi\varepsilon^{AQ} & 0 & \gamma\varepsilon^{AK} & -\alpha\varepsilon^{AD} & \eta_\Lambda{}^A{}_B \\
0 & 0 & -\xi\varepsilon_{CR} & 0 & -\gamma\varepsilon_{CM} & 0 & (\eta^T)_{\Lambda C}{}^D & \beta\varepsilon_{CB} \\
0 & \xi\varepsilon_{NY} & 0 & 0 & -\alpha\varepsilon_{NM} & (\eta^T)_{\Lambda N}{}^K & 0 & \delta\varepsilon_{NB} \\
-\xi\varepsilon^{LZ} & 0 & 0 & 0 & \eta_\Lambda{}^L{}_M & \beta\varepsilon^{LK} & -\delta\varepsilon^{LD} & 0 \\
0 & \gamma\varepsilon_{PY} & \alpha\varepsilon_{PR} & -(\eta^T)_{\Lambda P}{}^Q & 0 & 0 & 0 & -\zeta\varepsilon_{PB} \\
-\gamma\varepsilon^{SZ} & 0 & -\eta_\Lambda{}^S{}_R & -\beta\varepsilon^{SQ} & 0 & 0 & \zeta\varepsilon^{SD} & 0 \\
\alpha\varepsilon^{XZ} & -\eta_\Lambda{}^X{}_Y & 0 & \delta\varepsilon^{XQ} & 0 & -\zeta\varepsilon^{XK} & 0 & 0 \\
-(\eta^T)_{\Lambda T}{}^Z & -\beta\varepsilon_{TY} & -\delta\varepsilon_{TR} & 0 & \zeta\varepsilon_{TM} & 0 & 0 & 0
\end{pmatrix},$$

$$\begin{gathered}
\alpha := \tfrac{1}{2}(i\eta_{n+1} + \eta_{n+2}), \quad \beta := \tfrac{1}{2}(-i\eta_{n+1} + \eta_{n+2}), \\
\gamma := \tfrac{1}{2}(\eta_{n+3} + i\eta_{n+4}), \quad \delta := \tfrac{1}{2}(-\eta_{n+3} + i\eta_{n+4}), \\
\xi := \tfrac{1}{2}(\eta_{n+5} + i\eta_{n+6}), \quad \zeta := \tfrac{1}{2}(-\eta_{n+5} + i\eta_{n+6}).
\end{gathered} \tag{9.21}$$

6. Переход к связующим операторам пространства $\mathbb{C}^{n+8}$ осуществляется так:

$$\eta_\Lambda{}^{AB} := \begin{pmatrix} \eta_\alpha{}^{ab} & \phi\delta^a{}_d \\ \psi\delta_c{}^b & -(\eta^T)_{\alpha cd} \end{pmatrix}, \tag{9.22}$$

$$\phi := \tfrac{1}{2}(i\eta_{n+7} + \eta_{n+8}), \quad \psi := \tfrac{1}{2}(-i\eta_{n+7} + \eta_{n+8})$$

с метрическим спинором $\varepsilon^{XZ} := \begin{pmatrix} 0 & \delta^a{}_d \\ \delta_c{}^b & 0 \end{pmatrix}$, $\varepsilon_{XZ} := \begin{pmatrix} 0 & \delta_a{}^d \\ \delta^c{}_b & 0 \end{pmatrix}$. *После этого, можно перейти к связующим операторам пространства* $\mathbb{R}^{n+8} \subset \mathbb{C}^{n+8}$ *с помощью соответствующего оператора вложения. А уже такие операторы генерируют структурные константы алгебры размерности* $n+8$.

10 Инфинитизимальные преобразования

> *В этом параграфе рассказывается о том, как по связующим операторам, удовлетворяющим уравнению Клиффорда, построить инфинитизимальные преобразования. Вывод всех результатов этого параграфа сделан на основе [22, т. 1, с. 224-226].*

Согласно следствию 8.3 для собственных ортогональных преобразований верно разложение

$$\begin{aligned} &1).\quad S_\Lambda{}^\Psi \eta_\Psi{}^{AB} = -\eta_\Lambda{}^{CD} S_C{}^A E_D{}^K S_K{}^M E_M{}^B, \\ &2).\quad S_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \eta_\Lambda{}^{CD} S_C{}^A (\eta_I)^\Omega S_\Omega{}^\Phi \eta_\Phi{}^{KB} \varepsilon_{KL} S_X{}^L \varepsilon^{MX} (\varepsilon_I)_{MD}. \end{aligned} \tag{10.1}$$

Пусть имеется $S_\Lambda{}^\Psi(\lambda)$ - однопараметрическое семейство ортогональных преобразований

$$S_\Lambda{}^\Psi(\lambda) S_\Omega{}^\Phi(\lambda) g_{\Psi\Phi} = g_{\Lambda\Omega}, \quad S_\Lambda{}^\Psi(0) := \delta_\Lambda{}^\Psi, \tag{10.2}$$

что индуцирует однопараметрические преобразования в спинорном пространстве

$$S_C{}^A(\lambda), \quad S_C{}^A(0) = \delta_C{}^A. \tag{10.3}$$

Соответствующие инфинитизимальные преобразования определятся как

$$\begin{aligned} T_\Lambda{}^\Psi &= \left[\tfrac{d}{d\lambda} S_\Lambda{}^\Psi(\lambda)\right]\Big|_{\lambda=0}, & &T_{\Lambda\Psi} = -T_{\Psi\Lambda}, \\ & & &1).\quad T_C{}^A = -E_C{}^B T_B{}^D E_D{}^A, \\ T_C{}^A &= \left[\tfrac{d}{d\lambda} S_C{}^A(\lambda)\right]\Big|_{\lambda=0}, & & \\ & & &2).\quad T_C{}^A = -T^A{}_C. \end{aligned} \tag{10.4}$$

Дифференцируя в нуле (10.1), получим тождество (выкладки в приложении)

$$T_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \eta_\Lambda{}^{CB} T_C{}^A - \eta_\Lambda{}^{AD} (\varepsilon_I)^{KB} T_K{}^M (\varepsilon_I)_{MD} + \eta_\Lambda{}^{AD} (\varepsilon_I)_{MD} (\eta_I)^\Omega T_\Omega{}^\Phi \eta_\Phi{}^{MB}. \tag{10.5}$$

Частным решением этого уравнения будет

$$T_A{}^C = \frac{1}{2} T^{\Theta\Phi} \eta_\Phi{}^{AB} \eta_{\Theta CB}, \tag{10.6}$$

что проверяется прямой подстановкой. Однородное же решение уравнения

$$\eta_\Lambda{}^{CB} T_C{}^A - \eta_\Lambda{}^{AD} (\varepsilon_I)^{KB} T_K{}^M (\varepsilon_I)_{MD} = 0 \tag{10.7}$$

перепишется как (8.13), что означает тривиальность такого решения.

11 Комплексная и вещественная связности

В этом параграфе рассказывается о том, как по частным решениям уравнения Клиффорда построить продолжение римановой связности без кручения в спинорное расслоение. Рассматривается два различных варианта. Это даст возможность построить спинорные аналоги операторов Ли. Вывод всех результатов этого параграфа сделан на основе [9], [22, т. 1, с. 282-285].

Для вещественной реализации касательного расслоения $\tau^{\mathbb{R}}(V_{2n})$ уравнения геодезических примут вид

$$\frac{d^2 x^{\tilde{\varOmega}}}{ds^2} + \Gamma_{\tilde{\varPhi}\tilde{\varPsi}}{}^{\tilde{\varOmega}} \frac{dx^{\tilde{\varPhi}}}{ds} \frac{dx^{\tilde{\varPsi}}}{ds} = 0. \tag{11.1}$$

Для параметризации атласа $w^\Theta = \frac{1}{\sqrt{2}}(u^\Theta(x^{\tilde{\varTheta}}) + iv^\Theta(x^{\tilde{\varTheta}}))$

$$\begin{gathered}
\frac{d^2 x^{\tilde{\varOmega}}}{ds^2} = \frac{d}{ds}\Big(\frac{\partial x^{\tilde{\varOmega}}}{\partial w^\Theta} \frac{dw^\Theta}{ds} + \frac{\partial x^{\tilde{\varOmega}}}{\partial w^{\Theta'}} \frac{dw^{\Theta'}}{ds}\Big) = \\
\frac{\partial^2 x^{\tilde{\varOmega}}}{\partial w^\Theta ds} \frac{dw^\Theta}{ds} + \frac{\partial x^{\tilde{\varOmega}}}{\partial w^\Theta} \frac{d^2 w^\Theta}{ds^2} + \frac{\partial^2 x^{\tilde{\varOmega}}}{\partial w^{\Theta'} ds} \frac{dw^{\Theta'}}{ds} + \frac{\partial x^{\tilde{\varOmega}}}{\partial w^{\Theta'}} \frac{d^2 w^{\Theta'}}{ds^2}, \\
\frac{d^2 w^\Theta}{ds^2} \frac{\partial x^{\tilde{\varOmega}}}{\partial w^\Theta} + \frac{d^2 w^{\Theta'}}{ds^2} \frac{\partial x^{\tilde{\varOmega}}}{\partial w^{\Theta'}} + \frac{\partial^2 x^{\tilde{\varOmega}}}{\partial w^\Theta ds} \frac{dw^\Theta}{ds} + \frac{\partial^2 x^{\tilde{\varOmega}}}{\partial w^{\Theta'} ds} \frac{dw^{\Theta'}}{ds} + \\
+ \Gamma_{\tilde{\varPhi}\tilde{\varPsi}}{}^{\tilde{\varOmega}} \frac{dx^{\tilde{\varPhi}}}{ds} \frac{\partial x^{\tilde{\varPsi}}}{\partial w^\Theta} \frac{dw^\Theta}{ds} + \Gamma_{\tilde{\varPhi}\tilde{\varPsi}}{}^{\tilde{\varOmega}} \frac{dx^{\tilde{\varPhi}}}{ds} \frac{\partial x^{\tilde{\varPsi}}}{\partial w^{\Theta'}} \frac{dw^{\Theta'}}{ds} = 0.
\end{gathered} \tag{11.2}$$

Домножим обе части на $\frac{dw^\Lambda}{dx^{\tilde{\varOmega}}} = m^\Lambda{}_{\tilde{\varOmega}}$. Учитывая тождества (3.8) для операторов $m^\Lambda{}_{\tilde{\varOmega}}$ и $m^{\Lambda'}{}_{\tilde{\varOmega}}$, получим

$$\begin{gathered}
\Big(\frac{d^2 w^\Lambda}{ds^2} + \frac{\partial w^\Lambda}{\partial x^{\tilde{\varOmega}}} \frac{\partial^2 x^{\tilde{\varOmega}}}{\partial w^\Theta ds} \frac{dw^\Theta}{ds} + \Gamma_{\tilde{\varPhi}\tilde{\varPsi}}{}^{\tilde{\varOmega}} \frac{\partial x^{\tilde{\varPsi}}}{\partial w^\Theta} \frac{dx^{\tilde{\varPhi}}}{ds} \frac{dw^\Theta}{ds} \frac{\partial w^\Lambda}{\partial x^{\tilde{\varOmega}}}\Big) + \\
+ \Big(\frac{\partial w^\Lambda}{\partial x^{\tilde{\varOmega}}} \frac{\partial^2 x^{\tilde{\varOmega}}}{\partial w^{\Theta'} ds} \frac{dw^{\Theta'}}{ds} + \Gamma_{\tilde{\varPhi}\tilde{\varPsi}}{}^{\tilde{\varOmega}} \frac{\partial x^{\tilde{\varPsi}}}{\partial w^{\Theta'}} \frac{dx^{\tilde{\varPhi}}}{ds} \frac{dw^{\Theta'}}{ds} \frac{\partial w^\Lambda}{\partial x^{\tilde{\varOmega}}}\Big) = 0.
\end{gathered} \tag{11.3}$$

Для комплексной реализации касательного расслоения $\tau^{\mathbb{C}}(V_{2n})$ уравнения геодезических примут вид

$$\frac{d^2 w^\Lambda}{ds^2} + \Big(\Gamma_{\tilde{\varPhi}\Theta}{}^\Lambda \frac{dw^\Theta}{ds} + \Gamma_{\tilde{\varPhi}\Theta'}{}^\Lambda \frac{dw^{\Theta'}}{ds}\Big) \frac{dx^{\tilde{\varPhi}}}{ds} = 0. \tag{11.4}$$

Сравнивая эти два уравнения и определение (3.8) операторов $m^\Lambda{}_{\tilde{\varOmega}}$, $m_\Lambda{}^{\tilde{\varOmega}}$, получим

$$\begin{cases}
m^\Lambda{}_{\tilde{\varOmega}} \, (\partial_{\tilde{\varPhi}} m_\Theta{}^{\tilde{\varOmega}} + \Gamma_{\tilde{\varPhi}\tilde{\varPsi}}{}^{\tilde{\varOmega}} m_\Theta{}^{\tilde{\varPsi}}) = \Gamma_{\tilde{\varPhi}\Theta}{}^\Lambda, \\
m^\Lambda{}_{\tilde{\varOmega}} \, (\partial_{\tilde{\varPhi}} \bar{m}_{\Theta'}{}^{\tilde{\varOmega}} + \Gamma_{\tilde{\varPhi}\tilde{\varPsi}}{}^{\tilde{\varOmega}} \bar{m}_{\Theta'}{}^{\tilde{\varPsi}}) = \Gamma_{\tilde{\varPhi}\Theta'}{}^\Lambda,
\end{cases} \tag{11.5}$$

принимая во внимание тождество

$$\frac{\partial^2 x^{\tilde{\varOmega}}}{\partial w^{\Theta} ds} = \frac{\partial^2 x^{\tilde{\varOmega}}}{\partial w^{\Theta} \partial x^{\tilde{\varPhi}}} \frac{dx^{\tilde{\varPhi}}}{ds} = \frac{dx^{\tilde{\varPhi}}}{ds} \partial_{\tilde{\varPhi}} \frac{\partial x^{\tilde{\varOmega}}}{\partial w^{\Theta}} = \frac{dx^{\tilde{\varPhi}}}{ds} \partial_{\tilde{\varPhi}} {m_{\Theta}}^{\tilde{\varOmega}}. \tag{11.6}$$

Другая запись этой системы такая

$$\begin{cases} {m_{\Xi}}^{\tilde{\varPsi}} \ (-\partial_{\tilde{\varPhi}} {m^{\Theta}}_{\tilde{\varPsi}} + {\Gamma_{\tilde{\varPhi}\tilde{\varPsi}}}^{\tilde{\varOmega}} {m^{\Theta}}_{\tilde{\varOmega}}) = {\Gamma_{\tilde{\varPhi}\Xi}}^{\Theta} \ \text{, домножим на } \ {m^{\Xi}}_{\tilde{\varLambda}}, \\ {\bar{m}_{\Xi'}}^{\tilde{\varPsi}} \ (-\partial_{\tilde{\varPhi}} {m^{\Theta}}_{\tilde{\varPsi}} + {\Gamma_{\tilde{\varPhi}\tilde{\varPsi}}}^{\tilde{\varOmega}} {m^{\Theta}}_{\tilde{\varOmega}}) = {\Gamma_{\tilde{\varPhi}\Xi'}}^{\Theta} \ \text{, домножим на } \ {\bar{m}^{\Xi'}}_{\tilde{\varLambda}}, \end{cases} \\ \begin{cases} {\triangle_{\tilde{\varLambda}}}^{\tilde{\varPsi}} \ (-\partial_{\tilde{\varPhi}} {m^{\Lambda}}_{\tilde{\varPsi}} + {\Gamma_{\tilde{\varPhi}\tilde{\varPsi}}}^{\tilde{\varOmega}} {m^{\Lambda}}_{\tilde{\varOmega}}) = {\Gamma_{\tilde{\varPhi}\Xi}}^{\Lambda} {m^{\Xi}}_{\tilde{\varLambda}}, \\ {\bar{\triangle}_{\tilde{\varLambda}}}^{\tilde{\varPsi}} \ (-\partial_{\tilde{\varPhi}} {m^{\Lambda}}_{\tilde{\varPsi}} + {\Gamma_{\tilde{\varPhi}\tilde{\varPsi}}}^{\tilde{\varOmega}} {m^{\Lambda}}_{\tilde{\varOmega}}) = {\Gamma_{\tilde{\varPhi}\Xi'}}^{\Lambda} {\bar{m}^{\Xi'}}_{\tilde{\varLambda}}. \end{cases} \tag{11.7}$$

Сложим эти два уравнения

$$\partial_{\tilde{\varPhi}} {m^{\Lambda}}_{\tilde{\varLambda}} - {\Gamma_{\tilde{\varPhi}\tilde{\varLambda}}}^{\tilde{\varOmega}} {m^{\Lambda}}_{\tilde{\varOmega}} + {\Gamma_{\tilde{\varPhi}\Xi}}^{\Lambda} {m^{\Xi}}_{\tilde{\varLambda}} + {\Gamma_{\tilde{\varPhi}\Xi'}}^{\Lambda} {\bar{m}^{\Xi'}}_{\tilde{\varLambda}} = 0. \tag{11.8}$$

Свернем его с ${\bar{m}_{\Psi'}}^{\tilde{\varLambda}}$

$$\begin{gathered} {\bar{m}_{\Psi'}}^{\tilde{\varLambda}} \partial_{\tilde{\varPhi}} {m^{\Lambda}}_{\tilde{\varLambda}} - {\Gamma_{\tilde{\varPhi}\tilde{\varLambda}}}^{\tilde{\varOmega}} {m^{\Lambda}}_{\tilde{\varOmega}} {\bar{m}_{\Psi'}}^{\tilde{\varLambda}} + {\Gamma_{\tilde{\varPhi}\Psi'}}^{\Lambda} = 0, \\ {\Gamma_{\tilde{\varPhi}\Psi'}}^{\Lambda} = {\Gamma_{\tilde{\varPhi}\tilde{\varLambda}}}^{\tilde{\varOmega}} {m^{\Lambda}}_{\tilde{\varOmega}} {\bar{m}_{\Psi'}}^{\tilde{\varLambda}} - \tfrac{i}{2} {\bar{m}_{\Psi'}}^{\tilde{\varLambda}} {m^{\Lambda}}_{\tilde{\varTheta}} \partial_{\tilde{\varPhi}} {f_{\tilde{\varLambda}}}^{\tilde{\varTheta}}. \end{gathered} \tag{11.9}$$

Поэтому связность в касательном расслоении необходимо доопределить системой

$$\begin{cases} \nabla_{\tilde{\varPhi}} {m^{\Lambda}}_{\tilde{\varLambda}} = 0, \\ \\ \nabla_{\tilde{\varPhi}} {\bar{m}^{\Lambda'}}_{\tilde{\varLambda}} = 0. \end{cases} \tag{11.10}$$

Действительно, поскольку ${m^{\Psi}}_{\tilde{\varOmega}} = i \ {f_{\tilde{\varOmega}}}^{\tilde{\varLambda}} {m^{\Psi}}_{\tilde{\varLambda}}$, то

$$\begin{gathered} \nabla_{\tilde{\varPhi}} {f_{\tilde{\varOmega}}}^{\tilde{\varLambda}} = 0, \\ \partial_{\tilde{\varPhi}} {f_{\tilde{\varOmega}}}^{\tilde{\varLambda}} = ({\Gamma_{\tilde{\varPhi}\tilde{\varOmega}}}^{\tilde{\varTheta}} {f_{\tilde{\varTheta}}}^{\tilde{\varLambda}} - {\Gamma_{\tilde{\varPhi}\tilde{\varTheta}}}^{\tilde{\varLambda}} {f_{\tilde{\varOmega}}}^{\tilde{\varTheta}}). \end{gathered} \tag{11.11}$$

Домножим это уравнение на $\tfrac{i}{2} {\bar{m}_{\Psi'}}^{\tilde{\varOmega}} {m^{\Lambda}}_{\tilde{\varLambda}}$

$$\frac{i}{2} {\bar{m}_{\Psi'}}^{\tilde{\varOmega}} {m^{\Lambda}}_{\tilde{\varLambda}} \partial_{\tilde{\varPhi}} {f_{\tilde{\varOmega}}}^{\tilde{\varLambda}} = {\bar{m}_{\Psi'}}^{\tilde{\varOmega}} {m^{\Lambda}}_{\tilde{\varLambda}} {\Gamma_{\tilde{\varPhi}\tilde{\varOmega}}}^{\tilde{\varLambda}}. \tag{11.12}$$

Отсюда

$${\Gamma_{\tilde{\varPhi}\Psi'}}^{\Lambda} = 0. \tag{11.13}$$

Поэтому можно записать

$$\nabla_{\tilde{\varPhi}} {m^{\Lambda}}_{\tilde{\varLambda}} := \partial_{\tilde{\varPhi}} {m^{\Lambda}}_{\tilde{\varLambda}} - {\Gamma_{\tilde{\varPhi}\tilde{\varLambda}}}^{\tilde{\varOmega}} {m^{\Lambda}}_{\tilde{\varOmega}} + {\Gamma_{\tilde{\varPhi}\Psi}}^{\Lambda} {m^{\Psi}}_{\tilde{\varLambda}} = 0. \tag{11.14}$$

Таким образом, если известны коэффициенты связности в одной из реализации (вещественной или комплексной), то по этому уравнению однозначно восстанавливаются коэффициенты связности в другой реализации (комплексной или вещественной соответственно).

Если теперь в вещественной реализации касательного расслоения задана связность, согласованная с метрикой

$$\nabla_{\tilde{\Lambda}} G_{\tilde{\Psi}\tilde{\Omega}} = 0, \tag{11.15}$$

то в спинорное расслоение ее можно продолжить условием

$$\nabla_{\tilde{\Lambda}} \eta_{\tilde{\Psi}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}} = 0. \tag{11.16}$$

Пусть теперь I). n mod 4 =0, n>2, II). n mod 4 =2, n>4, тогда

$$\begin{gathered}
I). \qquad \nabla_{\tilde{\Psi}} \eta_{\tilde{\Lambda}\tilde{\tilde{C}}\tilde{\tilde{D}}} = 0, \\
\nabla_{\tilde{\Psi}} \varepsilon_{\tilde{\tilde{A}}\tilde{\tilde{C}}} \varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}} = 0, \\
\varepsilon_{\tilde{\tilde{A}}\tilde{\tilde{C}}} \nabla_{\tilde{\Psi}} \varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}} + \varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}} \nabla_{\tilde{\Psi}} \varepsilon_{\tilde{\tilde{A}}\tilde{\tilde{C}}} = 0, \\
\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}} (\delta_{\tilde{\tilde{A}}}{}^{\tilde{\tilde{K}}} \varepsilon^{\tilde{\tilde{L}}\tilde{\tilde{D}}} \nabla_{\tilde{\Psi}} \varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}} + \delta_{\tilde{\tilde{B}}}{}^{\tilde{\tilde{L}}} \varepsilon^{\tilde{\tilde{K}}\tilde{\tilde{C}}} \nabla_{\tilde{\Psi}} \varepsilon_{\tilde{\tilde{A}}\tilde{\tilde{C}}}) = 0, \\
\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{K}}\tilde{\tilde{B}}} \varepsilon^{\tilde{\tilde{L}}\tilde{\tilde{D}}} \nabla_{\tilde{\Psi}} \varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}} + \eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{B}}\tilde{\tilde{L}}} \varepsilon^{\tilde{\tilde{K}}\tilde{\tilde{C}}} \nabla_{\tilde{\Psi}} \varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{C}}} = 0, \\
II). \qquad \nabla_{\tilde{\Psi}} \eta_{\tilde{\Lambda}\tilde{\tilde{C}}\tilde{\tilde{D}}} = 0, \\
\tilde{\nabla}_{\tilde{\Psi}} E_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{A}}} E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} = 0, \\
E_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{A}}} \tilde{\nabla}_{\tilde{\Psi}} E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} + E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} \tilde{\nabla}_{\tilde{\Psi}} E_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{A}}} = 0, \\
\eta_{\tilde{\Lambda}\tilde{\tilde{A}}\tilde{\tilde{B}}} (\delta_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}} E_{\tilde{\tilde{L}}}{}^{\tilde{\tilde{C}}} \nabla_{\tilde{\Psi}} E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} + \delta_{\tilde{\tilde{L}}}{}^{\tilde{\tilde{B}}} E_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{D}}} \nabla_{\tilde{\Psi}} E_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{A}}}) = 0, \\
\eta_{\tilde{\Lambda}\tilde{\tilde{K}}\tilde{\tilde{B}}} E_{\tilde{\tilde{L}}}{}^{\tilde{\tilde{C}}} \nabla_{\tilde{\Psi}} E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} + \eta_{\tilde{\Lambda}\tilde{\tilde{A}}\tilde{\tilde{L}}} E_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{D}}} \nabla_{\tilde{\Psi}} E_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{A}}} = 0.
\end{gathered} \tag{11.17}$$

Обратимся к уравнению (8.13). Это даст

$$I). \quad \varepsilon^{\tilde{\tilde{L}}\tilde{\tilde{D}}} \nabla_{\tilde{\Psi}} \varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}} = 0, \qquad II). \quad E_{\tilde{\tilde{L}}}{}^{\tilde{\tilde{C}}} \nabla_{\tilde{\Psi}} E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} = 0. \tag{11.18}$$

Таким образом, получаем, что

$$\begin{aligned}
I). \quad & \nabla_{\tilde{\Psi}} \varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}} = 0, & II). \quad & \nabla_{\tilde{\Psi}} E_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} = 0, \\
& \nabla_{\tilde{\Psi}} \tilde{\varepsilon}_{\tilde{\tilde{B}}\tilde{\tilde{D}}} = 0, & & \nabla_{\tilde{\Psi}} \tilde{E}_{\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} = 0.
\end{aligned} \tag{11.19}$$

Поэтому, условием (11.16) налагаются ограничения

$$\begin{aligned}
I). \quad & \nabla_{\tilde{\Psi}} \eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}} \varepsilon_{\tilde{\tilde{A}}\tilde{\tilde{B}}} = 0, & II). \quad & \nabla_{\tilde{\Psi}} \eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}} \eta_{\tilde{\Phi}\tilde{\tilde{A}}\tilde{\tilde{C}}} E_{\tilde{\tilde{B}}}{}^{\tilde{\tilde{C}}} = 0, \\
& \nabla_{\tilde{\Psi}} \eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}} \tilde{\varepsilon}_{\tilde{\tilde{A}}\tilde{\tilde{B}}} = 0, & & \nabla_{\tilde{\Psi}} \eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}} \eta_{\tilde{\Phi}\tilde{\tilde{A}}\tilde{\tilde{C}}} \tilde{E}_{\tilde{\tilde{B}}}{}^{\tilde{\tilde{C}}} = 0.
\end{aligned} \tag{11.20}$$

Такой вид связности возможен при нормировке спинорного базиса согласно [22, т. 1, с. 284]. Однако, можно ослабить условия (11.15) и (11.16), вспомнив, что мы рассматриваем связность в касательном расслоении. Следовательно, необходимо туда и вернуться, получая соотношение

$$\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{B}}} \nabla_{\tilde{\Lambda}} \eta_{\tilde{\Omega}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}} = 0, \tag{11.21}$$

которое расписывается как

$$\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{B}}}\partial_{\tilde{\Lambda}}\eta_{\tilde{\Omega}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}} - \Gamma_{\tilde{\Lambda}\tilde{\Omega}}{}^{\tilde{\Theta}}\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta_{\tilde{\Theta}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}} + \Gamma_{\tilde{\Lambda}\tilde{\tilde{D}}}{}^{\tilde{\tilde{A}}}\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta_{\tilde{\Omega}}{}^{\tilde{\tilde{D}}\tilde{\tilde{B}}} + \Gamma_{\tilde{\Lambda}\tilde{\tilde{D}}}{}^{\tilde{\tilde{B}}}\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta_{\tilde{\Omega}}{}^{\tilde{\tilde{A}}\tilde{\tilde{D}}} = 0. \tag{11.22}$$

Поэтому для того, чтобы однозначно продолжить построенную связность в спинорное расслоение, необходимо потребовать

$$\begin{gathered}\Gamma_{\tilde{\Lambda}\tilde{\tilde{L}}}{}^{\tilde{\tilde{M}}} := \tfrac{1}{N^2}\Gamma_{\tilde{\Lambda}\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}(\eta_{\tilde{\Psi}\tilde{\tilde{L}}\tilde{\tilde{B}}}\eta_{\tilde{\Omega}}{}^{\tilde{\tilde{M}}\tilde{\tilde{B}}}\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{C}}}\eta^{\tilde{\Omega}\tilde{\tilde{K}}\tilde{\tilde{C}}} - \tfrac{(2n-2)}{4}\delta_{\tilde{\tilde{L}}}{}^{\tilde{\tilde{M}}}\delta_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}),\\ \Gamma_{\tilde{\Lambda}\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}} := -\tfrac{2}{2n-4}\Gamma_{\tilde{\Lambda}\tilde{\tilde{C}}}{}^{\tilde{\tilde{D}}}(\varepsilon_{\tilde{\tilde{D}}\tilde{\tilde{K}}}{}^{\tilde{\tilde{C}}\tilde{\tilde{A}}} - \tfrac{2n-2}{2N^2}\delta_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}\delta_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{C}}}).\end{gathered} \tag{11.23}$$

Действительно,

$$\begin{gathered}\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta_{\tilde{\Psi}\tilde{\tilde{K}}\tilde{\tilde{B}}}\cdot(-\tfrac{2}{2n-4})(\varepsilon_{\tilde{\tilde{D}}\tilde{\tilde{A}}}{}^{\tilde{\tilde{C}}\tilde{\tilde{K}}} - \tfrac{2n-2}{2N^2}\delta_{\tilde{\tilde{A}}}{}^{\tilde{\tilde{K}}}\delta_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{C}}}) = \eta_{[\tilde{\Lambda}}{}^{\tilde{\tilde{B}}\tilde{\tilde{C}}}\eta_{\tilde{\Psi}]\tilde{\tilde{B}}\tilde{\tilde{D}}} -\\ -\tfrac{1}{2n-4}(\tfrac{2n}{2} - \tfrac{2n-2}{2N^2}\cdot 2N^2)g_{\tilde{\Lambda}\tilde{\Psi}}\delta_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{C}}} = \eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{B}}\tilde{\tilde{C}}}\eta_{\tilde{\Psi}\tilde{\tilde{B}}\tilde{\tilde{D}}}.\end{gathered} \tag{11.24}$$

Замечание 11.1. *Следует отметить, что при n=4 все выводы остаются верными, а последнее тождество примет вид*

$$\begin{gathered}\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{C}}}\eta_{\tilde{\Psi}\tilde{\tilde{K}}\tilde{\tilde{C}}} = (\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{C}}\tilde{\tilde{A}}} + \eta_{\tilde{\Lambda}}\varepsilon^{\tilde{\tilde{A}}\tilde{\tilde{C}}})\eta_{\tilde{\Psi}\tilde{\tilde{K}}\tilde{\tilde{C}}} = \eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{C}}\tilde{\tilde{A}}}(\eta_{\tilde{\Psi}\tilde{\tilde{C}}\tilde{\tilde{K}}} + \eta_{\tilde{\Psi}}\varepsilon_{\tilde{\tilde{K}}\tilde{\tilde{C}}}) + \eta_{\tilde{\Lambda}}\eta_{\tilde{\Psi}\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}} =\\ = \eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{C}}\tilde{\tilde{A}}}\eta_{\tilde{\Psi}\tilde{\tilde{C}}\tilde{\tilde{K}}} - \eta_{\tilde{\Psi}}\eta_{\tilde{\Lambda}\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}} + \eta_{\tilde{\Lambda}}\eta_{\tilde{\Psi}\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}.\end{gathered} \tag{11.25}$$

Таким образом, свертка (11.22) с $\eta_{\tilde{\Psi}}\eta^{\tilde{\Omega}}$ *определит выражение* $\Gamma_{\tilde{\Lambda}\tilde{\tilde{A}}}{}^{\tilde{\tilde{A}}}$*, после подстановки которого в (11.22) однозначно определятся коэффициенты* $\Gamma_{\tilde{\Lambda}\tilde{\tilde{A}}}{}^{\tilde{\tilde{D}}}$*. Для этого достаточно свернуть (11.22) с* $\eta_{\tilde{\Psi}\tilde{\tilde{K}}\tilde{\tilde{L}}}\eta^{\tilde{\Omega}\tilde{\tilde{M}}\tilde{\tilde{L}}}$*. Это обусловлено тождеством* $\eta_{\Psi\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta^{\Psi}{}_{\tilde{\tilde{C}}\tilde{\tilde{D}}} = \varepsilon_{\tilde{\tilde{A}}\tilde{\tilde{B}}\tilde{\tilde{C}}\tilde{\tilde{D}}} = \varepsilon_{\tilde{\tilde{A}}\tilde{\tilde{C}}}\varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}}$*, где* $\varepsilon_{\tilde{\tilde{B}}\tilde{\tilde{D}}}$ *- как и прежде, кососимметрический метрический спинор (см. пример 8.1, который легко адаптируется для вложения* $\mathbb{R}^4 \subset \mathbb{C}^4$*). Но в таком случае условие (11.21) равносильно (11.16).*

Если выполнить симметризацию выражения (11.22) по Ω и опущенному Ψ, то из аналога (8.17) получим

$$\underbrace{\frac{N^2}{2}(\partial_{\tilde{\Lambda}}G_{\tilde{\Psi}\tilde{\Omega}} - 2\Gamma_{\tilde{\Lambda}(\tilde{\Omega}\tilde{\Psi})})}_{=\frac{N^2}{2}\nabla_{\Lambda}G_{\tilde{\Psi}\tilde{\Omega}}} + \underbrace{(\frac{1}{2N^2}\eta_{\tilde{\Psi}\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta_{\tilde{\Omega}\tilde{\tilde{C}}\tilde{\tilde{D}}}\partial_{\Lambda}\varepsilon^{\tilde{\tilde{A}}\tilde{\tilde{B}}\tilde{\tilde{C}}\tilde{\tilde{D}}} + \Gamma_{\tilde{\Lambda}\tilde{\tilde{B}}}{}^{\tilde{\tilde{B}}}G_{\tilde{\Psi}\tilde{\Omega}})}_{=\frac{1}{2N^2}\eta_{\tilde{\Psi}\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta_{\tilde{\Omega}\tilde{\tilde{C}}\tilde{\tilde{D}}}\nabla_{\Lambda}\varepsilon^{\tilde{\tilde{A}}\tilde{\tilde{B}}\tilde{\tilde{C}}\tilde{\tilde{D}}}} = 0. \tag{11.26}$$

Если метрический тензор ковариантно постоянен, тогда

$$\eta_{\tilde{\Psi}\tilde{\tilde{A}}\tilde{\tilde{B}}}\eta_{\tilde{\Omega}\tilde{\tilde{C}}\tilde{\tilde{D}}}\nabla_{\tilde{\Lambda}}\varepsilon^{\tilde{\tilde{A}}\tilde{\tilde{B}}\tilde{\tilde{C}}\tilde{\tilde{D}}} = 0. \tag{11.27}$$

Соответственно, комплексная связность определится соотношениями

$$\nabla_{\Lambda} := m_{\Lambda}{}^{\tilde{\Lambda}}\nabla_{\tilde{\Lambda}}, \quad \bar{\nabla}_{\Lambda'} := \bar{m}_{\Lambda'}{}^{\tilde{\Lambda}}\nabla_{\tilde{\Lambda}}, \tag{11.28}$$

что приведет в отсутствии кручения к связности

$$\nabla_\Lambda g_{\Psi\Phi} = 0, \quad \bar{\nabla}_{\Lambda'} \bar{g}_{\Psi'\Phi'} = 0, \tag{11.29}$$

которая продолжается в комплексное спинорное расслоение условиями (см. (6.33))

$$\begin{gathered}\nabla_\Lambda (\tilde{M}_K)_A{}^{\tilde{\tilde{B}}} = 0, \quad \nabla_\Lambda (\tilde{\tilde{M}}_K)_A{}^{\tilde{\tilde{B}}} = 0, \quad \bar{\nabla}_{\Lambda'} (\tilde{M}_K)_A{}^{\tilde{\tilde{B}}} = 0, \quad \bar{\nabla}_{\Lambda'} (\tilde{\tilde{M}}_K)_A{}^{\tilde{\tilde{B}}} = 0, \\ \sum_{K=1}^{2N} (\eta_K)^\Psi{}_{AB} \bar{\nabla}_{\Lambda'} (\eta_K)_\Omega{}^{AB} = 0, \quad \sum_{K=1}^{2N} (\eta_K)^\Psi{}_{AB} \nabla_\Lambda (\eta_K)_\Omega{}^{AB} = 0.\end{gathered} \tag{11.30}$$

При этом следует учесть, что верхние соотношения

$$\begin{gathered}\partial_\Lambda (\tilde{M}_K)_A{}^{\tilde{\tilde{B}}} + \Gamma_{\Lambda\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} (\tilde{M}_K)_A{}^{\tilde{\tilde{C}}} - (\tilde{\Gamma}_K)_{\Lambda A}{}^C (\tilde{M}_K)_C{}^{\tilde{\tilde{B}}} = 0, \\ \partial_\Lambda (\tilde{\tilde{M}}_K)_A{}^{\tilde{\tilde{B}}} + \Gamma_{\Lambda\tilde{\tilde{C}}}{}^{\tilde{\tilde{B}}} (\tilde{\tilde{M}}_K)_A{}^{\tilde{\tilde{C}}} - (\tilde{\tilde{\Gamma}}_K)_{\Lambda A}{}^C (\tilde{\tilde{M}}_K)_C{}^{\tilde{\tilde{B}}} = 0\end{gathered} \tag{11.31}$$

определяют различные коэффициенты связности $(\tilde{\Gamma}_K)_{\Lambda A}{}^C$ и $(\tilde{\tilde{\Gamma}}_K)_{\Lambda A}{}^C$, а нижние соотношение расписывается как

$$\begin{gathered}\sum_{K=1}^{2N} ((\eta_K)^\Psi{}_{AB} \partial_\Lambda (\eta_K)_\Omega{}^{AB} - \tfrac{N}{2} \cdot \Gamma_{\Lambda\Omega\Psi} + \\ + (\tilde{\Gamma}_K)_{\Lambda D}{}^A (\eta_K)^\Psi{}_{AB} (\eta_K)_\Omega{}^{DB} + (\tilde{\tilde{\Gamma}}_K)_{\Lambda D}{}^B (\eta_K)^\Psi{}_{AB} (\eta_K)_\Omega{}^{AD}) = 0.\end{gathered} \tag{11.32}$$

Чтобы однозначно продолжить связность, можно положить

$$\begin{gathered}(\tilde{\tilde{\Gamma}}_K)_{\Lambda L}{}^M := \tfrac{2}{N} (\tilde{\tilde{\Gamma}}_K)_{\Lambda D}{}^A ((\eta_K)_{\Psi LB} (\eta_K)_\Omega{}^{MB} (\eta_K)^\Psi{}_{AC} (\eta_K)^{\Omega DC} - \tfrac{(n-2)}{4} \delta_L{}^M \delta_D{}^A), \\ (\tilde{\Gamma}_K)_{\Lambda A}{}^C := -\tfrac{2}{n-4} (\tilde{\tilde{\Gamma}}_K)_{\Lambda K}{}^L ((\varepsilon_K)_{LA}{}^{KC} - \tfrac{n-2}{N} \delta_L{}^K \delta_A{}^C).\end{gathered} \tag{11.33}$$

Поэтому, из ковариантного постоянства метрического тензора

$$\begin{gathered}\nabla_\Lambda g_{\Psi\Omega} = (\tfrac{2}{N})^2 \nabla_\Lambda ((\eta_K)_{\Psi AB} (\eta_K)_{\Omega CD} (\varepsilon_K)^{ABCD}) = \\ = \tfrac{2}{N} ((\eta_K)_\Psi{}^{AB} \nabla_\Lambda (\eta_K)_{\Omega AB} + (\eta_K)_\Omega{}^{AB} \nabla_\Lambda (\eta_K)_{\Psi AB}) + \\ + (\tfrac{2}{N})^2 (\eta_K)_{\Psi AB} (\eta_K)_{\Omega CD} \nabla_\Lambda (\varepsilon_K)^{ABCD} = 0\end{gathered} \tag{11.34}$$

будет следовать

$$\sum_{K=1}^{2N} (\eta_K)_{\Psi AB} (\eta_K)_{\Omega CD} \nabla_\Lambda (\varepsilon_K)^{ABCD} = 0. \tag{11.35}$$

Теперь обратимся к вещественным вложениям. Уравнение

$$\nabla_\Lambda H_i{}^\Psi = 0 \tag{11.36}$$

распишется как

$$\partial_\Lambda H_i{}^\Psi - \Gamma_{\Lambda i}{}^j H_j{}^\Psi + \Gamma_{\Lambda\Phi}{}^\Psi H_i{}^\Phi = 0. \tag{11.37}$$

Соответственно в спинорное расслоение вещественная связность продолжается условием

$$\sum_{I=1}^{2N} (\eta_K)^i{}_{AB} \nabla_k (\eta_K)_j{}^{AB} = 0. \tag{11.38}$$

Это означает, что на вещественном пространстве связность, согласованная с метрикой будет иметь вид

$$\nabla_i g_{ik} = 0, \tag{11.39}$$

где

$$\nabla_i := H_i{}^{\Lambda}\nabla_{\Lambda}. \tag{11.40}$$

Example 11.1. *n=4. Индекс метрики равен 1.*

Пусть

$$g^i{}_{AB'} := (\eta_1)^i{}_{AB}\bar{S}_{B'}{}^{B}, \quad g_i{}^{AB'} := (\eta_1)_i{}^{AB}S_B{}^{B'}, \quad \overline{g_j{}^{A'B}} = \bar{g}_j{}^{B'A} = g_j{}^{AB'}. \tag{11.41}$$

Тогда частный случай условий согласования связностей может иметь вид

$$g^i{}_{AB'}\nabla_k g_j{}^{AB'} + \bar{g}^i{}_{B'A}\nabla_k \bar{g}_j{}^{B'A} = 0. \tag{11.42}$$

Это условие расписывается как

$$\begin{gathered} g^i{}_{AB'}(\partial_k g_j{}^{AB'} - \Gamma_{kj}{}^{l}g_l{}^{AB'} + \tilde{\tilde{\Gamma}}_{kC}{}^{A}g_j{}^{CB'} + \tilde{\Gamma}_{kC'}{}^{B'}g_j{}^{AC'} + \partial_k g_j{}^{AB'} - \Gamma_{kj}{}^{l}g_l{}^{AB'} + \\ + \bar{\tilde{\tilde{\Gamma}}}_{kC'}{}^{B'}g_j{}^{AC'} + \bar{\tilde{\Gamma}}_{kC}{}^{A}g_j{}^{CB'}) = 0. \end{gathered} \tag{11.43}$$

Поэтому это соотношение ввиду замечания 11.1 перепишется так

$$\partial_k g_j{}^{AB'} - \Gamma_{kj}{}^{l}g_l{}^{AB'} + \frac{1}{2}(\tilde{\tilde{\Gamma}}_{kC}{}^{A} + \bar{\tilde{\Gamma}}_{kC}{}^{A})g_j{}^{CB'} + \frac{1}{2}\overline{(\tilde{\tilde{\Gamma}}_{kC}{}^{B} + \bar{\tilde{\Gamma}}_{kC}{}^{B})}g_j{}^{AC'} = 0. \tag{11.44}$$

Определим $\Gamma_{kC}{}^{A} := \frac{1}{2}(\tilde{\tilde{\Gamma}}_{kC}{}^{A} + \bar{\tilde{\Gamma}}_{kC}{}^{A})$, что приведет к связности [22, т. 1, с. 284]

$$\nabla_k g_j{}^{AB'} = 0. \tag{11.45}$$

Этим доказана теорема.

Теорема 11.1. *Рассмотрим псевдориманово многообразие V_{2n} ($n \geq 4$). Пусть в касательном расслоении со слоями, изоморфными $R^{2n}_{(n,n)}$, введена некоторая связность без кручения. Тогда условие согласования связностей в спинорном и касательном расслоении может иметь вид*

$$\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{B}}}\nabla_{\tilde{\Lambda}}\eta_{\tilde{\Omega}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}} = 0. \tag{11.46}$$

Чтобы продолжение было однозначным, потребуем

$$\begin{gathered} \tilde{\tilde{\Gamma}}_{\tilde{\Lambda}\tilde{\tilde{L}}}{}^{\tilde{\tilde{M}}} := \tfrac{1}{N^2}\tilde{\tilde{\Gamma}}_{\tilde{\Lambda}\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}(\eta_{\tilde{\Psi}\tilde{\tilde{L}}\tilde{\tilde{B}}}\eta_{\tilde{\Omega}}{}^{\tilde{\tilde{M}}\tilde{\tilde{B}}}\eta^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}\tilde{\tilde{C}}}\eta^{\tilde{\Omega}\tilde{\tilde{K}}\tilde{\tilde{C}}} - \tfrac{(2n-2)}{4}\delta_{\tilde{\tilde{L}}}{}^{\tilde{\tilde{M}}}\delta_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}), \\ \tilde{\Gamma}_{\tilde{\Lambda}\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}} := -\tfrac{2}{2n-4}\tilde{\tilde{\Gamma}}_{\tilde{\Lambda}\tilde{\tilde{C}}}{}^{\tilde{\tilde{D}}}(\varepsilon_{\tilde{\tilde{D}}\tilde{\tilde{K}}}{}^{\tilde{\tilde{C}}\tilde{\tilde{A}}} - \tfrac{2n-2}{2N^2}\delta_{\tilde{\tilde{K}}}{}^{\tilde{\tilde{A}}}\delta_{\tilde{\tilde{D}}}{}^{\tilde{\tilde{C}}}). \end{gathered} \tag{11.47}$$

Для продолжения связности на комплексную реализацию $\mathbb{C}V_n$ *со слоями, изоморфными* $\mathbb{C}^n$, *необходимо, чтобы*

$$\nabla_{\tilde{\Lambda}} m_{\tilde{\Psi}}{}^{\Psi} = 0, \quad \nabla_{\tilde{\Lambda}} \bar{m}_{\tilde{\Psi}}{}^{\Psi'} = 0, \quad \nabla_{\Lambda} := m^{\tilde{\Lambda}}{}_{\Lambda} \nabla_{\tilde{\Lambda}}, \quad \bar{\nabla}_{\Lambda'} := \bar{m}^{\tilde{\Lambda}}{}_{\Lambda'} \nabla_{\tilde{\Lambda}}. \tag{11.48}$$

Тогда условие согласования связностей примет вид

$$\begin{gathered} \nabla_{\Lambda} (\tilde{M}_K)_A{}^{\tilde{\tilde{B}}} = 0, \quad \nabla_{\Lambda} (\tilde{\tilde{M}}_K)_A{}^{\tilde{\tilde{B}}} = 0, \quad \bar{\nabla}_{\Lambda'} (\tilde{M}_K)_A{}^{\tilde{\tilde{B}}} = 0, \quad \bar{\nabla}_{\Lambda'} (\tilde{\tilde{M}}_K)_A{}^{\tilde{\tilde{B}}} = 0, \\ \sum_{K=1}^{2N} (\eta_K)^{\Psi}{}_{AB} \bar{\nabla}_{\Lambda'} (\eta_K)_{\Omega}{}^{AB} = 0, \quad \sum_{K=1}^{2N} (\eta_K)^{\Psi}{}_{AB} \nabla_{\Lambda} (\eta_K)_{\Omega}{}^{AB} = 0. \end{gathered} \tag{11.49}$$

Чтобы продолжение было однозначным, потребуем

$$\begin{gathered} (\tilde{\tilde{\Gamma}}_K)_{\Lambda L}{}^{M} := \tfrac{2}{N} (\tilde{\tilde{\Gamma}}_K)_{\Lambda D}{}^{A} ((\eta_K)_{\Psi LB} (\eta_K)_{\Omega}{}^{MB} (\eta_K)^{\Psi}{}_{AC} (\eta_K)^{\Omega DC} - \tfrac{(n-2)}{4} \delta_L{}^{M} \delta_D{}^{A}), \\ (\tilde{\Gamma}_K)_{\Lambda A}{}^{C} := -\tfrac{2}{n-4} (\tilde{\tilde{\Gamma}}_K)_{\Lambda K}{}^{L} ((\varepsilon_K)_{LA}{}^{KC} - \tfrac{n-2}{N} \delta_L{}^{K} \delta_A{}^{C}). \end{gathered} \tag{11.50}$$

В случае вложения действительного пространства в комплексное необходимо выполнение соотношений

$$\nabla_{\Lambda} H_i{}^{\Psi} = 0, \quad \bar{\nabla}_{\Lambda'} \bar{H}_i{}^{\Psi'} = 0, \quad \nabla_i := H_i{}^{\Lambda} \nabla_{\Lambda}. \tag{11.51}$$

Тогда условие согласования связностей примет вид

$$\sum_{I=1}^{2N} (\eta_K)^{i}{}_{AB} \nabla_k (\eta_K)_j{}^{AB} = 0. \tag{11.52}$$

Следствие 11.1. *В случае, если в касательном расслоении многообразия* V_{2n} *задана связность без кручения, согласованная с метрикой*

$$\nabla_{\tilde{\Lambda}} G_{\tilde{\Psi}\tilde{\Omega}} = 0, \tag{11.53}$$

то в касательном расслоении комплексной реализации $\mathbb{C}V_n$ *индуцируется связность*

$$\nabla_{\Lambda} g_{\Omega\Psi} = 0, \quad \bar{\nabla}_{\Lambda'} \bar{g}_{\Omega'\Psi'} = 0. \tag{11.54}$$

Для вложения $V_n \subset \mathbb{C}V_n$ *индуцируется связность*

$$\nabla_i g_{ik} = 0. \tag{11.55}$$

Построим операторы $P^{\tilde{\Psi}}{}_{\tilde{\tilde{A}}} := \eta^{\tilde{\Psi}}{}_{\tilde{\tilde{B}}\tilde{\tilde{A}}} X^{\tilde{\tilde{B}}}$ и $P_{\tilde{\Psi}}{}^{\tilde{\tilde{A}}} := \eta_{\tilde{\Psi}}{}^{\tilde{\tilde{B}}\tilde{\tilde{A}}} Y_{\tilde{\tilde{B}}}$ такие, что $X^{\tilde{\tilde{A}}} Y_{\tilde{\tilde{A}}} = 2$ и $P_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}} P_{\tilde{\Psi}\tilde{\tilde{A}}} = G_{\tilde{\Lambda}\tilde{\Psi}}$. Это всегда можно сделать при $n \geq 8$. Тогда условие согласования связностей для вещественной реализации может иметь и такой вид

$$\nabla_{\tilde{\Lambda}} P_{\tilde{\Psi}}{}^{\tilde{\tilde{A}}} = 0, \tag{11.56}$$

которое распишется как

$$\partial_{\tilde{\Lambda}} P_{\tilde{\Psi}}{}^{\tilde{\tilde{A}}} - \Gamma_{\tilde{\Lambda}\tilde{\Psi}}{}^{\tilde{\Phi}} P_{\tilde{\Phi}}{}^{\tilde{\tilde{A}}} + \Gamma_{\tilde{\Lambda}\tilde{\tilde{C}}}{}^{\tilde{\tilde{A}}} P_{\tilde{\Psi}}{}^{\tilde{\tilde{C}}} = 0. \tag{11.57}$$

Поэтому можно однозначно определить связность в спинорном расслоении как

$$\Gamma_{\tilde{\varLambda}\tilde{\tilde{C}}}{}^{\tilde{\tilde{A}}} := P^{\tilde{\varPsi}}{}_{\tilde{\tilde{C}}}\partial_{\tilde{\varLambda}}P_{\tilde{\varPsi}}{}^{\tilde{\tilde{A}}} + \Gamma_{\tilde{\varLambda}\tilde{\varPsi}}{}^{\tilde{\varPhi}}P^{\tilde{\varPsi}}{}_{\tilde{\tilde{C}}}P_{\tilde{\varPhi}}{}^{\tilde{\tilde{A}}}. \tag{11.58}$$

Для комплексной реализации определены операторы

$$\begin{gathered}(P_K)^{\Psi}{}_A := m_{\tilde{\varPsi}}{}^{\Psi}P^{\tilde{\varPsi}}{}_{\tilde{\tilde{A}}}(\tilde{M}_K)_A{}^{\tilde{\tilde{A}}}, \quad (P_K)^{\Psi'}{}_A := \bar{m}_{\tilde{\varPsi}}{}^{\Psi'}P^{\tilde{\varPsi}}{}_{\tilde{\tilde{A}}}(\tilde{M}_K)_A{}^{\tilde{\tilde{A}}},\\ (P^*_K)_{\Psi}{}^A := m^{\tilde{\varPsi}}{}_{\Psi}P_{\tilde{\varPsi}}{}^{\tilde{\tilde{A}}}(\tilde{M}^*_K)^A{}_{\tilde{\tilde{A}}}, \quad (P^*_K)_{\Psi'}{}^A := \bar{m}^{\tilde{\varPsi}}{}_{\Psi'}P_{\tilde{\varPsi}}{}^{\tilde{\tilde{A}}}(\tilde{M}^*_K)^A{}_{\tilde{\tilde{A}}}.\end{gathered} \tag{11.59}$$

Теорема 11.2. *Рассмотрим псевдориманово многообразие V_{2n} $(n \geq 4)$. Пусть в касательном расслоении со слоями, изоморфными $R^{2n}_{(n,n)}$, введена некоторая связность без кручения. Тогда условие согласования связностей в спинорном и касательном расслоении может иметь вид*

$$\nabla_{\tilde{\varLambda}}P_{\tilde{\varPsi}}{}^{\tilde{\tilde{A}}} = 0. \tag{11.60}$$

Чтобы продолжение было однозначным, потребуем

$$\Gamma_{\tilde{\varLambda}\tilde{\tilde{C}}}{}^{\tilde{\tilde{A}}} := -P^{\tilde{\varPsi}}{}_{\tilde{\tilde{C}}}\partial_{\tilde{\varLambda}}P_{\tilde{\varPsi}}{}^{\tilde{\tilde{A}}} + \Gamma_{\tilde{\varLambda}\tilde{\varPsi}}{}^{\tilde{\varPhi}}P^{\tilde{\varPsi}}{}_{\tilde{\tilde{C}}}P_{\tilde{\varPhi}}{}^{\tilde{\tilde{A}}}. \tag{11.61}$$

Для продолжения связности на комплексную реализацию $\mathbb{C}V_n$ со слоями, изоморфными $\mathbb{C}^n$, необходимо, чтобы

$$\nabla_{\tilde{\varLambda}}m_{\tilde{\varPsi}}{}^{\Psi} = 0, \quad \nabla_{\tilde{\varLambda}}\bar{m}_{\tilde{\varPsi}}{}^{\Psi'} = 0, \quad \nabla_{\Lambda} := m^{\tilde{\varLambda}}{}_{\Lambda}\nabla_{\tilde{\varLambda}}, \quad \bar{\nabla}_{\Lambda'} := \bar{m}^{\tilde{\varLambda}}{}_{\Lambda'}\nabla_{\tilde{\varLambda}}. \tag{11.62}$$

Тогда условие согласования связностей примет вид

$$\begin{gathered}\nabla_{\Lambda}(\tilde{M}_K)_A{}^{\tilde{\tilde{B}}} = 0, \quad \bar{\nabla}_{\Lambda'}(\tilde{M}_K)_A{}^{\tilde{\tilde{B}}} = 0,\\ \nabla_{\Lambda}(P_K)^{\Psi}{}_A = 0, \quad \bar{\nabla}_{\Lambda'}(P_K)^{\Psi}{}_A = 0, \quad \nabla_{\Lambda}(\bar{P}_K)^{\Psi'}{}_A = 0, \quad \bar{\nabla}_{\Lambda'}(\bar{P}_K)^{\Psi'}{}_A = 0.\end{gathered} \tag{11.63}$$

В случае вложения действительного пространства в комплексное необходимо выполнение соотношений

$$\nabla_{\Lambda}H_i{}^{\Psi} = 0, \quad \bar{\nabla}_{\Lambda'}\bar{H}_i{}^{\Psi'} = 0, \quad \nabla_i := H_i{}^{\Lambda}\nabla_{\Lambda}. \tag{11.64}$$

Тогда условие согласования связностей примет вид

$$\nabla_i(P_K)^j{}_A = 0, \quad \nabla_i(P_K)_j{}^A = 0. \tag{11.65}$$

Теперь появилась возможность построить операторы

$$\begin{gathered}(Y_K)^A := y^{\Psi}(P^*_K)_{\Psi}{}^A, \quad y^{\Psi} = \sum_{J=1}^{2N}(Y_J)^B(P_J)^{\Psi}{}_B,\\ (\bar{Y}_K)^A := \bar{y}^{\Psi'}(\bar{P}^*_K)_{\Psi'}{}^A, \quad \bar{y}^{\Psi'} = \sum_{J=1}^{2N}(\bar{Y}_J)^B(\bar{P}_J)^{\Psi'}{}_B,\\ L_x(Y_K)^A := x^{\Omega}\nabla_{\Omega}(Y_K)^A - \sum_{J=1}^{2N}(Y_J)^B(P_J)^{\Psi}{}_B(P^*_K)_{\Omega}{}^A\nabla_{\Psi}x^{\Omega},\\ \bar{L}_{\bar{x}}(\bar{Y}_K)^A := \bar{x}^{\Omega'}\bar{\nabla}_{\Omega'}(\bar{Y}_K)^A - \sum_{J=1}^{2N}(\bar{Y}_J)^B(\bar{P}_J)^{\Psi'}{}_B(\bar{P}^*_K)_{\Omega'}{}^A\bar{\nabla}_{\Psi'}\bar{x}^{\Omega'}.\end{gathered} \tag{11.66}$$

Эти операторы будут инвариантами связности только в том случае, когда

$$\begin{gathered}\partial_{\Lambda}(P_K)^{\Psi}{}_{B}=0,\quad \bar\partial_{\Lambda'}(P_K)^{\Psi}{}_{B}=0,\quad \partial_{\Lambda}(P^*_K)_{\Psi}{}^{B}=0,\quad \bar\partial_{\Lambda'}(P^*_K)_{\Psi}{}^{B}=0,\\ \partial_{\Lambda}(\bar P_K)^{\Psi'}{}_{B}=0,\quad \bar\partial_{\Lambda'}(\bar P_K)^{\Psi'}{}_{B}=0,\quad \partial_{\Lambda}(\bar P^*_K)_{\Psi'}{}^{B}=0,\quad \bar\partial_{\Lambda'}(\bar P^*_K)_{\Psi'}{}^{B}=0,\end{gathered}\tag{11.67}$$

$$\partial_{\tilde\Lambda}P^{\tilde\Psi}{}_{\tilde{\tilde B}}=0,\quad \partial_{\tilde\Lambda}P_{\tilde\Psi}{}^{\tilde{\tilde B}}=0,\quad \partial_{\tilde\Lambda}m^{\tilde\Psi}{}_{\Psi}=0,\quad \partial_{\tilde\Lambda}m_{\tilde\Psi}{}^{\Psi}=0,\quad \partial_{\tilde\Lambda}\bar m^{\tilde\Psi}{}_{\Psi'}=0,\quad \partial_{\tilde\Lambda}\bar m_{\tilde\Psi}{}^{\Psi'}=0.\tag{11.68}$$

Тогда из (11.58) будет следовать

$$(\Gamma_K)_{\Lambda C}{}^{A}=\Gamma_{\Lambda\Psi}{}^{\Phi}\underbrace{(P_K)^{\Psi}{}_{C}(P^*_K)_{\Phi}{}^{A}}_{:=(P_K)_{\Phi}{}^{\Psi}{}_{C}{}^{A}},\quad (\bar\Gamma_K)_{\Lambda' C}{}^{A}=\Gamma_{\Lambda'\Psi}{}^{\Phi}\underbrace{(P_K)^{\Psi}{}_{C}(P^*_K)_{\Phi}{}^{A}}_{:=(P_K)_{\Phi}{}^{\Psi}{}_{C}{}^{A}}=0,\tag{11.69}$$

что обеспечит инвариантность операторов. Поэтому будут выполнены тождества

$$\begin{aligned}L_{[x,y]}(Y_K)^A&=L_xL_y(Y_K)^A-L_yL_x(Y_K)^A,\\ \bar L_{[\bar x,\bar y]}(\bar Y_K)^A&=\bar L_{\bar x}\bar L_{\bar y}(\bar Y_K)^A-\bar L_{\bar y}\bar L_{\bar x}(\bar Y_K)^A.\end{aligned}\tag{11.70}$$

При n=4 подобные связности рассматривались в [9]. Однако, там в качестве $(P_K)_{\Phi}{}^{\Psi}{}_{C}{}^{A}$ рассматривалось выражение вида $(\eta_K)^{\Psi}{}_{DC}(\eta_K)_{\Phi}{}^{DA}$, что вызывало кручение таких операторов.

Теорема 11.3. *Операторы*

$$\begin{aligned}L_x(Y_K)^A&:=x^{\Omega}\partial_{\Omega}(Y_K)^A-\sum_{J=1}^{2N}(Y_J)^B(P_J)^{\Psi}{}_{B}(P^*_K)_{\Omega}{}^{A}\partial_{\Psi}x^{\Omega},\\ \bar L_{\bar x}(\bar Y_K)^A&:=\bar x^{\Omega'}\bar\partial_{\Omega'}(\bar Y_K)^A-\sum_{J=1}^{2N}(\bar Y_J)^B(\bar P_J)^{\Psi'}{}_{B}(\bar P^*_K)_{\Omega'}{}^{A}\bar\partial_{\Psi'}\bar x^{\Omega'}\end{aligned}\tag{11.71}$$

будут являться инвариантами комплексной связности из теоремы 11.2 при условии постоянства операторов

$$\begin{gathered}\partial_{\Lambda}(P_K)^{\Psi}{}_{B}=0,\quad \bar\partial_{\Lambda'}(P_K)^{\Psi}{}_{B}=0,\quad \partial_{\Lambda}(P^*_K)_{\Psi}{}^{B}=0,\quad \bar\partial_{\Lambda'}(P^*_K)_{\Psi}{}^{B}=0,\\ \partial_{\Lambda}(\bar P_K)^{\Psi'}{}_{B}=0,\quad \bar\partial_{\Lambda'}(\bar P_K)^{\Psi'}{}_{B}=0,\quad \partial_{\Lambda}(\bar P^*_K)_{\Psi'}{}^{B}=0,\quad \bar\partial_{\Lambda'}(\bar P^*_K)_{\Psi'}{}^{B}=0,\\ \partial_{\tilde\Lambda}P^{\tilde\Psi}{}_{\tilde{\tilde B}}=0,\quad \partial_{\tilde\Lambda}P_{\tilde\Psi}{}^{\tilde{\tilde B}}=0,\\ \partial_{\tilde\Lambda}m^{\tilde\Psi}{}_{\Psi}=0,\quad \partial_{\tilde\Lambda}m_{\tilde\Psi}{}^{\Psi}=0,\quad \partial_{\tilde\Lambda}\bar m^{\tilde\Psi}{}_{\Psi'}=0,\quad \partial_{\tilde\Lambda}\bar m_{\tilde\Psi}{}^{\Psi'}=0,\\ \sum_{K=1}^{2N}(P_K)^{\Psi}{}_{B}(P^*_K)_{\Phi}{}^{B}=\delta_{\Psi}{}^{\Phi},\quad \sum_{K=1}^{2N}(\bar P_K)^{\Psi'}{}_{B}(\bar P^*_K)_{\Phi'}{}^{B}=\delta_{\Psi'}{}^{\Phi'}.\end{gathered}\tag{11.72}$$

Такие операторы будут удовлетворять тождествам

$$\begin{aligned}L_{[x,y]}(Y_K)^A&=L_xL_y(Y_K)^A-L_yL_x(Y_K)^A,\\ \bar L_{[\bar x,\bar y]}(\bar Y_K)^A&=\bar L_{\bar x}L_{\bar y}(\bar Y_K)^A-\bar L_{\bar y}L_{\bar x}(\bar Y_K)^A\end{aligned}\tag{11.73}$$

и являться естественным продолжением операторов $L_x(y)^{\Psi}=x^{\Omega}\partial_{\Omega}y^{\Psi}-y^{\Omega}\partial_{\Omega}x^{\Psi}$, $\bar L_{\bar x}(\bar y)^{\bar\Psi}=\bar x^{\Omega'}\bar\partial_{\Omega'}\bar y^{\Psi'}-\bar y^{\Omega'}\bar\partial_{\Omega'}\bar x^{\Psi'}$ *в спинорное расслоение*

$$\begin{gathered}L_x(y)^{\Psi}=\sum_{K=1}^{2N}L_x(Y_K)^A(P_K)^{\Psi}{}_{A},\quad \bar L_{\bar x}(\bar y)^{\Psi'}=\sum_{K=1}^{2N}\bar L_{\bar x}(\bar Y_K)^A(\bar P_K)^{\Psi'}{}_{A},\\ L_x(Y_K)^A=L_x(y)^{\Psi}(P^*_K)_{\Psi}{}^{A},\quad \bar L_{\bar x}(\bar Y_K)^A=\bar L_{\bar x}(\bar y)^{\Psi'}(\bar P^*_K)_{\Psi'}{}^{A}.\end{gathered}\tag{11.74}$$

12 О классификации тензоров с симметриями тензора кривизны. Спиноры кривизны

В этом параграфе рассказывается о том, как сопоставить тензорам, обладающим симметриями тензора кривизны, их спинорные аналоги. Это упрощает проблему классификации таких тензоров для малых размерностей. Вывод всех результатов этого параграфа сделан на основе [22, т. 1, с. 285-303].

Ограничим себя случаем, когда $(\Gamma_K)_{\Lambda A}{}^B = (\tilde{\Gamma}_K)_{\Lambda A}{}^B = (\tilde{\tilde{\Gamma}}_K)_{\Lambda A}{}^B$. Тогда тензоры кривизны можно вычислить по формулам

$$\nabla_{[\Lambda}\nabla_{\Psi]}r^\Omega = R_{\Lambda\Psi\Phi}{}^\Omega r^\Phi, \quad \nabla_{[\Lambda}\nabla_{\Psi]}X^A = \sum_{K=1}^{2N}(\mathcal{R}_K)_{\Lambda\Psi C}{}^A X^C. \tag{12.1}$$

Для этого рассмотрим согласование комплексных связностей, индуцируемое формулами (11.30), (11.33) при ковариантном постоянстве метрического тензора. Потребуем, чтобы

$$\nabla_\Lambda(\varepsilon_K)_{AB}{}^{CD} = \partial_\Lambda(\varepsilon_K)_{AB}{}^{CD} = 0. \tag{12.2}$$

Следовательно,

$$\sum_{K=1}^{2N}(\nabla_\Theta(\eta_K)^\Psi{}_{AB})(\nabla_\Lambda(\eta_K)_\Omega{}^{AB}) = 0. \tag{12.3}$$

Поэтому условия интегрируемости (11.30) примут вид

$$\begin{gathered}\sum_{K=1}^{2N}(\eta_K)^\Theta{}_{AB}\nabla_{[\Lambda}\nabla_{\Psi]}(\eta_K)_\Omega{}^{AB} = 0,\\ N^2 R_{\Lambda\Psi\Omega}{}^\Theta = \sum_{K=1}^{2N}((\mathcal{R}_K)_{\Lambda\Psi C}{}^A(\eta_K)^\Theta{}_{AB}(\eta_K)_\Omega{}^{CB} + (\mathcal{R}_K)_{\Lambda\Psi C}{}^B(\eta_K)^\Theta{}_{AB}(\eta_K)_\Omega{}^{AC}).\end{gathered} \tag{12.4}$$

Определим

$$(A_K)_{\Theta\Phi L}{}^X := \frac{1}{2}(\eta_K)_{[\Theta}{}^{MX}(\eta_K)_{\Phi]ML}. \tag{12.5}$$

Тогда из (8.27) будет следовать

$$(A_K)_{\Phi\Theta L}{}^X(A_K)_{\Omega\Gamma X}{}^L = \frac{N}{8}g_{\Phi[\Gamma}g_{\Omega]\Theta}. \tag{12.6}$$

Поэтому можно положить

$$\begin{gathered}(R_K)_{\Lambda\Psi C}{}^N := -R_{\Lambda\Psi\Theta\Phi}(A_K)^{\Theta\Phi}{}_C{}^N, \quad R_{\Lambda\Psi\Theta\Phi} = \tfrac{8}{N}(A_K)_{\Theta\Phi C}{}^N(R_K)_{\Lambda\Psi N}{}^C,\\ (R_K)_{\Lambda\Psi C}{}^M = -\tfrac{2}{n-4}(R_K)_{\Lambda\Psi L}{}^D\varepsilon_{CD}{}^{ML}.\end{gathered} \tag{12.7}$$

Очевидно, что $(R_K)_{\Lambda\Psi C}{}^N = (\mathcal{R}_K)_{\Lambda\Psi C}{}^N$, например, если условие согласования связностей имеет вид

$$\nabla_\Lambda(\eta_K)_\Psi{}^{AB} = 0. \tag{12.8}$$

Для дальнейших выкладок следует воспользоваться тождествами (номер K будем опускать, поскольку все остальные выкладки не зависят от этого номера)

$$\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_2} = \frac{N}{2}g_{\Lambda_1[\Lambda_4}g_{\Lambda_3]\Lambda_2}. \tag{12.9}$$

$$\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} = Ng_{[\Lambda_3|[\Lambda_2}g_{\Lambda_1]|[\Lambda_6}g_{\Lambda_5]|\Lambda_4]}. \tag{12.10}$$

$$\begin{gathered}\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2} = \\ = \tfrac{N}{4}\big(g_{\Lambda_1[\Lambda_8}g_{\Lambda_7]\Lambda_2}g_{\Lambda_3[\Lambda_6}g_{\Lambda_5]\Lambda_4} - g_{\Lambda_1[\Lambda_6}g_{\Lambda_5]\Lambda_2}g_{\Lambda_4[\Lambda_8}g_{\Lambda_7]\Lambda_3} + \\ + g_{\Lambda_1[\Lambda_4}g_{\Lambda_3]\Lambda_2}g_{\Lambda_5[\Lambda_8}g_{\Lambda_7]\Lambda_6}\big) + N\big(g_{[\Lambda_3|[\Lambda_2}g_{\Lambda_1][\Lambda_8}g_{\Lambda_7][\Lambda_6}g_{\Lambda_5]|\Lambda_4]} + \\ + g_{[\Lambda_5|[\Lambda_4}g_{\Lambda_3][\Lambda_7}g_{\Lambda_8][\Lambda_2}g_{\Lambda_1]|\Lambda_6]} + g_{[\Lambda_3|[\Lambda_7}g_{\Lambda_8][\Lambda_5}g_{\Lambda_6][\Lambda_1}g_{\Lambda_2]|\Lambda_4]}\big).\end{gathered} \tag{12.11}$$

Доказательство в приложении.

Теорема 12.1. *Классификацию битензора, обладающего свойствами ($\Lambda, \Psi, ... = \overline{1, n}$, $A, B, ... = \overline{1, N}$)*

$$R_{\Lambda\Psi\Phi\Theta} = R_{[\Lambda\Psi][\Phi\Theta]}, \quad R_{\Lambda\Psi\Phi\Theta} = R_{\Phi\Theta\Lambda\Psi} \tag{12.12}$$

и принадлежащего касательному расслоению $\tau(\mathbb{C}V_n)$ над аналитическим псевдоримановым пространством $\mathbb{C}V_n$, можно свести к классификации тензора $R_A{}^B{}_C{}^D$ N-мерного комплексного спинорного пространства такого, что

$$R_C{}^D{}_S{}^R := R_{\Lambda\Psi\Phi\Theta}A^{\Lambda\Psi}{}_C{}^DA^{\Phi\Theta}{}_S{}^R, \quad A_{\Lambda\Psi C}{}^D := \frac{1}{2}\eta_{[\Lambda}{}^{AD}\eta_{\Psi]AC}. \tag{12.13}$$

Кроме того, выполнены соотношения

$$R_K{}^K{}_S{}^R = R_S{}^R{}_K{}^K = 0, \quad R_C{}^D{}_S{}^R = R_S{}^R{}_C{}^D. \tag{12.14}$$

Разложение

$$\begin{gathered}R_C{}^K{}_M{}^A = C_C{}^K{}_M{}^A - \tfrac{4}{N(n-2)}R_G{}^D{}_N{}^P\varepsilon^{ABKL}\varepsilon^{GN}{}_{ML}\varepsilon_{DPCB} - \\ -\big(\tfrac{1}{2(n-1)(n-2)} - \tfrac{1}{4(n-2)}\big)R\varepsilon^{ABKL}\varepsilon_{CBML} - \big(\tfrac{1}{4(n-2)} - \tfrac{n}{8(n-1)(n-2)}\big)R\delta_C{}^A\delta_M{}^K\end{gathered} \tag{12.15}$$

соответствует разложению тензора $R_{\Lambda\Psi}{}^{\Phi\Theta}$

$$R_{\Lambda\Psi}{}^{\Phi\Theta} = C_{\Lambda\Psi}{}^{\Phi\Theta} + \frac{4}{n-2}R_{[\Lambda}{}^{[\Phi}g_{\Psi]}{}^{\Theta]} - \frac{2}{(n-1)(n-2)}Rg_{[\Lambda}{}^{[\Phi}g_{\Psi]}{}^{\Theta]} \tag{12.16}$$

на неприводимые ортогональными преобразованиями компоненты. При этом тождество Бианки

$$R_{\Lambda\Psi\Phi\Theta} + R_{\Lambda\Theta\Psi\Phi} + R_{\Lambda\Phi\Theta\Psi} = 0 \tag{12.17}$$

будет выглядеть как

$$R_L{}^D{}_S{}^L = -\frac{1}{8}\cdot R\delta_S{}^D. \tag{12.18}$$

Доказательство.

1. Первый шаг. Тождество Бианки $R_{[\Lambda\Psi\Phi]\Theta}=0$.

$$\begin{gathered}
R_{[\Lambda\Psi\Phi]\Theta}\eta^{\Lambda}{}_{AB}\eta^{\Psi AC}\eta^{\Phi}{}_{DC}\eta^{\Theta DL}= \\
=R_{\Lambda\Psi\Theta\Phi}\tfrac{1}{3}(\eta^{\Lambda}{}_{AB}\eta^{\Psi AC}\eta^{\Phi}{}_{DC}+\eta^{\Phi}{}_{AB}\eta^{\Lambda AC}\eta^{\Psi}{}_{DC}+\eta^{\Psi}{}_{AB}\eta^{\Phi AC}\eta^{\Lambda}{}_{DC})\eta^{\Theta DL}= \\
=R_{\Lambda\Psi\Phi\Theta}\tfrac{1}{3}(\eta^{\Lambda}{}_{AB}\eta^{\Psi AC}\eta^{\Phi}{}_{DC}+g^{\Phi\Lambda}\eta^{\Psi}{}_{DB}- \\
-\eta^{\Lambda}{}_{AB}\eta^{\Phi AC}\eta^{\Psi}{}_{DC}+\eta^{\Lambda}{}_{AB}\eta^{\Psi AC}\eta^{\Phi}{}_{DC}-g^{\Phi\Psi}\eta^{\Lambda}{}_{DB})\eta^{\Theta DL}= \\
=R_{\Lambda\Psi\Phi\Theta}(\eta^{\Lambda}{}_{AB}\eta^{\Psi AC}\eta^{\Phi}{}_{DC}+g^{\Phi\Lambda}\eta^{\Psi}{}_{DB})\eta^{\Theta DL}=4R_{B}{}^{C}{}_{C}{}^{L}+\tfrac{1}{2}R\delta_{B}{}^{L}=0.
\end{gathered}\tag{12.19}$$

Обратно,

$$\begin{gathered}
(4R_{A_1}{}^{C}{}_{C}{}^{A_5}+\tfrac{1}{2}R\delta_{A_1}{}^{A_5})\eta_{[\Lambda_5|A_5A_6|}\eta_{\Lambda_6]}{}^{A_7A_6}\eta_{[\Lambda_7|A_7A_8|}\eta_{\Lambda_8]}{}^{A_1A_8}= \\
=R_{\Lambda_1\Lambda_2\Lambda_3\Lambda_4}\eta^{\Lambda_1}{}_{A_1A_2}\eta^{\Lambda_2A_3A_2}\eta^{\Lambda_3}{}_{A_3A_4}\eta^{\Lambda_4A_5A_4}\eta_{[\Lambda_5|A_5A_6|}\eta_{\Lambda_6]}{}^{A_7A_6}\eta_{[\Lambda_7|A_7A_8|}\eta_{\Lambda_8]}{}^{A_1A_8}+ \\
+\tfrac{1}{2}R\eta_{[\Lambda_5|A_1A_6|}\eta_{\Lambda_6]}{}^{A_7A_6}\eta_{[\Lambda_7|A_7A_8|}\eta_{\Lambda_8]}{}^{A_1A_8}=\tfrac{N}{4}(R_{\Lambda_8\Lambda_7\Lambda_6\Lambda_5}-R_{\Lambda_6\Lambda_5\Lambda_7\Lambda_8}- \\
-Rg_{\Lambda_5[\Lambda_8}g_{\Lambda_7]\Lambda_6})+N(-R_{[\Lambda_8|[\Lambda_5}g_{\Lambda_6]\Lambda_7]}+R_{[\Lambda_6|[\Lambda_8\Lambda_7]|\Lambda_5]}+ \\
+R_{[\Lambda_6|[\Lambda_7}g_{\Lambda_8]|\Lambda_5]})+\tfrac{1}{2}\cdot\tfrac{N}{2}Rg_{\Lambda_5[\Lambda_8}g_{\Lambda_7]\Lambda_6}=\tfrac{N}{2}(R_{\Lambda_8\Lambda_7\Lambda_6\Lambda_5}+2R_{\Lambda_6[\Lambda_8\Lambda_7]\Lambda_5})=0.
\end{gathered}\tag{12.20}$$

2. Второй шаг. Тензор Риччи $R_{\Lambda\Psi}$.

Воспользуемся двумя вспомогательными тождествами.

$$\begin{gathered}
R_{\Lambda\Psi C}{}^{M}=-\tfrac{2}{n-4}R_{\Lambda\Psi L}{}^{D}\varepsilon_{CD}{}^{ML}, \\
\eta_{[\Gamma}{}^{CL}\eta_{\Omega]CD}=-\tfrac{2}{n-4}\eta_{[\Gamma}{}^{CN}\eta_{\Omega]MN}\varepsilon_{CD}{}^{ML}.
\end{gathered}\tag{12.21}$$

Это определит тождество

$$\eta_{[\Gamma|CD|}\eta_{\Omega]}{}^{CL}R_{\Lambda\Psi L}{}^{D}=-\eta_{[\Gamma}{}^{CN}\eta_{\Omega]MN}R_{\Lambda\Psi C}{}^{M}.\tag{12.22}$$

Поэтому

$$\begin{gathered}
\tfrac{4}{N}\eta_{\Lambda}{}^{AB}\eta_{\Psi CD}R_{A}{}^{C}{}_{B}{}^{D}-\tfrac{1}{4}Rg_{\Lambda\Psi}= \\
=\tfrac{1}{N}\eta_{\Lambda}{}^{AB}\eta_{\Psi CD}\eta_{\Gamma AL}\eta_{\Phi}{}^{CL}\eta_{\Omega NB}\eta_{\Theta}{}^{ND}R^{\Gamma\Phi\Omega\Theta}-\tfrac{1}{4}Rg_{\Lambda\Psi}= \\
=\tfrac{1}{N}(g_{\Phi\Psi}\eta_{\Lambda}{}^{AB}\eta_{\Gamma AL}\eta_{[\Theta}{}^{NL}\eta_{\Omega]NB}-\eta_{\Lambda}{}^{AB}\eta_{\Gamma AL}\eta_{\Psi}{}^{CL}\eta_{\Phi CD}\eta_{\Theta}{}^{ND}\eta_{\Omega NB})R^{\Gamma\Phi\Omega\Theta}- \\
-\tfrac{1}{4}Rg_{\Lambda\Psi}=\tfrac{1}{2}g_{\Phi\Psi}g_{\Lambda\Omega}g_{\Gamma\Theta}R^{\Gamma\Phi\Omega\Theta}-\tfrac{1}{N}(g_{\Gamma\Psi}\eta_{\Lambda}{}^{AB}\eta_{\Phi AD}\eta_{[\Theta}{}^{ND}\eta_{\Omega]NB}- \\
-\eta_{(\Lambda}{}^{AB}\eta_{\Psi)AL}\eta_{\Gamma}{}^{CL}\eta_{\Phi CD}\eta_{[\Theta}{}^{ND}\eta_{\Omega]NB})R^{\Gamma\Phi\Omega\Theta}-\tfrac{1}{4}Rg_{\Lambda\Psi}= \\
=-\tfrac{1}{2}R_{\Lambda\Psi}-\tfrac{1}{2}g_{\Phi\Theta}g_{\Lambda\Omega}g_{\Gamma\Psi}R^{\Gamma\Phi\Omega\Theta}+\tfrac{1}{4}g_{\Lambda\Psi}g_{\Gamma\Omega}g_{\Theta\Phi}R^{\Gamma\Phi\Omega\Theta}-\tfrac{1}{4}Rg_{\Lambda\Psi}=-R_{\Lambda\Psi}.
\end{gathered}\tag{12.23}$$

Откуда

$$P_{KL}{}^{MN}:=R_{\Lambda\Psi}\eta^{\Lambda}{}_{KL}\eta^{\Psi MN}=\frac{4}{N}\varepsilon_{KL}{}^{AB}\varepsilon^{MN}{}_{CD}R_{A}{}^{C}{}_{B}{}^{D}-\frac{1}{4}R\varepsilon_{KL}{}^{MN}.\tag{12.24}$$

Кроме того,

$$\begin{gathered}
g^{[\Delta|[\Psi}R^{\Phi]|\Lambda]}\eta_{[\Delta}{}^{AB}\eta_{\Lambda]CB}\eta_{[\Psi}{}^{KL}\eta_{\Phi]ML}= \\
=g^{\Delta\Psi}R^{\Phi\Lambda}(\eta_{\Delta}{}^{AB}\eta_{\Lambda CB}-\tfrac{1}{2}g_{\Delta\Lambda}\delta_C{}^A)\eta_{[\Psi}{}^{KL}\eta_{\Phi]ML}= \\
=g^{\Delta\Psi}R^{\Phi\Lambda}\eta_{\Delta}{}^{AB}\eta_{\Lambda CB}(\eta_{\Psi}{}^{KL}\eta_{\Phi ML}-\tfrac{1}{2}g_{\Psi\Phi}\delta_M{}^K)= \\
=\varepsilon^{ABKL}R^{\Phi\Lambda}\eta_{\Phi ML}\eta_{\Lambda CB}-\tfrac{1}{2}R^{\Delta\Lambda}\eta_{\Delta}{}^{AB}\eta_{\Lambda CB}\delta_M{}^K= \\
=\varepsilon^{ABKL}R^{\Phi\Lambda}\eta_{\Phi ML}\eta_{\Lambda CB}-\tfrac{1}{4}R\delta_C{}^A\delta_M{}^K= \\
=-\varepsilon^{ABKL}(\tfrac{4}{N}\eta^{\Phi GN}\eta^{\Lambda}{}_{DP}R_G{}^D{}_N{}^P-\tfrac{1}{4}Rg^{\Phi\Lambda})\eta_{\Lambda CB}\eta_{\Phi ML}-\tfrac{1}{4}R\delta_C{}^A\delta_M{}^K= \\
=-\tfrac{4}{N}R_G{}^D{}_N{}^P\varepsilon^{ABKL}\varepsilon^{GN}{}_{ML}\varepsilon_{DPCB}+\tfrac{1}{4}R(\varepsilon^{ABKL}\varepsilon_{CBML}-\delta_C{}^A\delta_M{}^K).
\end{gathered}\tag{12.25}$$

3. Третий шаг. Скалярная кривизна R.

$$\begin{gathered}
g^{[\Delta|[\Psi}g^{\Phi]|\Lambda]}\eta_{[\Delta}{}^{AB}\eta_{\Lambda]CB}\eta_{[\Psi}{}^{KL}\eta_{\Phi]ML}= \\
=g^{\Delta\Psi}g^{\Phi\Lambda}\eta_{\Delta}{}^{AB}\eta_{\Lambda CB}(\eta_{\Psi}{}^{KL}\eta_{\Phi ML}-\tfrac{1}{2}g_{\Psi\Phi}\delta_M{}^K)=\varepsilon^{ABKL}\varepsilon_{CBML}-\tfrac{n}{4}\delta_C{}^A\delta_M{}^K.
\end{gathered}\tag{12.26}$$

Теперь остается собрать все результаты вместе и получить аналог разложения на неприводимые ортогональными преобразованиями компоненты. □

Следствие 12.1.

1. *Условия простоты бивектора n-мерного пространства $\mathbb{C}^n$ записываются в виде*

$$p^{[\Lambda\Psi}p^{\Phi\Omega]}=0.\tag{12.27}$$

   Координатам такого бивектора можно сопоставить бесследовую комплексную матрицу такую, что

$$p_L{}^Dp_S{}^L-\frac{1}{N}(p_L{}^Kp_K{}^L)\delta_S{}^D=0.\tag{12.28}$$

2. *Простому бивектору пространства $\mathbb{C}^n$ с условием $p^{\Lambda\Psi}p_{\Lambda\Psi}=0$ можно поставить в соответствие нильпотентный оператор с индексом 2: $p_L{}^Dp_S{}^L=0$.*

Доказательство.

1). Бивектор прост тогда и только тогда, когда имеет место разложение

$$p^{\Lambda\Psi}=x^{\Lambda}y^{\Psi}-y^{\Lambda}x^{\Psi}.\tag{12.29}$$

Поэтому условия (12.27) выполнены автоматически. Обратно, если выполнены условия (12.27), то их можно расписать как

$$p^{\Lambda\Psi}p^{\Phi\Omega}-p^{\Lambda\Phi}p^{\Psi\Omega}+p^{\Psi\Phi}p^{\Lambda\Omega}=0.\tag{12.30}$$

Свернем это уравнение с такими ненулевыми ковекторами t_Ω и z_Φ, что $p^{\Phi\Omega}z_\Phi t_\Omega \neq 0$

$$p^{\Lambda\Psi} = \frac{1}{p^{\Theta\Xi}z_\Theta t_\Xi}(p^{\Lambda\Phi}z_\Phi p^{\Psi\Omega}t_\Omega - p^{\Psi\Phi}z_\Phi p^{\Lambda\Omega}t_\Omega). \tag{12.31}$$

Положим

$$x^\Lambda := \frac{1}{p^{\Theta\Xi}z_\Theta t_\Xi}p^{\Lambda\Phi}z_\Phi, \quad y^\Psi := \frac{1}{p^{\Theta\Xi}z_\Theta t_\Xi}p^{\Psi\Omega}t_\Omega, \tag{12.32}$$

откуда и будет следовать условие простоты бивектора. Поскольку тензор $R_{\Lambda\Psi\Phi\Omega} = p_{\Lambda\Psi}p_{\Phi\Omega}$ удовлетворяет условиям теоремы классификации, то формула (12.28) есть прямое следствие тождества Бианки.

2). В условиях первого пункта добавится соотношение $p^{\Lambda\Psi}p_{\Lambda\Psi} = 0$, которое примет вид ${p_L}^K{p_K}^L = 0$, откуда и следует существование нильпотентного оператора с индексом 2. □

Построим аналог дифференциального тождества Бианки при условии ковариантного постоянства связующих операторов

$$\nabla_{[\Lambda}R_{\Psi\Phi]\Theta\Omega} = 0. \tag{12.33}$$

Для этого свернем его с выражением ${\eta^\Lambda}_{AB}{A^{\Psi\Phi}}_C{}^D{A^{\Theta\Omega}}_K{}^L$

$$\begin{gathered}{\eta^\Lambda}_{AB}{A^{\Psi\Phi}}_C{}^D({\eta_\Lambda}^{MN}A_{\Psi\Phi P}{}^Q - {\eta_\Psi}^{MN}{\eta_\Lambda}^{XQ}\eta_{\Phi XP})\nabla_{MN}{R_Q}^P{}_K{}^L = \\ = -\tfrac{N^2}{16}\nabla_{AB}{R_C}^D{}_K{}^L - \tfrac{N}{8}(\varepsilon_{YCXP}{\varepsilon_{AB}}^{XQ}\nabla^{YD} - {\varepsilon^{YD}}_{XP}{\varepsilon_{AB}}^{XQ}\nabla_{YC}){R_Q}^P{}_K{}^L = 0.\end{gathered} \tag{12.34}$$

Это приведет к дифференциальному спинорному тождеству Бианки

$$\nabla_{AB}{R_C}^D{}_K{}^L = \frac{4}{N}({\varepsilon^{YD}}_{XP}{\varepsilon_{AB}}^{XQ}\nabla_{YC} - \frac{1}{2}{\delta_C}^D{\varepsilon_{AB}}^{XQ}\nabla_{XP}){R_Q}^P{}_K{}^L. \tag{12.35}$$

13 Твисторное уравнение

В этом параграфе рассказывается о том, как построить и решить n-мерное твисторное уравнение, а затем исследовать его свойства. Вывод всех результатов этого параграфа сделан на основе [22, т. 1, с. 419-426, т. 2, с. 56-60, с. 545].

Определим твисторное уравнение как

$$\eta_{\Lambda AB}\nabla_\Psi X^A + \eta_{\Psi AB}\nabla_\Lambda X^A = \frac{2}{n}g_{\Lambda\Psi}{\eta^\Phi}_{AB}\nabla_\Phi X^A. \tag{13.1}$$

Свернем его с $\eta^{\Psi CB}$

$$\begin{gathered}\eta_{\Lambda AB}\nabla^{CB}X^A + \tfrac{n}{2}\nabla_\Lambda X^C = \tfrac{2}{n}\eta^{\Lambda CB}\nabla_{AB}X^A, \\ \nabla_\Lambda X^C - {\eta_\Lambda}^{CB}\nabla_{AB}X^A + \tfrac{n}{2}\nabla_\Lambda X^C = \tfrac{2}{n}{\eta_\Lambda}^{CB}\nabla_{AB}X^A,\end{gathered} \tag{13.2}$$

$$\nabla_\Lambda X^C = \frac{2}{n}\eta_\Lambda{}^{CB}\nabla_{AB}X^A. \tag{13.3}$$

Условия интегрируемости данного уравнения для произвольного X^A имеют вид (доказательство в приложении)

$$C_{\Phi\Psi\Lambda\Delta} = 0, \tag{13.4}$$

что соответствует конформно-плоскому пространству. Пусть Ω - произвольное скалярное поле. Рассмотрим конформное изменение масштаба метрики

$$g_{\Lambda\Psi} \to \hat{g}_{\Lambda\Psi} = \Omega g_{\Lambda\Psi}. \tag{13.5}$$

Поэтому можно положить

$$\hat{\eta}_\Lambda{}^{AB} := \eta_\Lambda{}^{AB}, \quad \hat{\eta}_{\Lambda AB} := \Omega\eta_{\Lambda AB}. \tag{13.6}$$

Потребуем выполнения

$$\nabla_\Lambda\eta_\Psi{}^{AB} = 0, \quad \nabla_\Lambda\eta_{\Psi AB} = 0, \quad \hat{\nabla}_\Lambda\hat{\eta}_\Psi{}^{AB} = 0, \quad \hat{\nabla}_\Lambda\hat{\eta}_{\Psi AB} = 0. \tag{13.7}$$

Это приведет к системе

$$\begin{gathered}
\begin{cases}
(\hat{\nabla}_\Lambda - \nabla_\Lambda)\eta_\Psi{}^{AB} = 0, \\
(\hat{\nabla}_\Lambda - \nabla_\Lambda)\eta_{\Psi AB} = -(\Omega^{-1}\nabla_\Lambda\Omega)\eta_{\Psi AB},
\end{cases} \\
\begin{cases}
Q_{\Lambda\Psi}{}^{\Theta}\eta_\Theta{}^{AB} = Q_{\Lambda K}{}^{A}\eta_\Psi{}^{KB} + Q_{\Lambda K}{}^{B}\eta_\Psi{}^{AK}, \\
Q_{\Lambda\Psi}{}^{\Theta}\eta_{\Theta AB} = -Q_{\Lambda A}{}^{K}\eta_{\Psi KB} - Q_{\Lambda B}{}^{K}\eta_{\Psi AK} + (\Omega^{-1}\nabla_\Lambda\Omega)\eta_{\Psi AB},
\end{cases}
\end{gathered} \tag{13.8}$$

где $Q_{\Lambda\Psi}{}^{\Theta}$ - тензор деформации. Тогда

$$\begin{gathered}
\begin{cases}
Q_{\Lambda(\Psi\Omega)} = \frac{2}{N}g_{\Psi\Omega}Q_{\Lambda K}{}^{K}, \\
Q_{\Lambda(\Psi\Omega)} = -\frac{2}{N}g_{\Psi\Omega}Q_{\Lambda K}{}^{K} + (\Omega^{-1}\nabla_\Lambda\Omega)g_{\Psi\Omega},
\end{cases} \\
\Upsilon_\Lambda := \tfrac{1}{2}\Omega^{-1}\nabla_\Lambda\Omega, \quad Q_{\Lambda(\Psi\Omega)} = g_{\Psi\Omega}\Upsilon_\Lambda, \quad Q_{\Lambda K}{}^{K} = \tfrac{N}{2}\Upsilon_\Lambda, \\
\tilde{Q}_{\Lambda\Psi\Omega} := Q_{\Lambda[\Psi\Omega]}, \quad \tilde{Q}_{\Lambda K}{}^{A} := Q_{\Lambda K}{}^{A} - \tfrac{1}{N}Q_{\Lambda L}{}^{L}\delta_K{}^{A}.
\end{gathered} \tag{13.9}$$

Поэтому

$$\tilde{Q}_{\Lambda\Psi\Omega} = \frac{8}{N}A_{\Psi\Omega A}{}^{K}\tilde{Q}_{\Lambda K}{}^{A}. \tag{13.10}$$

Если потребовать сохранения вида твисторного уравнения

$$\begin{cases}
\nabla_\Lambda X^C = \frac{2}{n}\eta_\Lambda{}^{CB}\nabla_{AB}X^A, \\
\hat{\nabla}_\Lambda\hat{X}^C = \frac{2}{n}\hat{\eta}_\Lambda{}^{CB}\hat{\nabla}_{AB}\hat{X}^A,
\end{cases} \quad \hat{X}^C := X^C, \tag{13.11}$$

то это наложит на тензор деформации условие

$$Q_{\Lambda K}{}^{N} = \frac{2}{n}\eta_\Lambda{}^{CN}\eta_{\Psi CB}Q^{\Psi}{}_{K}{}^{B}. \tag{13.12}$$

Тогда

$$
\begin{gathered}
(\tilde{Q}_{\Lambda K}{}^{N} + \tfrac{1}{2}\Upsilon_{\Lambda}\delta_{K}{}^{N})\eta_{\Phi}{}^{LK}\eta_{\Theta LN} = \tfrac{2}{n}\eta_{\Lambda}{}^{CN}\eta_{\Psi CB}\eta_{\Phi}{}^{LK}\eta_{\Theta LN}(\tilde{Q}^{\Psi}{}_{K}{}^{B} + \tfrac{1}{2}\Upsilon^{\Psi}\delta_{K}{}^{B}),\\
\underbrace{\tilde{Q}_{\Lambda\Phi\Theta} + \Upsilon_{\Lambda}g_{\Phi\Theta}}_{=Q_{\Lambda\Phi\Theta}} = \tfrac{8}{nN}\eta_{\Lambda}{}^{CN}\eta_{\Psi CB}\eta_{\Phi}{}^{LK}\eta_{\Theta LN}\tilde{Q}^{\Psi}{}_{K}{}^{B} + \tfrac{1}{n}(g_{\Lambda\Psi}g_{\Phi\Theta} + g_{\Lambda\Theta}g_{\Phi\Psi} - g_{\Lambda\Phi}g_{\Theta\Psi})\Upsilon^{\Psi},\\
Q_{\Lambda\Phi\Theta} - \tfrac{1}{n}(\Upsilon_{\Lambda}g_{\Phi\Theta} + \Upsilon_{\Phi}g_{\Lambda\Theta} - \Upsilon_{\Theta}g_{\Lambda\Phi}) = \tfrac{8}{nN}\eta_{\Lambda}{}^{CN}\eta_{\Psi CB}\eta_{\Phi}{}^{LK}\eta_{\Theta LN}\tilde{Q}^{\Psi}{}_{K}{}^{B},\\
Q_{\Lambda(\Phi\Theta)} - \tfrac{1}{n}\Upsilon_{\Lambda}g_{\Phi\Theta} = \tfrac{1}{2n}(Q^{\Psi}{}_{\Lambda\Psi} - Q^{\Psi}{}_{\Psi\Lambda})g_{\Phi\Theta},\\
Q^{\Psi}{}_{\Lambda\Psi} - Q^{\Psi}{}_{\Psi\Lambda} = 2(n-1)\Upsilon_{\Lambda},\\
Q_{\Lambda[\Phi\Theta]} + \tfrac{1}{n}(-\Upsilon_{\Phi}g_{\Lambda\Theta} + \Upsilon_{\Theta}g_{\Lambda\Phi}) = \tfrac{4}{nN}\eta_{\Lambda}{}^{CN}\eta_{\Psi CB}\eta_{[\Phi}{}^{LK}\eta_{\Theta]LN}\eta_{[\Gamma|SK|}\eta_{\Omega]}{}^{SB}\tilde{Q}^{\Psi[\Gamma\Omega]} =\\
= \tfrac{1}{2n}(g_{\Lambda\Psi}g_{\Omega\Theta}g_{\Phi\Gamma} - g_{\Lambda\Psi}g_{\Omega\Phi}g_{\Theta\Gamma})\tilde{Q}^{\Psi[\Gamma\Omega]} + \tfrac{4}{n}g_{[\Theta|[\Lambda}g_{\Psi][\Omega}g_{\Gamma]|\Phi]}Q^{\Psi\Gamma\Omega},\\
(1-\tfrac{1}{n})Q_{\Lambda[\Phi\Theta]} + \tfrac{1}{n}(-\Upsilon_{\Phi}g_{\Lambda\Theta} + \Upsilon_{\Theta}g_{\Lambda\Phi}) =\\
= \tfrac{1}{n}(g_{\Lambda\Theta}g_{\Omega\Psi}g_{\Phi\Gamma} - g_{\Psi\Theta}g_{\Omega\Lambda}g_{\Phi\Gamma} + g_{\Phi\Psi}g_{\Omega\Lambda}g_{\Theta\Gamma} - g_{\Lambda\Phi}g_{\Omega\Psi}g_{\Theta\Gamma})Q^{\Psi[\Gamma\Omega]} =\\
= \tfrac{1}{n}((n-1)\Upsilon_{\Phi}g_{\Theta\Lambda} - Q_{\Theta[\Phi\Lambda]} + Q_{\Phi[\Theta\Lambda]} - (n-1)\Upsilon_{\Theta}g_{\Phi\Lambda}).
\end{gathered}
\tag{13.13}
$$

Откуда окончательно имеем

$$
Q_{[\Lambda\Phi\Theta]} = 0,\quad Q_{\Lambda[\Phi\Theta]} = \Upsilon_{\Phi}g_{\Lambda\Theta} - \Upsilon_{\Theta}g_{\Lambda\Phi},\quad Q_{\Lambda\Phi\Theta} = \Upsilon_{\Lambda}g_{\Phi\Theta} + \Upsilon_{\Phi}g_{\Lambda\Theta} - \Upsilon_{\Theta}g_{\Lambda\Phi}. \tag{13.14}
$$

Попробуем решить твисторное уравнение в плоском пространстве ($R_{\Lambda\Psi K}{}^{A}X^{K} = \nabla_{[\Lambda}\nabla_{\Psi]}X^{A} = 0$) при условии ковариантного постоянства связующих операторов ($\nabla_{\Lambda}\eta_{\Psi}{}^{CB} = 0$)

$$
\begin{gathered}
\nabla_{\Lambda}X^{C} = \tfrac{2}{n}\eta_{\Lambda}{}^{CB}\nabla_{AB}X^{A},\\
\nabla_{\Psi}\nabla_{\Lambda}X^{C} = \tfrac{2}{n}\eta_{\Lambda}{}^{CB}\nabla_{AB}(\nabla_{\Psi}X^{A}),\\
\nabla_{\Psi}\nabla_{\Lambda}X^{C} = \tfrac{2}{n}\eta_{\Lambda}{}^{CB}\nabla_{AB}(\tfrac{2}{n}\eta_{\Psi}{}^{AK}\nabla_{LK}X^{L}),\\
\nabla_{(\Psi}\nabla_{\Lambda)}X^{C} = \tfrac{4}{n^{2}}(\eta_{(\Lambda}{}^{CK}\nabla_{\Psi)} - \eta_{(\Lambda}{}^{CB}\eta_{\Psi)AB}\nabla^{AK})\nabla_{LK}X^{L},\\
(1-\tfrac{2}{n})\nabla_{(\Psi}\nabla_{\Lambda)}X^{C} = -\tfrac{2}{n^{2}}g_{\Lambda\Psi}\nabla^{CK}\nabla_{LK}X^{L} = -\tfrac{1}{n}g_{\Lambda\Psi}\nabla^{\Omega}\nabla_{\Omega}X^{C},\\
\nabla_{\Psi}\nabla_{\Lambda}X^{C} = 0.
\end{gathered}
\tag{13.15}
$$

Таким образом $\nabla_{\Lambda}X^{C}$ является константой и представимо как

$$
\nabla_{\Lambda}X^{C} := i\eta_{\Lambda}{}^{CA}\dot{Y}_{A}, \tag{13.16}
$$

где через $\dot{Y}_{A}$, $\dot{X}^{C}$ будем обозначать постоянные спинорные поля. Проинтегрировав это уравнение, получим

$$
X^{C} := \dot{X}^{C} + iR^{CA}\dot{Y}_{A}. \tag{13.17}
$$

Нас будет интересовать случай, когда $X^{C} = 0$. Опуская точки над спинорами, получаем следующее соотношение

$$
X^{C} = -iR^{CA}Y_{A}. \tag{13.18}
$$

С геометрической точки зрения уравнение (13.16), переписанное в терминах операторов γ_Λ, есть ничто иное как деривационное уравнение нормализованного грассманиана [16, уравнение (1.2)], а (13.17) есть уравнение [16, уравнение (2.6)]. Подобная нормализация называется спинорной нормализацией. При n=6 она построена в [7].

Вернемся к твисторному уравнению (13.1), но воспользуемся ковариантно постоянными операторами (11.59), построенными в предыдущем параграфе. Тогда твисторное уравнение (13.16) для действительной реализации примет вид

$$\nabla_{\tilde{\Lambda}} X^{\tilde{\tilde{C}}} := i\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{C}}\tilde{\tilde{A}}}\dot{Y}_{\tilde{\tilde{A}}} =: P_{\tilde{\Lambda}}{}^{\tilde{\tilde{C}}}, \tag{13.19}$$

а для комплексной реализации

$$\begin{gathered}\nabla_\Lambda (X_K)^C = i\eta_\Lambda{}^{C\tilde{\tilde{A}}}\dot{Y}_{\tilde{\tilde{A}}} = (P^*_K)_\Lambda{}^C, \\ (P_K)_{\Lambda A}\nabla_\Psi (X_K)^A + (P_K)_{\Psi A}\nabla_\Lambda (X_K)^A = \tfrac{2}{n} g_{\Lambda\Psi}(P_K)^\Phi{}_A \nabla_\Phi (X_K)^A.\end{gathered} \tag{13.20}$$

Выполним суммирование по K и, определяя $x_\Lambda := \sum\limits_{K=1}^{2N} (P_K)_{\Lambda A}(X_K)^A$, получим конформное уравнение Киллинга

$$\nabla_\Lambda x_\Psi + \nabla_\Psi x_\Lambda = \frac{2}{n} g_{\Lambda\Psi}\nabla_\Phi x^\Phi. \tag{13.21}$$

14 Спинорный формализм для n=6 и n=8

В этом параграфе рассказывается о том, как построить спинорный формализм при малых размерностях. Для наглядности строятся различные геометрические интерпретации некоторых алгебраических соотношений. Показывается, как на основании принципа тройственности Картана перейти к структурным константам алгебры октав. Вывод всех результатов этого параграфа сделан на основе [1] и литературе к ней, [11], [12], [13], [24], [25], [26], [28].

14.1 Основные изоморфизмы

В этом параграфе рассказывается о том, как построить основные изоморфизмы спинорного формализма при n=6. Вывод всех результатов этого параграфа сделан на основе [1], [3].

1. $\mathbb{C}^6 \cong \Lambda^2\mathbb{C}^4$.

Пусть $\alpha, \beta, ... = \overline{1,6}$, $a, b, a_1, b_1, ... = \overline{1,4}$. Тогда из (8.17) будут следовать тождества

$$\frac{1}{2}\eta^\alpha{}_{aa_1}\eta_\beta{}^{aa_1} = \delta_\alpha{}^\beta, \quad \eta^\alpha{}_{aa_1}\eta_\alpha{}^{bb_1} = \delta_{aa_1}{}^{bb_1} := 2\delta_{[a}{}^{[b}\delta_{a_1]}{}^{b_1]}, \tag{14.1}$$

$$r^{\alpha} = 1/2 \cdot \eta^{\alpha}{}_{aa_1} R^{aa_1}, \quad R^{aa_1} = \eta_{\alpha}{}^{aa_1} r^{\alpha}, \tag{14.2}$$

$$\begin{gathered} g^{\alpha\beta} = 1/4 \cdot \eta^{\alpha}{}_{aa_1} \eta^{\beta}{}_{bb_1} \varepsilon^{aa_1bb_1}, \quad \varepsilon^{aa_1bb_1} = \eta_{\alpha}{}^{aa_1} \eta_{\beta}{}^{bb_1} g^{\alpha\beta}, \\ g_{\alpha\beta} = 1/4 \cdot \eta_{\alpha}{}^{aa_1} \eta_{\beta}{}^{bb_1} \varepsilon_{aa_1bb_1}, \quad \varepsilon_{aa_1bb_1} = \eta^{\alpha}{}_{aa_1} \eta^{\beta}{}_{bb_1} g_{\alpha\beta}, \end{gathered} \tag{14.3}$$

определяющие изоморфизм между пространством $\mathbb{C}^6$ и пространством бивекторов $\Lambda^2\mathbb{C}^4$, а $g_{\alpha\beta}$ будет метрическим тензором пространства $\mathbb{C}^6$.

2. $SO(6, \mathbb{C}) \cong SL(4, \mathbb{C}) / \pm 1$.

   Согласно (6.42) собственные ортогональные преобразования представимы как

$$S_{\alpha}{}^{\beta} \eta_{\beta}{}^{aa_1} = \eta_{\alpha}{}^{bb_1} S_b{}^a S_{b_1}{}^{a_1}. \tag{14.4}$$

   При этом $\tilde{E}_a{}^b$ из $(8.19')$ будет тождественным преобразованием, домноженным на мнимую единицу. Для несобственных преобразований справедливо тождество

$$S_{\alpha}{}^{\beta} \eta_{\beta}{}^{aa_1} = \eta_{\alpha bb_1} S^{ba} S^{b_1a_1}. \tag{14.5}$$

3. $so(6, \mathbb{C}) \cong sl(4, \mathbb{C})$.

   Определим

$$A_{\alpha\beta d}{}^{c} = \eta_{[\alpha}{}^{ca} \eta_{\beta]da}. \tag{14.6}$$

   Тогда по аналогии с (10.6) и (12.5) получим

$$T_{\alpha\beta} = A_{\alpha\beta d}{}^{c} T_c{}^d, \quad T_k{}^k = 0, \quad T_{\alpha\beta} = -T_{\alpha\beta}, \tag{14.7}$$

$$\begin{gathered} A_{\alpha\beta d}{}^{c} A^{\alpha\beta}{}_r{}^s = \tfrac{1}{2} \delta_r{}^s \delta_d{}^c - 2\delta_r{}^c \delta_d{}^s, \quad A_{\alpha\beta d}{}^{c} A^{\lambda\mu}{}_c{}^d = 2\delta_{[\alpha}{}^{\mu} \delta_{\beta]}{}^{\lambda}, \\ A_{\alpha\beta d}{}^{c} A_{\gamma}{}^{\beta}{}_r{}^s = \\ = (\eta_{\alpha}{}^{cs} \eta_{\gamma rd} + \eta_{\alpha}{}^{ck} \eta_{\gamma kr} \delta_d{}^s) + 1/2(\eta_{\alpha}{}^{sn} \eta_{\gamma rn} \delta_d{}^c + \eta_{\alpha}{}^{ck} \eta_{\gamma dk} \delta_r{}^s) - 1/4 g_{\alpha\gamma} \delta_r{}^s \delta_d{}^c, \\ T_m{}^n = \tfrac{1}{2} A^{\alpha\beta}{}_m{}^n T_{\beta\alpha}, \quad T_{\beta\alpha} = -T_{\alpha\beta}. \end{gathered} \tag{14.8}$$

Кроме того, интересны тождества для 4-вектора

$$\begin{gathered} e_{\alpha\beta\gamma\delta} = e_{[\alpha\beta\gamma\delta]}, \quad e_{\alpha\beta\gamma\delta} = A_{\alpha\beta b}{}^{a} A_{\gamma\delta d}{}^{c} e_a{}^b{}_c{}^d, \\ e_m{}^n{}_r{}^s = (e_m{}^s \delta_r{}^n + e_r{}^n \delta_m{}^s) - \tfrac{1}{2}(e_r{}^s \delta_m{}^n + e_m{}^n \delta_r{}^s), \\ e_a{}^b := \tfrac{1}{3} e_a{}^k{}_k{}^b, \quad B_{\alpha\beta\gamma\delta r}{}^{k} := A_{\alpha\beta r}{}^{d} A_{\gamma\delta d}{}^{k} + A_{\alpha\beta c}{}^{k} A_{\gamma\delta r}{}^{c}, \quad e_{\alpha\beta\gamma\delta} := B_{\alpha\beta\gamma\delta r}{}^{k} e_k{}^r, \quad e_k{}^k = 0 \end{gathered} \tag{14.9}$$

и 6-вектора

$$\begin{gathered} e_{\alpha\beta\gamma\delta\lambda\mu} = A_{\alpha\beta b}{}^{a} A_{\gamma\delta d}{}^{c} A_{\lambda\mu l}{}^{k} e_a{}^b{}_c{}^d{}_k{}^l, \\ e_a{}^b{}_c{}^d{}_k{}^l = \tfrac{i}{8}(2((4\delta_k{}^b \delta_c{}^l - \delta_k{}^l \delta_c{}^b)\delta_a{}^d + (4\delta_k{}^d \delta_a{}^l - \delta_k{}^l \delta_a{}^d)\delta_c{}^b) - \\ -(4\delta_k{}^b \delta_a{}^l - \delta_k{}^l \delta_a{}^b)\delta_c{}^d - (4\delta_k{}^d \delta_c{}^l - \delta_k{}^l \delta_c{}^d)\delta_a{}^b), \\ e_{\alpha\gamma\lambda\nu\pi\sigma} = 2\eta_{[\alpha}{}^{bb_1} \eta_{\gamma]dd_1} \eta_{[\lambda}{}^{mm_1} \eta_{\nu]xx_1} \eta_{[\pi}{}^{rr_1} \eta_{\sigma]}{}^{ss_1} \cdot i\ \delta_r{}^d \delta_m{}^s \delta_b{}^x \delta_{b_1}{}^{d_1} \delta_{m_1}{}^{x_1} \delta_{r_1}{}^{s_1} = \\ = \tfrac{1}{4} \eta_{[\alpha}{}^{bb_1} \eta_{\gamma]}{}^{dd_1} \eta_{[\lambda}{}^{mm_1} \eta_{\nu]}{}^{xx_1} \eta_{[\pi}{}^{rr_1} \eta_{\sigma]}{}^{ss_1} \cdot i\ \varepsilon_{rb_1dd_1} \varepsilon_{mr_1ss_1} \varepsilon_{bm_1xx_1}. \end{gathered} \tag{14.10}$$

Доказательство этих тождеств вынесено в приложение.

14.2 О классификации тензоров с симметриями тензора кривизны для n=6. Спиноры кривизны

В этом параграфе рассказывается о том, как сопоставить тензорам, обладающим симметриями тензора кривизны, их спинорные аналоги. Это упрощает проблему классификации таких тензоров для малых размерностей. Вывод всех результатов этого параграфа сделан на основе [2].

Теорема 14.1. *Классификацию битензора, обладающего свойствами*

$$R_{\alpha\beta\gamma\delta} = R_{[\alpha\beta][\gamma\delta]}\,, \quad R_{\alpha\beta\gamma\delta} = R_{\gamma\delta\alpha\beta}, \quad R_{\alpha\beta\gamma\delta} + R_{\alpha\delta\beta\gamma} + R_{\alpha\gamma\delta\beta} = 0 \tag{14.11}$$

и принадлежащего касательному расслоению $\tau(\mathbb{C}V_6)$ над шестимерным аналитическим псевдоримановым пространством $\mathbb{C}V_6$, можно свести к классификации тензора ${R_a}^b{}_c{}^d$ 4-мерного комплексного векторного пространства такого, что

$$R_{\alpha\beta\gamma\delta} = A_{\alpha\beta d}{}^c A_{\gamma\delta r}{}^s {R_c}^d{}_s{}^r. \tag{14.12}$$

Кроме того, выполнены следующие соотношения

$${R_k}^k{}_s{}^r = {R_s}^r{}_k{}^k = 0, \quad {R_c}^d{}_s{}^r = {R_s}^r{}_c{}^d. \tag{14.13}$$

Разложение

$${R_c}^d{}_s{}^r = {C_c}^d{}_s{}^r - P_{cs}{}^{dr} - \frac{1}{40} \cdot R(3{\delta_s}^d{\delta_c}^r - 2{\delta_s}^r{\delta_c}^d) \tag{14.14}$$

соответствует разложению тензора

$$R_{\alpha\beta}{}^{\gamma\delta} = C_{\alpha\beta}{}^{\gamma\delta} + R_{[\alpha}{}^{[\gamma} g_{\beta]}{}^{\delta]} - 1/10 R g_{[\alpha}{}^{[\gamma} g_{\beta]}{}^{\delta]} \tag{14.15}$$

на неприводимые ортогональными преобразованиями компоненты, которые будут удовлетворять следующим соотношениям

$$P_{cs}{}^{rd} = -4(R_{[c}{}^{[r}{}_{s]}{}^{d]} + R_k{}^{[r}{}_{[c}{}^{|k|}\delta_{s]}{}^{d]}), \tag{14.16}$$

$${C_c}^d{}_s{}^r = R_{(c}{}^{(d}{}_{s)}{}^{r)} + \frac{1}{40} \cdot R\delta_{(s}{}^d \delta_{c)}{}^r, \quad {C_c}^d{}_s{}^r = C_{(c}{}^{(d}{}_{s)}{}^{r)}, \tag{14.17}$$

$$R = R_\beta{}^\beta = -2 \cdot {R_k}^r{}_r{}^k, \quad P_{kc}{}^{kd} = 1/2 \cdot R{\delta_c}^d, \tag{14.18}$$

$${R_l}^d{}_s{}^l = -\frac{1}{8} \cdot R{\delta_s}^d, \tag{14.19}$$

последнее из которых является эквивалентом тождества Бианки.

Доказательство. Верно следующее равенство

$$R_{\alpha\beta\gamma\delta} = 1/16 \cdot \eta_\alpha{}^{aa_1}\eta_\beta{}^{bb_1}\eta_\gamma{}^{cc_1}\eta_\delta{}^{dd_1}R_{aa_1bb_1cc_1dd_1}. \tag{14.20}$$

Положим

$$R_c{}^d{}_s{}^r := \frac{1}{4}R_{ck}{}^{dk}{}_{st}{}^{rt}, \quad R_{\beta\gamma} = \frac{1}{4}\eta_\beta{}^{cs}\eta_{\gamma rd} \cdot P_{cs}{}^{rd}. \tag{14.21}$$

Из этого следует, что

$$R_{\alpha\beta\gamma\delta} = A_{\alpha\beta d}{}^c A_{\gamma\delta r}{}^s R_c{}^d{}_s{}^r, \tag{14.22}$$

$$\begin{aligned} R_{\beta\delta} = R_{\alpha\beta}{}^\alpha{}_\delta = A_{\alpha\beta d}{}^c A^\alpha{}_{\delta r}{}^s R_c{}^d{}_s{}^r = (\eta_\beta{}^{cs}\eta_{\delta rd} + \eta_\beta{}^{ck}\eta_{\delta kr}\delta_d{}^s)R_c{}^d{}_s{}^r = \\ = \tfrac{1}{4}\eta_\beta{}^{cs}\eta_{\delta rd} \cdot 4(R_{[c}{}^{[d}{}_{s]}{}^{r]} - R_{[c}{}^k{}_{|k|}{}^{[r}\delta_{s]}{}^{d]}). \end{aligned} \tag{14.23}$$

Положим

$$P_{cs}{}^{rd} := -4(R_{[c}{}^{[r}{}_{s]}{}^{d]} - R_{[c}{}^k{}_{|k|}{}^{[r}\delta_{s]}{}^{d]}) \ , \tag{14.24}$$

тогда

$$R_{\beta\delta} = \frac{1}{4}\eta_\beta{}^{cs}\eta_{\delta rd}P_{cs}{}^{rd}. \tag{14.25}$$

Поэтому скалярная кривизна имеет вид

$$R = R_\beta{}^\beta = \tfrac{1}{4}\eta_\beta{}^{cc_1}\eta^\beta{}_{aa_1}P_{cc_1}{}^{aa_1} = \tfrac{1}{4}\varepsilon_{aa_1}{}^{cc_1}P_{cc_1}{}^{aa_1} = \tfrac{1}{2}P_{aa_1}{}^{aa_1} = -2R_k{}^r{}_r{}^k, \tag{14.26}$$

и выполнено условие

$$\begin{aligned} P_{ks}{}^{kd} = -4(R_{[k}{}^{[k}{}_{s]}{}^{r]} + R_{[r}{}^k{}_{|k|}{}^{[r}\delta_{s]}{}^{d]}) = \\ = -4(-\tfrac{1}{2}R_k{}^d{}_s{}^k + \tfrac{1}{4}(R_k{}^r{}_r{}^k\delta_s{}^d + 4R_k{}^d{}_s{}^k - 2R_k{}^d{}_s{}^k)) = -R_k{}^r{}_r{}^k\delta_s{}^d = \tfrac{1}{2}R\delta_s{}^d. \end{aligned} \tag{14.27}$$

Тождество Бианки можно переписать следующим образом

$$(A_{\alpha\beta d}{}^c A_{\gamma\delta r}{}^s + A_{\alpha\gamma d}{}^c A_{\delta\beta r}{}^s + A_{\alpha\delta d}{}^c A_{\beta\gamma r}{}^s) \cdot R_c{}^d{}_s{}^r = 0. \tag{14.28}$$

Свернув это уравнение с $A^{\alpha\beta}{}_t{}^l A^{\gamma\delta}{}_m{}^n$, получим, что

$$4R_k{}^l{}_m{}^k\delta_t{}^n + 4R_r{}^n{}_t{}^r\delta_m{}^l - 2R_k{}^l{}_t{}^k\delta_m{}^n - 2R_k{}^n{}_m{}^k\delta_t{}^l - 2R_k{}^r{}_r{}^k\delta_t{}^n\delta_m{}^l + R_r{}^k{}_k{}^r\delta_m{}^n\delta_t{}^l = 0. \tag{14.29}$$

Свертка этого уравнения с $\delta_n{}^t$ и приведет нас к спинорному аналогу тождества Бианки. При этом все 15 существенных уравнений сохранены. Пусть

$$C_{\alpha\beta}{}^{\gamma\delta} := A_{\alpha\beta d}{}^c A^{\gamma\delta}{}_r{}^s C_c{}^d{}_s{}^r, \tag{14.30}$$

тогда

$$C_{\alpha\beta}{}^{\gamma\delta} := R_{\alpha\beta}{}^{\gamma\delta} - R_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]} + 1/10Rg_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]}, \tag{14.31}$$

$$R_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]} = A_{\alpha\beta d}{}^c A^{\gamma\delta}{}_r{}^s \cdot \frac{1}{4}(P_{sc}{}^{dr} - 1/2R\delta_s{}^d\delta_c{}^r + \frac{1}{4}R\delta_s{}^r\delta_c{}^d), \tag{14.32}$$

$$g_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]} = A_{\alpha\beta d}{}^c A^{\gamma\delta}{}_r{}^s \cdot \frac{1}{4}(1/2\delta_s{}^r\delta_c{}^d - 2\delta_s{}^d\delta_c{}^r), \tag{14.33}$$

откуда получим разложение на неприводимые ортогональными преобразованиями компоненты. Все выкладки приведены в приложении. □

Замечание 14.1. *При n=6 возможно два варианта классификационных схем спинора Вейля:*

$$1.\quad C_k{}^l{}_m{}^n \phi_l{}^k = \lambda \phi_t{}^n, \qquad 2.\quad C_k{}^l{}_m{}^n \phi^{km} = \lambda \phi^{ln}. \tag{14.34}$$

Следствие 14.1. *1. Условия простоты бивектора 6-мерного пространства $\mathbb{C}^6$ записываются как*

$$p_{[\alpha\beta} p_{\gamma\beta]} = 0. \tag{14.35}$$

Координатам такого бивектора можно сопоставить бесследовую комплексную матрицу 4×4 такую, что выполнено условие

$$p_l{}^d p_s{}^l - 1/4(p_l{}^k p_k{}^l)\delta_s{}^d = 0. \tag{14.36}$$

2. Простому бивектору пространства $\mathbb{C}^6$, построенному на изотропных векторах ($p^{\alpha\beta}p_{\alpha\beta} = 0$), можно сопоставить вырожденную нуль-пару Розенфельда: ковектор и вектор пространства $\mathbb{C}^4$, свертка которых есть нуль. При этом указанные вектор и ковектор определятся с точностью до комплексного множителя.

Доказательство. Доказательство аналогично доказательству следствия 12.1. Единственное отличие заключается в том, что изотропные векторы r^α пространства $\mathbb{C}^6$ представимы в виде $r^\alpha \eta_\alpha{}^{ab} = R^{ab} = X^a Y^b - X^b Y^a$. Такое представление возможно из-за выполнения тождества для изотропного r^α ($r^\alpha r_\alpha = 0$)

$$24 R^{[ab} R^{cd]} = \varepsilon_{klmn} R^{kl} R^{mn} \varepsilon^{abcd} = 4(r^\alpha r_\alpha)\varepsilon^{abcd} = 0, \tag{14.37}$$

где ε_{abcd} - тензор вида (8.17). При этом такой тензор кососимметричен по всем 4 индексам и удовлетворяет соотношениям

$$\begin{gathered}\varepsilon_{abcd}\varepsilon^{klmn} = 24\delta_{[a}{}^k \delta_b{}^l \delta_c{}^m \delta_{n]}{}^d, \quad \varepsilon_{abcd}\varepsilon^{almn} = 6\delta_{[b}{}^l \delta_c{}^m \delta_{n]}{}^d, \quad \varepsilon_{abcd}\varepsilon^{abmn} = 2\delta_{[c}{}^m \delta_{n]}{}^d, \\ \varepsilon_{abcd}\varepsilon^{abcn} = \delta_d{}^n, \quad \varepsilon_{abcd}\varepsilon^{abcd} = 24.\end{gathered} \tag{14.38}$$

Положим $x^\alpha \eta_\alpha{}^{ab} := X^a Y^b - Y^a X^b$, $y^\alpha \eta_\alpha{}^{ab} := Z^a T^b - T^a Z^b$. Из условия $p^{\alpha\beta}p_{\alpha\beta} = 0$ будет следовать $x^\alpha y_\alpha = 0$. Это означает, что $\varepsilon_{abcd} X^a Y^b Z^c T^d = 0$. Поэтому векторы $X^a,\ \ Y^b,\ \ Z^c,\ \ T^d$ линейно зависимы. Положим $T^a := \alpha X^a + \beta Y^a + \gamma Z^a$ и получим

$$\begin{gathered} p_a{}^b := \tfrac{1}{2} A_{\alpha\beta a}{}^b p_{\alpha\beta} = \tfrac{1}{4}\eta_\alpha{}^{kb}\eta_\beta{}^{cd}\varepsilon_{cdka}(x^\alpha y^\beta - y^\alpha x^\beta) = \\ = \tfrac{1}{4}((X^k Y^b - X^b Y^k) Z^c T^d - X^c Y^d (Z^k T^b - T^k Z^b))\varepsilon_{cdka} = \\ = Z^c(\beta Y^d) X^k \varepsilon_{cdka} Y^b - Z^c(\alpha X^d) Y^k \varepsilon_{cdka} X^b - X^c Y^d Z^k \varepsilon_{cdka}(\alpha X^b + \beta Y^b + \gamma Z^b) + \\ + X^c Y^d (\gamma Z^k)\varepsilon_{cdka} Z^b = \underbrace{X^c Y^d Z^k \varepsilon_{cdka}}_{:=M_a} \underbrace{(-2\alpha X^b - 2\beta Y^b)}_{:=N^b} = M_a N^b. \end{gathered} \tag{14.39}$$

При этом N^b и M_a определены с точностью до преобразования

$$N^b \longmapsto e^{\phi} N^b, \quad M_a \longmapsto e^{-\phi} M_a. \tag{14.40}$$

□

Отметим, что пара (N^b, M_a) будет является нуль-парой Розенфельда [19]. В пространстве $\mathbb{C}P^4 =' \mathbb{C}^4/'\mathbb{C}$ (где $'\mathbb{C}^s = \mathbb{C}^s/0$) N^b определит точку, а M_a - плоскость с условием инцидентности $M^b N_a = 0$. Поэтому можно построить пространство $\mathbb{C}\Pi^4 =' \mathbb{C}^{*4}/'\mathbb{C}$, двойственное пространству $\mathbb{C}P^4$. Тогда пространство $\mathbb{C}P^4 \times \mathbb{C}\Pi^4$ будет пространством нуль-пар Розенфельда. Следует отметить, что такие пространства изучались впервые Синцовым [23] и Котельниковым [14].

Подставим теперь $\varepsilon_{ab}{}^{cd} = 2\delta_{[a}{}^{c}\delta_{b]}{}^{d}$ в (12.35). Тогда дифференциальные тождества Бианки упростятся до

$$\nabla_{[cm} R_{t]}{}^{k}{}_{r}{}^{s} = \delta_{[m}{}^{k} \nabla_{c|n|} R_{t]}{}^{n}{}_{r}{}^{s}. \tag{14.41}$$

Свернем это уравнение с $\delta_k{}^c$

$$\nabla_{c(m} R_{t)}{}^{c}{}_{r}{}^{s} = 0, \tag{14.42}$$

свертка которого с $\delta_s{}^m$ даст

$$\nabla_{cm} R_{t}{}^{c}{}_{r}{}^{m} = 1/8 \nabla_{rt} R, \tag{14.43}$$

что является спинорным аналогом

$$\nabla^{\alpha}(R_{\alpha\beta} - 1/2 R g_{\alpha\beta}) = 0. \tag{14.44}$$

14.3 Геометрическое представление твистора в $\mathbb{R}^6_{(2,4)}$

В этом параграфе рассказывается о том, как построить геометрическую интерпретацию изотропного спинора (твистора) на изотропном конусе $\mathbb{R}^6_{(2,4)}$. *Это представление аналогично представлению спинора в* $\mathbb{R}^4_{(1,3)}$ *с той лишь разницей, что размерность флага и полотнища увеличиваются на единицу. Вывод всех результатов этого параграфа сделан на основе [5].*

Пусть метрика пространства $\mathbb{R}^6_{(2,4)}$ имеет вид

$$dS^2 = dT^2 + dV^2 - dW^2 - dX^2 - dY^2 - dZ^2, \tag{14.45}$$

и пусть задано сечение светового конуса K_6

$$T^2 + V^2 - W^2 - X^2 - Y^2 - Z^2 = 0 \tag{14.46}$$

плоскостью V+W=1. Рассмотрим стереографическую проекцию этого сечения на плоскость (V=0,W=1) с полюсом $N(0,\frac{1}{2},\frac{1}{2},0,0,0)$ так, что точке P(T,V,W,X,Y,Z) соответствует $p(t,0,1,x,y,z)$ на плоскости V=0. Тогда

$$T/t = X/x = Y/y = Z/z = -\frac{(V-\frac{1}{2})}{\frac{1}{2}}. \tag{14.47}$$

Сделаем замену

$$\varsigma = ix - y, \quad \omega = i(t+z), \quad \eta = i(t-z) \tag{14.48}$$

и получим

$$\varsigma = \frac{-iX+Y}{2V-1}, \quad \eta = \frac{-i(T+Z)}{2V-1}, \quad \omega = \frac{i(Z-T)}{2V-1}. \tag{14.49}$$

Поэтому индуцированная в сечении метрика имеет вид

$$ds^2 := dT^2 - dX^2 - dY^2 - dZ^2 = -\frac{d\varsigma d\bar{\varsigma} + d\omega d\eta}{(\varsigma\bar{\varsigma} + \eta\omega)^2}. \tag{14.50}$$

Доказательство этого факта вынесено в приложение. Положим

$$X := \begin{pmatrix} \omega & \varsigma \\ -\bar{\varsigma} & \eta \end{pmatrix}, \quad dX := \begin{pmatrix} d\omega & d\varsigma \\ -d\bar{\varsigma} & d\eta \end{pmatrix}, \quad \frac{\partial}{\partial X} := \begin{pmatrix} \frac{\partial}{\partial\omega} & \frac{\partial}{\partial\varsigma} \\ -\frac{\partial}{\partial\bar{\varsigma}} & \frac{\partial}{\partial\eta} \end{pmatrix}. \tag{14.51}$$

Тогда

$$ds^2 = -\frac{det(dX)}{(det(X))^2}, \quad \bar{X}^T + X = 0. \tag{14.52}$$

Рассмотрим группу дробно-линейных преобразований L

$$\tilde{X} = (AX+B)(CX+D)^{-1}, \quad S := \begin{pmatrix} A & B \\ C & D \end{pmatrix}, \quad detS = 1. \tag{14.53}$$

Условие действительности, накладываемое на X ($X^* + X = 0$), даст подгруппу унитарных дробно-линейных преобразований $LU(2,2)$ так, что матрица S удовлетворяет условию (см. приложение)

$$S^*\hat{E}S = \hat{E}, \quad \hat{E} := \begin{pmatrix} 0 & E \\ E & 0 \end{pmatrix}. \tag{14.54}$$

Определим далее в специальном базисе [22, т. 2, с. 83, с. 361]

$$\begin{array}{rcccl}
R^{12} & = & 1/\sqrt{2}(V+W) & = & \omega^0\xi^1 - \omega^1\xi^0, \\
R^{14} & = & i/\sqrt{2}(T+Z) & = & \omega^0\bar{\eta}^{0'} - \xi^0\bar{\pi}^{0'}, \\
R^{34} & = & 1/\sqrt{2}(V-W) & = & \bar{\pi}^{0'}\bar{\eta}^{1'} - \bar{\pi}^{1'}\bar{\eta}^{0'}, \\
R^{23} & = & i/\sqrt{2}(Z-T) & = & \xi^1\bar{\pi}^{1'} - \omega^1\bar{\eta}^{1'}, \\
R^{24} & = & 1/\sqrt{2}(Y+iX) & = & \omega^1\bar{\eta}^{0'} - \xi^1\bar{\pi}^{0'}, \\
R^{13} & = & 1/\sqrt{2}(Y-iX) & = & \xi^0\bar{\pi}^{1'} - \omega^0\bar{\eta}^{1'}.
\end{array} \tag{14.55}$$

Эта формула примечательна тем, что в ней показано выражение бивектора R^{ab} через его спинорные компоненты. Положим

$$X := YZ^{-1}, \quad \tilde{Y} = AY + BZ, \quad \tilde{Z} = CY + DZ, \quad Y = \begin{pmatrix} \omega^0 & \xi^0 \\ \omega^1 & \xi^1 \end{pmatrix}, \quad Z = \begin{pmatrix} \bar{\pi}_0 & \bar{\eta}_0 \\ \bar{\pi}_1 & \bar{\eta}_1 \end{pmatrix}, \tag{14.56}$$

тогда

$$R :=\| R^{ab} \|= \begin{pmatrix} (detY)J & YZ^{-1}(detZ)J \\ -(YZ^{-1}(detZ)J)^T & (detZ)J \end{pmatrix} = \begin{pmatrix} Y \\ Z \end{pmatrix} J \begin{pmatrix} Y^T & Z^T \end{pmatrix},$$
$$J := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \tilde{R} = SRS^T. \tag{14.57}$$

Положим

$$\tilde{S} := \tilde{I}S\tilde{I}^{-1}, \quad \tilde{I} := \frac{1}{\sqrt{2}} \begin{pmatrix} E & E \\ -E & E \end{pmatrix}, \tag{14.58}$$

тогда

$$\tilde{S}^* \tilde{E} \tilde{S} = \tilde{E}. \tag{14.59}$$

Матрицы S образуют группу, изоморфную $SU(2,2)$, следовательно матрицы $\tilde{S}$ образуют саму группу $SU(2,2)$. Назовем преобразования с матрицей из группы $LU(2,2)$ твисторными преобразованиями. Ввиду двулистности накрытия связной компоненты единицы группы $SO(2,4)$ (которая обозначается через $SO^+(2,4)$) группой $SU(2,2)$ и двулистности накрытия группы конформных преобразований $\mathbb{C}_+^{\uparrow 4}(1,3)$ [22, т.2, с.359] группой $SO^+(2,4)$ следует существование цепочки изоморфизмов

$$SU(2,2)/\{\pm 1; \pm i\} \cong LU(2,2) \cong \mathbb{C}_+^{\uparrow 4}(1,3) \cong SO^e(2,4)/\pm 1. \tag{14.60}$$

Это означает, что группа $LU(2,2)$ исчерпывает все конформные преобразования из группы $\mathbb{C}_+^{\uparrow 4}(1,3)$. При этом матрица S восстанавливается с точностью до множителя λ такого, что $\lambda^4 = 1$ (det(S)=1), откуда и появляется указанная неоднозначность. На основании того, что верны тождества

$$Y = AX + B \quad \Rightarrow \quad dX = AdY, \quad Y = X^{-1} \quad \Rightarrow \quad dX = -X^{-1}dXX^{-1}, \tag{14.61}$$

где A и B - некоторые постоянные матрицы, имеем

$$\tilde{Z}^* \, d\tilde{X} \, \tilde{Z} = Z^* \, dX \, Z. \tag{14.62}$$

Это уравнение инвариантно относительно преобразований из группы LU(2,2). Доказательство этого факта рассмотрено в приложении. Другой инвариант можно получить, рассматривая тождества

$$Y = AX + B \quad \Rightarrow \quad \frac{\partial}{\partial X} = A^T \frac{\partial}{\partial Y}, \quad Y = X^{-1} \quad \Rightarrow \quad \frac{\partial}{\partial X} = -Y^T \frac{\partial}{\partial Y} Y^T, \tag{14.63}$$

где A и B - тоже некоторые постоянные матрицы. Он будет иметь вид

$$\tilde{Z}^{-1} \frac{\partial}{\partial \tilde{X}^T} \tilde{Z}^{*\ -1} = Z^{-1} \frac{\partial}{\partial X^T} Z^{*\ -1}. \tag{14.64}$$

Доказательство можно найти в приложении. Это означает, что существует действительный касательный вектор $\tilde{L}$ к гиперболоиду, полученному сечением конуса K_6 плоскостью V+W=1. Этот вектор инвариантен относительно преобразований базиса из группы LU(2,2) (т.е. координатно-независимый на касательном пространстве к данному гиперболоиду) и однозначно определенный матрицей

$$\begin{gathered}\hat{L} := \tfrac{1}{\sqrt{2}}(Z^{-1} \tfrac{\partial}{\partial X^T} Z^{*\ -1} - \bar{Z}^{-1} \tfrac{\partial}{\partial X} Z^{T\ -1}) = \\ = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} (\tfrac{\partial}{\partial \omega}(-\bar{\eta}^0 \pi^0 + \eta^0 \bar{\pi}^0) + \tfrac{\partial}{\partial \eta}(-\bar{\eta}^1 \pi^1 + \eta^1 \bar{\pi}^1) + \\ + \tfrac{\partial}{\partial \xi}(-\bar{\eta}^1 \pi^0 + \eta^0 \bar{\pi}^1) + \tfrac{\partial}{\partial \bar{\xi}}(\bar{\eta}^0 \pi^1 - \eta^1 \bar{\pi}^0)) \cdot \tfrac{1}{(det(Z))^2 \sqrt{2}} := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \tilde{L}\end{gathered} \tag{14.65}$$

Норма этого вектора в нашей метрике будет такой

$$\parallel \tilde{L} \parallel = -\frac{1}{2(det(Y))^2} = -\frac{1}{(V+W)^2} \tag{14.66}$$

Назовем изотропный вектор k вектором, имеющим единичную протяженность первого типа [22, т. 1, с. 57] в том случае, когда k будет задавать точку на изотропном конусе, принадлежащую сечению плоскостью V+W=1. Тогда $\parallel \tilde{L} \parallel = -1$ и любой изотропной вектор K, коллинеарный k, определится как

$$K = (- \parallel \tilde{L} \parallel)^{\frac{1}{2}} k. \tag{14.67}$$

Однако, при V=-W получаются векторы с бесконечной протяженностью первого типа. Чтобы научиться их различать можно задавать сечение K_6 плоскостью T+Z=1 и ввести подобным образом некоторый вектор $\tilde{\tilde{L}}$ с нормой

$$\parallel \tilde{\tilde{L}} \parallel = -\frac{1}{(T+Z)^2}. \tag{14.68}$$

Назовем изотропный вектор k вектором, имеющим единичную протяженность второго типа в том случае, когда k будет задавать точку на изотропном конусе, принадлежащую сечению плоскостью T+Z=1 и протяженность первого типа не будет конечной. Определим протяженность вектора K как конечную протяженность первого типа, а если такой не существует, то как протяженность второго типа. Отметим, что вектор $\tilde{L}$ не является координатно-независимым в пространстве $\mathbb{R}^6_{(2,4)}$, хотя и является инвариантом касательного пространства к гиперболоиду, полученному сечением конуса K_6 плоскостью V+W=1. Теперь появилась возможность наглядно изобразить твистор в пространстве $\mathbb{R}^6{}_{(2,4)}$. Рассмотрим пару векторов из $\mathbb{R}^6{}_{(2,4)}$ равной протяженности

$$K^\alpha = \eta^\alpha{}_{ab}\; iT^{[a}X^{b]}, \quad N^\alpha = \eta^\alpha{}_{ab}\; T^{[a}Z^{b]} \;\Rightarrow\; K^\alpha K_\alpha = 0, \quad N^\alpha K_\alpha = 0, \quad N^\alpha N_\alpha = 0. \tag{14.69}$$

Выберем вектор Y^a таким образом, чтобы были выполнены условия

$$\begin{gathered} Y^aY_a = 0, \quad Y^aX_a = 0, \quad Y^aZ_a = 0, \\ \varepsilon^{abcd}X_aY_bZ_cT_d = X^cZ_cY^dT_d = 1, \quad \varepsilon^{abcd} = 24X^{[a}Y^bZ^cT^{d]}. \end{gathered} \tag{14.70}$$

Таким образом получится базис из векторов X^a, Y^a, Z^a, T^a (напомним, что $Y_a = s_{aa'}Y^{a'}$)

$$\begin{gathered} Y^aY_a = 0, \quad Y^aX_a = 0, \quad Y^aZ_a = 0, \quad X^aX_a = 0, \\ X^aT_a = 0, \quad Z^aZ_a = 0, \quad Z^aT_a = 0, \quad T^aT_a = 0. \end{gathered} \tag{14.71}$$

Откуда

$$\varepsilon_{abcd}T^{[c}Y^{d]} = -2X_{[a}Z_{b]}. \tag{14.72}$$

Поэтому векторы

$$L^\alpha = \eta^\alpha{}_{ab}(-T^{[a}Y^{b]} + X^{[a}Z^{b]}), \quad M^\alpha = \eta^\alpha{}_{ab}(-i)(T^{[a}Y^{b]} + X^{[a}Z^{b]}) \tag{14.73}$$

удовлетворяют соотношениям

$$\begin{gathered} L^\alpha = \bar{L}^\alpha, \quad M^\alpha = \bar{M}^\alpha, \quad L^\alpha K_\alpha = 0, \quad L^\alpha M_\alpha = 0, \quad M^\alpha K_\alpha = 0, \\ L^\alpha N_\alpha = 0, \quad M^\alpha N_\alpha = 0, \quad L^\alpha L_\alpha = -2, \quad M^\alpha M_\alpha = -2. \end{gathered} \tag{14.74}$$

Вот теперь мы можем построить тривектор

$$P^{\alpha\beta\gamma} = 6K^{[\alpha}N^\beta L^{\gamma]}. \tag{14.75}$$

Зная K^α, мы знаем T^a и X^a с точностью до

$$X^a \longmapsto \lambda_1 X^a + \mu_1 T^a, \quad T^a \longmapsto \nu_1 X^a + \xi_1 T^a, \quad det\begin{pmatrix} \lambda_1 & \mu_1 \\ \nu_1 & \xi_1 \end{pmatrix} = 1. \tag{14.76}$$

А если нам известен N^α, то произвол в выборе T^a и Z^a таков

$$Z^a \longmapsto \lambda_2 Z^a + \mu_2 T^a, \quad T^a \longmapsto \nu_2 Z^a + \xi_2 T^a, \quad det\begin{pmatrix} \lambda_2 & \mu_2 \\ \nu_2 & \xi_2 \end{pmatrix} = 1. \tag{14.77}$$

Поэтому $\nu_1 = \nu_2 = 0$ и $\xi_2 = \xi_1$. Для Y^a получим

$$Y^a \longmapsto \alpha X^a + \beta Y^a + \gamma Z^a + \delta T^a. \tag{14.78}$$

Если теперь потребовать, чтобы два таких базиса связаны преобразованием из группы LU(2,2), то получим

$$\begin{aligned} X^a &\longmapsto \tau^{-1} X^a + \mu T^a, & T^a &\longmapsto \tau T^a, \\ Z^a &\longmapsto \tau^{-1} Z^a + \chi T^a, & Y^a &\longmapsto -\bar\chi X^a + \tau Y^a - \bar\mu Z^a + \delta T^a, \end{aligned} \tag{14.79}$$

$$\bar\chi\mu + \bar\mu\chi + \tau\bar\delta + \bar\tau\delta = 0, \quad \tau\bar\tau = 1. \tag{14.80}$$

Откуда

$$\begin{aligned} X^{[a}T^{b]} &\longmapsto X^{[a}T^{b]} \Leftrightarrow K^\alpha \longmapsto K^\alpha \\ Z^{[a}T^{b]} &\longmapsto Z^{[a}T^{b]} \Leftrightarrow N^\alpha \longmapsto N^\alpha, \end{aligned} \tag{14.81}$$

$$\begin{aligned} T^{[a}Y^{b]} &\longmapsto -\tau\bar\chi T^{[a}X^{b]} + \tau^2 T^{[a}Y^{b]} - \bar\mu\tau T^{[a}Z^{b]} \\ X^{[a}Z^{b]} &\longmapsto \tau^{-2} X^{[a}Z^{b]} + \tau^{-1}\chi X^{[a}T^{b]} + \tau^{-1}\mu T^{[a}Z^{b]}. \end{aligned} \tag{14.82}$$

Положим $\tau =: e^{i\Theta}$, тогда

$$\begin{aligned} L^\alpha &\longmapsto L^\alpha \cos(2\Theta) + M^\alpha \sin(2\Theta) - i(\bar\chi\tau - \bar\tau\chi)K^\alpha + (\mu\bar\tau + \tau\bar\mu)N^\alpha, \\ M^\alpha &\longmapsto M^\alpha \cos(2\Theta) - L^\alpha \sin(2\Theta) + (\bar\chi\tau + \bar\tau\chi)K^\alpha - i(\mu\bar\tau - \tau\bar\mu)N^\alpha. \end{aligned} \tag{14.83}$$

Таким образом 3-полуплоскость натянутая, на векторы $K^\alpha, N^\alpha, L^\alpha$, будет координатно-независима в пространстве $\mathbb{R}^6_{(2,4)}$. Итак, нашу конструкцию можно представить в следующем виде. Протяженность первого типа векторов K^α и N^α должна быть одинакова. K^α и N^α определяют 2-плоскость, множество векторов которой с протяженностью первого типа, равной протяженности вектора K^α и началом, совпадающим с началом вектора K^α, назовем флагштоком. K^α,N^α,L^α определят 3-полуплоскость, которую назовем полотнищем флага. Таким образом, зная K^α и N^α, мы знаем твистор T^a с точностью до фазы Θ. В свою очередь, 2Θ - это угол поворота полотнища флага - 3-полуплоскости $P^{\alpha\beta\gamma}$ - в 2-плоскости (L^α, M^α) вокруг флагштока - 2-плоскости (N^α, K^α). Поэтому, поворот флага на 2π приведет к твистору $-T^a$, и только поворот на 4π вернет нашу конструкцию к исходному состоянию. Кроме того, коллинеарные твисторы определяются протяженностью вектора K^α так, что при преобразовании $T^a \longmapsto rT^a$, $Y^a \longmapsto r^{-1}Y^a$ $(r \in R\backslash\{0\})$ флагшток умножается на r, а полотнище остается неизменным. Следует, наконец, отметить тот факт, что указанная геометрическая структура однозначно восстанавливается по твистору T^a. В случае бесконечной протяженности первого типа вектора K^α рассматривается конус $K_4 \subset K_6$, на котором лежит вектор N^α. Но ненулевые векторы K^α и N^α имеют конечную протяженность второго типа, давая геометрическую интерпретацию спинора на изотропном конусе K_4 для K^α и N^α.

14.4 О структурных константах алгебры октонионов. Спинорный формализм для n = 8

В этом параграфе рассказывается о том, как по связующим операторам, удовлетворяющим уравнению Клиффорда, построить структурные константы алгебры октонионов. Такие алгебры являются частным случаем нормируемой алгебры с делением (n=8). Рассматривается аксиоматика таких алгебр. Вывод всех результатов этого параграфа сделан на основе [4].

Пусть операторы $\eta_\alpha{}^{ab}$, $(\alpha\ ,\beta\ ,... = \overline{1,6}; a\ ,b\ ,... = \overline{1,4})$ - это кососимметричные операторы, построенные выше для спинорного формализма при n=6. Тогда операторы $\eta_\Lambda{}^{AB}$ $(\Lambda\ ,\Psi\ ,... = \overline{1,8}; A\ ,B\ ,... = \overline{1,8})$ для спинорного формализма при n=8 строятся согласно схеме (8.20). Поэтому тензор (8.17) будет обладать симметриями

$$\varepsilon_{AB(CD)} = \frac{1}{2}\varepsilon_{AB}\varepsilon_{CD}, \quad \varepsilon_{A(B|C|D)} = \frac{1}{2}\varepsilon_{AC}\varepsilon_{BD}, \tag{14.84}$$

где согласно (8.19) $\varepsilon_{AC} = \tilde{\varepsilon}_{AC}$ является метрическим на спинорном пространстве $\mathbb{C}^8$.

$$\eta_{\Lambda AB} = \frac{1}{4}\eta_\Lambda{}^{CD}\varepsilon_{ABCD} = \eta_\Lambda{}^{CD}\varepsilon_{AD}\varepsilon_{BC}, \quad \eta_\Lambda := \frac{1}{4}\eta_\Lambda{}^{CD}\varepsilon_{CD}. \tag{14.85}$$

Вторoе тождество (14.84) очень важно. Оно обеспечивает невырожденность оператора $\varepsilon_{XA}{}^{YB}X^XX_Y$ и, как следствие, нормируемость алгебры октав. При этом тождество (9.11) распадается на три: два тождества альтернативности (левое и правое) и центральное тождество эластичности. Как это происходит.

Теорема 14.2. *Для любого* $r^\Lambda \in \mathbb{C}^8$ *имеет место разложение*

$$r^\Lambda = \eta^\Lambda{}_{AB}X^AY^B \tag{14.86}$$

для некоторых X^A, $\ Y^B \in \mathbb{C}^8$.

Доказательство. Из уравнения Клиффорда (8.1) будет следовать, что

$$R^{AB}R_{CB} = \frac{1}{2}r^\Lambda r_\Lambda\delta_C{}^A, \quad R_{KL}\delta_A{}^B = R_{AR}\varepsilon_{KL}{}^{BR} + R^{BR}\varepsilon_{KLAR}. \tag{14.87}$$

Свернем это уравнение с P^KP^A, выбирая P^K так, что $p := \frac{1}{2}\varepsilon_{AK}P^KP^A \neq 0$, и полагая, что $Q_L := R_{KL}P^K$,

$$Q_LP^B = Q_RP^K\varepsilon_{KL}{}^{BR} + pR^B{}_L. \tag{14.88}$$

Тогда из (14.84) будет следовать

$$R^{BL} = \frac{1}{p}Q^RP^K(\delta_R{}^L\delta_K{}^B - \varepsilon_K{}^{LB}{}_R) = \frac{1}{p}Q^RP^K\varepsilon^{BL}{}_{KR}. \tag{14.89}$$

Остается положить

$$X^A := \frac{1}{\sqrt{p}} P^A, \quad Y^B := \frac{1}{\sqrt{p}} Q^B, \tag{14.90}$$

что и докажет теорему. □

Определим

$$P^{\Lambda}{}_A := \eta^{\Lambda}{}_{AB} X^A, \quad X^A X_A := 2, \tag{14.91}$$

тогда

$$P^{\Lambda}{}_A P^{\Psi}{}_B \varepsilon^{AB} = g^{\Lambda\Psi}, \quad P^{\Lambda}{}_A P^{\Psi}{}_B g_{\Lambda\Psi} = \varepsilon_{AB}. \tag{14.92}$$

Таким образом оператор $P^{\Lambda}{}_A$ определяет изоморфизм между пространствами $\mathbb{C}^n \cong \mathbb{C}^N$ ($n = N = 8$)

$$r^{\Lambda} = P^{\Lambda}{}_B Y^B. \tag{14.93}$$

Положим

$$\eta_{\Lambda}{}^{\Psi\Theta} := \sqrt{2} \eta_{\Lambda}{}^{AB} P^{\Psi}{}_A P^{\Theta}{}_B, \tag{14.94}$$

тогда для таких структурных констант выполнены тождества альтернативности алгебры октав

$$\eta_{\Lambda\Theta}{}^{\Phi} \eta_{(\Psi\Upsilon)}{}^{\Theta} = \eta_{\Lambda(\Psi}{}^{\Theta} \eta_{|\Theta|\Upsilon)}{}^{\Phi}, \quad \eta_{(\Lambda\Psi)}{}^{\Theta} \eta_{\Theta\Phi}{}^{\Upsilon} = \eta_{(\Lambda|\Theta|}{}^{\Upsilon} \eta_{\Psi)\Phi}{}^{\Theta} \tag{14.95}$$

и центральное тождество Муфанг

$$r^{\Phi} \eta_{\Phi\Theta}{}^{\Omega} \eta_{\Lambda\Psi}{}^{\Theta} \eta_{\Omega\Upsilon}{}^{\Gamma} r^{\Upsilon} = r^{\Phi} \eta_{\Phi\Lambda}{}^{\Theta} \eta_{\Theta\Omega}{}^{\Gamma} \eta_{\Psi\Upsilon}{}^{\Omega} r^{\Upsilon}. \tag{14.96}$$

Доказательство в приложении. Теперь, для того, чтобы перейти к структурным константам алгебры октав, необходимо воспользоваться оператором вложения $H_i{}^{\Lambda} : \mathbb{R}^6 \subset \mathbb{C}^6$. Нормируемость алгебры октав будет следовать из уравнения Клиффорда (8.1), выписанного для структурных констант

$$(\eta_{\Phi\Theta}{}^{\Omega} \eta_{\Psi\Xi}{}^{\Delta} + \eta_{\Psi\Theta}{}^{\Omega} \eta_{\Phi\Xi}{}^{\Delta}) g_{\Omega\Delta} = 2 g_{\Phi\Psi} g_{\Theta\Xi}. \tag{14.97}$$

Операторы же вида (12.5)

$$A_{\Theta\Phi L}{}^{X} := \tfrac{1}{2} \eta_{[\Theta}{}^{MX} \eta_{\Phi]ML}, \quad \tilde{A}_{\Theta\Phi L}{}^{X} := \tfrac{1}{4} (\eta_{[\Theta}{}^{MX} \eta_{\Phi]ML} + \eta_{[\Theta}{}^{XM} \eta_{\Phi]LM}) \tag{14.98}$$

будут удовлетворять тождествам

$$\begin{aligned} A_{\Lambda\Psi}{}^{AB} A^{\Lambda\Psi}{}_{CD} &= \delta_{[C}{}^{A} \delta_{D]}{}^{B}, & \tilde{A}_{\Lambda\Psi}{}^{AB} \tilde{A}^{\Lambda\Psi}{}_{CD} &= \tfrac{1}{2} \delta_{[C}{}^{A} \delta_{D]}{}^{B} + \tfrac{1}{4} \varepsilon_{[C}{}^{A}{}_{D]}{}^{B}, \\ A_{\Lambda\Psi}{}^{AB} A^{\Upsilon\Omega}{}_{AB} &= \delta_{[\Lambda}{}^{\Upsilon} \delta_{\Psi]}{}^{\Omega}, & \tilde{A}_{\Lambda\Psi}{}^{AB} \tilde{A}^{\Upsilon\Omega}{}_{AB} &= \delta_{[\Lambda}{}^{\Upsilon} \delta_{\Psi]}{}^{\Omega} + \eta^{[\Upsilon} \delta_{[\Lambda}{}^{\Omega]} \eta_{\Psi]}. \end{aligned} \tag{14.99}$$

Доказательство в приложении.

14.5 Геометрия индуктивного перехода от n=6 к n=8

В этом параграфе рассказывается о том, как построить две квадрики, связанные принципом тройственности Картана. Построение проводится в явном виде, обосновывая стандартную схему индуктивного перехода от n=6 к n=8. Вывод всех результатов этого параграфа сделан на основе [6].

14.5.1 Нуль-пары Розенфельда

В этом параграфе рассказывается о том, как построить явное решение твисторного уравнения при n=6. Вывод всех результатов этого параграфа сделан на основе [6], [19].

Обозначим через $\mathbb{A}^{\mathbb{C}*}$ пространство, двойственное к спинорному 4-мерному комплексному векторному пространству $\mathbb{A}^{\mathbb{C}}$, и образуем 8-мерное комплексное пространство $\mathbb{T}^2$ как прямую сумму $\mathbb{A}^{\mathbb{C}} \oplus \mathbb{A}^{\mathbb{C}*}$. То есть, если X^a $(a, b, ... = \overline{1,4})$ - координаты вектора в $A^{\mathbb{C}}$, а Y_b - координаты ковектора в $A^{\mathbb{C}*}$, то $X^A := (X^a, Y_b), \quad (A, B, ... = \overline{1,8})$ будут координатами вектора из $\mathbb{T}^2$. Будем рассматривать координаты бивектора r^{ab} из (13.18)

$$X^a = ir^{ab}Y_b \tag{14.100}$$

как координаты точки комплексного аффинного пространства $\mathbb{C}\mathbb{A}_6$. Это есть система из 4 линейных уравнений с 6 неизвестными. Для выяснения ее ранга рассмотрим однородное уравнение $r^{ab}Y_b = 0$, которое имеет ненулевые решения тогда и только тогда, когда бивектор простой: $r_{kc}r^{ac} = \frac{1}{4}r^{ab}r_{ab}\delta_k{}^d = 0$ - и, следовательно, представим в виде

$$r^{ab}_{\text{однородное}} = P^aQ^b - P^bQ^a, \tag{14.101}$$

причем P^a и Q^a определены с точностью до их линейных комбинаций. Из этого следует, что $P^aY_a = 0, \quad Q^aY_a = 0$. Обозначим через X^a,S^a,Z^a все решения уравнения $X^aY_a = 0$, которые образуют базис. Тогда наше решение примет вид

$$r^{ab}_{\text{однородное}} = \lambda_1 S^{[a}X^{b]} + \lambda_2 X^{[a}Z^{b]} + \lambda_3 S^{[a}Z^{b]} \tag{14.102}$$

и, следовательно, определит в пространстве бивекторов 3-мерное подпространство. Отсюда получается общее решение

$$r^{ab} = r^{ab}_{\text{частное}} + \lambda_1 S^{[a}X^{b]} + \lambda_2 X^{[a}Z^{b]} + \lambda_3 S^{[a}Z^{b]}, \tag{14.103}$$

где $r^{ab}_{\text{частное}}$-произвольный бивектор, являющийся частным решением.

14.5.2 Построение квадрик $\mathbb{C}Q_6$ и $\mathbb{C}\tilde{Q}_6$

> *В этом параграфе рассказывается о том, как построить квадрики, удовлетворяющие принципу тройственности Картана. Вывод всех результатов этого параграфа сделан на основе [6].*

Покажем, что определенное ранее пространство $\mathbb{T}^2$ будет комплексным пространством, в котором скалярный квадрат вектора определится квадратичной формой

$$\varepsilon_{AB}X^AX^B = 2X^aY_a \tag{14.104}$$

так, что матрица тензора ε_{AB} имеет вид

$$\parallel \varepsilon_{AB} \parallel = \begin{pmatrix} 0 & \delta_a{}^c \\ \delta^b{}_d & 0 \end{pmatrix}. \tag{14.105}$$

При фиксированном r^{ab} уравнение (14.100) определит в $\mathbb{T}^2$ 4-мерное пространство, которое будет являться 4-мерной плоской образующей конуса $\varepsilon_{AB}X^AX^B = 0$. Таким образом, мы можем рассматривать квадрику $\mathbb{C}Q_6$, задаваемую уравнением (14.104), в проективном пространстве $\mathbb{CP}_7$. Ее 4 базисные точки будут удовлетворять условию $\varepsilon_{AB}X^A_IX^B_J = 0, \quad (I, J, ... = \overline{1,4})$. Положим

$$X^A_1 := (X^a, Y_b), \quad X^A_2 := (Z^a, T_b), \quad X^A_3 := (L^a, N_b), \quad X^A_4 := (K^a, M_b). \tag{14.106}$$

Исходя из общего решения уравнения (14.100), каждой точке квадрики $\mathbb{C}Q_6$ можно поставить в соответствие 3-мерную изотропную плоскость пространства $\mathbb{CA}_6$. Точку (t,v,w,x,y,z) пространства $\mathbb{CA}_6$ можно представить прямой $(\lambda T, \lambda V, \lambda U, \lambda S, \lambda W, \lambda X, \lambda Y, \lambda Z)$ пространства $\mathbb{C}^8$, обладающего метрикой

$$dL^2 = dT^2 + dV^2 + dU^2 + dS^2 + dW^2 + dX^2 + dY^2 + dZ^2. \tag{14.107}$$

Эти прямые будут образующими изотропного конуса $\mathbb{C}K_8$: $T^2 + V^2 + U^2 + S^2 + W^2 + X^2 + Y^2 + Z^2 = 0$. Пересечение 7-плоскости $U - iS = 1$ с указанным конусом $\mathbb{C}K_8$ обладает индуцированной метрикой

$$d\tilde{L}^2 = dT^2 + dV^2 + dW^2 + dX^2 + dY^2 + dZ^2. \tag{14.108}$$

Это пространство имеет вид параболоида на $\mathbb{C}K_8$ и тождественно пространству $\mathbb{CR}^6$: $U = 1 + iS = \frac{1}{2}(1 - T^2 - V^2 - W^2 - X^2 - Y^2 - Z^2)$ Всякая образующая этого конуса (множество точек, принадлежащих $\mathbb{C}K_8$ с постоянным отношением T:V:U:S:W:X:Y:Z), не лежащая на гиперплоскости $U = iS$, пересекает параболоид в единственной точке. Образующим конуса, лежащим на гиперплоскости $U = iS$, соответствуют точки, принадлежащие бесконечности пространства $\mathbb{C}^6$. Таким образом, прямым $\mathbb{C}^8$, проходящим через начало $\mathbb{C}^8$, соответствуют точки проективного пространства $\mathbb{CP}_7$. Стереографическая проекция указанного сечения на плоскость (U=1, S=0) с полюсом $N(0,0,\frac{1}{2},\frac{i}{2},0,0,0,0)$ отображает точку P(T,V,U,S,W,X,Y,Z) гиперболоида на точку p(t,v,1,0,w,x,y,z) плоскости (U=1, S=0)

$$\begin{aligned} &\lambda T = t, \ \ \lambda V = v, \ \ \lambda W = w, \ \ \lambda X = x, \ \ \lambda Y = y, \ \ \lambda Z = z, \ \ \lambda = \tfrac{1}{U+iS}, \\ &r^\alpha r_\alpha = -\tfrac{U-iS}{U+iS}, \ \ \lambda U = \tfrac{1}{2}(1 - t^2 - v^2 - w^2 - x^2 - y^2 - z^2) = -\lambda iS + 1. \end{aligned} \tag{14.109}$$

Образующим же конуса $\mathbb{C}K_8$ соответствует квадрика $\mathbb{C}\tilde{Q}_6$ в проективном пространстве $\mathbb{CP}_7$

$$g_{\Lambda\Psi} r^\Lambda r^\Psi = 0. \tag{14.110}$$

14.5.3 Соответствие $\mathbb{C}Q_6 \longmapsto \mathbb{C}\tilde{Q}_6$

> *В этом параграфе рассказывается о том, как построить соответствие между образующими квадрики различного ранга согласно принципу тройственности Картана. Вывод всех результатов этого параграфа сделана основе [6].*

1. Общее решение уравнения (14.100)

   $$r^{ab} = r^{ab}_{\text{частное}} + r^{ab}_{\text{однородное}} = r^{ab}_{\text{частное}} + \lambda_1 S^{[a}X^{b]} + \lambda_2 X^{[a}Z^{b]} + \lambda_3 S^{[a}Z^{b]} \tag{14.111}$$

   определит 4-мерную плоскую образующую конуса $\mathbb{C}K_8$. Тогда для такой образующей определится система

   $$\left\{ \begin{array}{rcl} ir^{ab}Y_b &=& X^a, \\ ir^{ab}T_b &=& Z^a, \\ ir^{ab}N_b &=& L^a, \\ ir^{ab}M_b &=& K^a \end{array} \right. \tag{14.112}$$

   с условиями

   $$X^aY_a = 0, \ \ Z^aT_a = 0, \ \ L^aN_a = 0, \ \ K^aM_a = 0,$$

   $$X^aT_a = -Z^aY_a, \ \ X^aN_a = -L^aY_a, \ \ X^aM_a = -K^aY_a, \tag{14.113}$$

   $$Z^aN_a = -L^aT_a, \ \ Z^aM_a = -K^aT_a, \ \ K^aN_a = -L^aM_a.$$

   Таким образом из 16 уравнений с 6 неизвестными r^{ab} существенными будут только 6 уравнений (10 условий связи), что определит точку $\mathbb{CA}_6$, а значит и точку квадрики $\mathbb{C}\tilde{Q}_6$.

2. Если же нам известно только одно уравнение
$$ir^{ab}Y_b = X^a \tag{14.114}$$
с условием
$$X^aY_a = 0, \tag{14.115}$$
то из 4 уравнений существенными будут лишь 3 (одно условие связи). Это означает, что точке квадрики $\mathbb{C}Q_6$ будет соответствовать плоская 3-мерная образующая $\mathbb{CP}_3$, принадлежащая квадрике $\mathbb{C}\tilde{Q}_6$.

3. Если нам известны два уравнения
$$\begin{cases} ir^{ab}Y_b & = & X^a, \\ ir^{ab}T_b & = & Z^a \end{cases} \tag{14.116}$$
с условиями
$$X^aY_a = 0, \quad Z^aT_a = 0, \quad X^aT_a = -Z^aY_a, \tag{14.117}$$
то из 8 уравнений существенными будут лишь 5 (неизвестных же 6 и 3 условия связи). Это означает, что прямолинейной образующей $\mathbb{CP}_1$ квадрики $\mathbb{C}Q_6$ будет соответствовать прямолинейная образующая $\mathbb{CP}_1$, принадлежащая квадрике $\mathbb{C}\tilde{Q}_6$. При этом многообразие образующих $\mathbb{CP}_1(\mathbb{C}Q_6)$, принадлежащих одной и той же образующей $\mathbb{CP}_3(\mathbb{C}Q_6)$, определит пучок образующих $\mathbb{CP}_1(\mathbb{C}\tilde{Q}_6)$, принадлежащий квадрике $\mathbb{C}\tilde{Q}_6$ (этот пучок является на самом деле конусом). Центр пучка определится системой (14.112).

4. Если нам известны три уравнения
$$\begin{cases} ir^{ab}Y_b & = & X^a, \\ ir^{ab}T_b & = & Z^a, \\ ir^{ab}N_b & = & L^a, \end{cases} \tag{14.118}$$
с условиями
$$\begin{gathered} X^aY_a = 0, \quad Z^aT_a = 0, \quad L^aN_a = 0, \\ X^aT_a = -Z^aY_a, \quad X^aN_a = -L^aY_a, \quad Z^aN_a = -L^aT_a, \end{gathered} \tag{14.119}$$
то из 12 уравнений существенными будут лишь 6 (и неизвестных 6 с 6 условиями связи). Это означает, что 2-мерной образующей $\mathbb{CP}_2$ квадрики $\mathbb{C}Q_6$ будет соответствовать точка квадрики $\mathbb{C}\tilde{Q}_6$. При этом многообразие образующих $\mathbb{CP}_2(\mathbb{C}Q_6)$, принадлежащих одной и той же образующей $\mathbb{C}P_3(\mathbb{C}Q_6)$, определит единственную точку квадрики $\mathbb{C}\tilde{Q}_6$. Эта точка определится системой (14.112).



Рис. 1: Соответствие $\forall CP_2 \subset CP_3 \leftrightarrow R$



Рис. 2: Соответствие $CP_3 \supset CP_1 \leftrightarrow CP_1 \subset K_6$

14.5.4 Геометрия перехода к связующим операторам ${\eta_\Lambda}^{KL}$ для n=8

В этом параграфе рассказывается о том, как построить связующие операторы для спинорного формализма при n=8. Вывод всех результатов этого параграфа сделана основе [6].

Рассмотрим далее бивектор $\hat{R}^{AB}$ такой, что его составляющие векторы ${X_1}^A, {X_2}^A$ определены как

$$X_1^A := (X^a, Y_b), \quad X_2^A := (Z^a, T_b), \quad \left\{ \begin{array}{rcl} ir^{ab}Y_b & = & X^a, \\ ir^{ab}T_b & = & Z^a. \end{array} \right. \tag{14.120}$$

Этим определится цепочка тождеств

$$\begin{gathered} ir^{ab}Y_b = X^a, \quad ir_{ab}Z^b = pf(r)T_a, \\[2em] \tfrac{1}{2} ir_{cd}\varepsilon^{abcd}Y_b\varepsilon_{aklm} = X^a\varepsilon_{aklm}, \\[2em] 3ir_{[kl}Y_{m]} = X^a\varepsilon_{aklm}, \\[2em] ir_{kl}Y_m + 2ir_{m[k}Y_{l]} = X^a\varepsilon_{aklm}. \end{gathered} \tag{14.121}$$

Свернем последнее тождество с Z^m, что даст

$$\begin{gathered} ir_{kl}Y_mZ^m = X^mZ^n\varepsilon_{klmn} + pf(r)\cdot 2T_{[k}Y_{l]}, \\ X^mZ^n - X^nZ^m = -ir^{mn}T_kX^k + \varepsilon^{klmn}pf(r)Y_{[k}T_{l]}. \end{gathered} \tag{14.122}$$

Исходя из вышесказанного, рассмотрим прямолинейную образующую квадрики $\mathbb{C}Q_6$, определенную бивектором $(T_kX^k \neq 0)$

$$\begin{gathered} \hat{R}^{AB} = 2X_1^{[A}X_2^{B]} = \begin{pmatrix} X^aZ^b - X^bZ^a & X^aT_d - Y_dZ^a \\ Y_cZ^b - T_cX^b & Y_cT_d - Y_dT_c \end{pmatrix} = \\[2em] = \begin{pmatrix} -ir^{ab}T_kX^k + \varepsilon^{klab}pf(r)Y_kT_l & 2ir^{ar}Y_{[r}T_{b]} \\ i\varepsilon^{kbmn}r_{ck}Y_{[m}T_{n]} - {\delta_c}^bX^kT_k & 2Y_{[c}T_{d]} \end{pmatrix} = \\[2em] = \underbrace{T_lX^l \begin{pmatrix} -\frac{1}{2}i{\delta^a}_kr^\gamma r_\gamma & r^{ar} \\ r_{ck} & -i{\delta_c}^r \end{pmatrix}}_{:={R^A}_K} \cdot \underbrace{\frac{1}{T_lX^l} \begin{pmatrix} \varepsilon^{kbmn}iY_mT_n & 0 \\ -i{\delta_r}^bX^nT_n & 2iY_{[r}T_{d]} \end{pmatrix}}_{:=P^{KB}} := {R^A}_KP^{KB}, \end{gathered} \tag{14.123}$$

$$
\begin{aligned}
R^{AB} &:= \varepsilon^{BC} R^{A}{}_{C} = T_l X^l \begin{pmatrix} r^{an} & -\frac{1}{2} i \delta^{a}{}_{k} r^{\gamma} r_{\gamma} \\ -i \delta_{c}{}^{n} & r_{ck} \end{pmatrix}, \\
R_{AB} &:= \varepsilon_{AC} R^{C}{}_{B} = T_l X^l \begin{pmatrix} r_{ck} & -i \delta_{c}{}^{n} \\ -\frac{1}{2} i \delta^{a}{}_{k} r^{\gamma} r_{\gamma} & r^{an} \end{pmatrix},
\end{aligned}
\tag{14.124}
$$

где ε_{AC} все тот же метрический симметрический спин-тензор. При этом будет верно уравнение

$$
R_{A}{}^{C} \hat{R}^{AB} = 0, \tag{14.125}
$$

которое означает, что любой тензор $\hat{R}^{AB}$, представляющий образующую $\mathbb{CP}_1(\mathbb{C}Q_6)$, будет содержать один и тот же тензор $R^{A}{}_{K}$ в своем разложении, при этом второй тензор разложения P^{KB} будет отвечать за положение $\mathbb{CP}_1$ в $\mathbb{CP}_3$. Поэтому есть резон поставить в соответствие точке квадрики $\mathbb{C}\tilde{Q}_6$ биспинор R^{AB}, которым она определится однозначно. При переходе к пространству $\mathbb{CR}^8$, исходя из

$$
\begin{array}{lllllllll}
r^{12} & = & \frac{1}{\sqrt{2}}(v+iw), & r^{13} & = & \frac{1}{\sqrt{2}}(x+iy), & r^{14} & = & \frac{i}{\sqrt{2}}(t+iz), \\
r^{23} & = & \frac{i}{\sqrt{2}}(iz-t), & r^{24} & = & \frac{1}{\sqrt{2}}(-x+iy), & r^{34} & = & \frac{1}{\sqrt{2}}(v-iw),
\end{array}
\tag{14.126}
$$

определим однородные координаты $\mathbb{CR}^8$ следующим образом

$$
\lambda = \left\{
\begin{array}{ccccccccccccccc}
R^{12} & : & R^{13} & : & R^{14} & : & R^{23} & : & R^{24} & : & R^{34} & : & R^{15} & : & R^{51}, \\
\\
r^{12} & : & r^{13} & : & r^{14} & : & r^{23} & : & r^{24} & : & r^{34} & : & -\frac{1}{2} i r^{\gamma} r_{\gamma} & : & -i,
\end{array}
\right.
\tag{14.127}
$$

$$
\begin{array}{lllllllll}
R^{12} & = & \frac{1}{\sqrt{2}}(V+iW), & R^{13} & = & \frac{1}{\sqrt{2}}(iY+X), & R^{14} & = & \frac{1}{\sqrt{2}}(-Z+iT), \\
R^{23} & = & \frac{1}{\sqrt{2}}(-Z-iT), & R^{24} & = & \frac{1}{\sqrt{2}}(-X+iY), & R^{34} & = & \frac{1}{\sqrt{2}}(V-iW), \\
R^{15} & = & \frac{1}{2}(iU+S), & R^{51} & = & -iU+S, & & &
\end{array}
\tag{14.128}
$$

$$
\begin{array}{lllllll}
R^{12} & = & -R^{21} & = & R^{78} & = & -R^{87}, \\
R^{13} & = & -R^{31} & = & R^{86} & = & -R^{68}, \\
R^{14} & = & -R^{41} & = & R^{67} & = & -R^{76}, \\
R^{23} & = & -R^{32} & = & R^{58} & = & -R^{85}, \\
R^{24} & = & -R^{42} & = & R^{75} & = & -R^{57}, \\
R^{34} & = & -R^{43} & = & R^{56} & = & -R^{65}, \\
R^{15} & = & R^{26} & = & R^{37} & = & R^{48}, \\
R^{51} & = & R^{62} & = & R^{73} & = & R^{84}.
\end{array}
\tag{14.129}
$$

$$R^{AB}R_{AB} = 8(R^{12}R^{34} - R^{13}R^{24} + R^{14}R^{23} + R^{15}R^{51}) =$$

$$= 4(T^2 + V^2 + U^2 + S^2 + W^2 + X^2 + Y^2 + Z^2) := 4nf(R), \tag{14.130}$$

$$R^{AB}R_{CB} = \tfrac{1}{2}nf(R)\delta_C{}^A \quad \Rightarrow \quad R^{AB}R_{AB} = 4nf(R).$$

Для того, чтобы $(r_I)^\Lambda$ определяли образующую квадрики $\mathbb{C}\tilde{Q}_6$ необходимо и достаточно выполнение условия

$$g_{\Lambda\Psi}(r_I)^\Lambda(r_J)^\Psi = 0. \tag{14.131}$$

Определим некоторые связующие операторы $\eta_\Lambda{}^{BC}$ так, чтобы выполнялись условия

$$r_\Lambda = \frac{1}{4}\eta_\Lambda{}^{BC}R_{BC}, \quad r^\Lambda = \frac{1}{4}\eta^\Lambda{}_{BC}R^{BC}. \tag{14.132}$$

Тогда эти операторы удовлетворяют уравнению Клиффорда

$$g_{\Lambda\Psi}\delta_K{}^L = \eta_{\Lambda K}{}^R\eta_\Psi{}^L{}_R + \eta_{\Lambda K}{}^R\eta_\Psi{}^L{}_R. \tag{14.133}$$

Определим

$$\varepsilon_{PQRT} := \eta^\Lambda{}_{PQ}\eta^\Psi{}_{RT}g_{\Lambda\Psi}, \quad \varepsilon_{PQRT} = \varepsilon_{RTPQ}, \tag{14.134}$$

что даст еще один метрический спинор ε_{PQRT}, с помощью которого можно поднимать и опускать парные индексы

$$\begin{gathered}
g_{\Lambda\Psi} = \tfrac{1}{4}\eta_\Lambda{}^{PQ}\eta_{\Psi PQ}, \quad \eta_{\Lambda RT} = \tfrac{1}{4}\eta_\Lambda{}^{PQ}\varepsilon_{PQRT}, \quad \varepsilon_{PQ} = \tfrac{1}{4}\varepsilon_{PQRT}\varepsilon^{RT},\\
\varepsilon_{STPQ}\delta_K{}^L = \varepsilon_{STK}{}^R\varepsilon_{PQ}{}^L{}_R + \varepsilon_{PQK}{}^R\varepsilon_{ST}{}^L{}_R, \quad \varepsilon_{STPQ} = \tfrac{1}{4}\varepsilon_{ST}{}^{KR}\varepsilon_{PQKR},\\
\varepsilon_{PQ(KL)} = \tfrac{1}{2}\varepsilon_{PQ}\varepsilon_{KL}, \quad \varepsilon_{[PQ](KL)} = 0, \quad \eta_A{}^{(MN)} = \tfrac{1}{8}\eta_A{}^{KL}\varepsilon_{KL}\varepsilon^{MN},
\end{gathered} \tag{14.135}$$

при этом результат применения двух метрических тензоров должен быть одинаков. Если же матрицы тензоров $g_{\Lambda\Psi}$ и ε_{KL} имеют вид

$$\| g_{\Lambda\Psi} \| = \begin{pmatrix} 1&0&0&0&0&0&0&0\\ 0&1&0&0&0&0&0&0\\ 0&0&1&0&0&0&0&0\\ 0&0&0&1&0&0&0&0\\ 0&0&0&0&1&0&0&0\\ 0&0&0&0&0&1&0&0\\ 0&0&0&0&0&0&1&0\\ 0&0&0&0&0&0&0&1 \end{pmatrix}, \quad \| \varepsilon_{KL} \| = \begin{pmatrix} 0&0&0&0&1&0&0&0\\ 0&0&0&0&0&1&0&0\\ 0&0&0&0&0&0&1&0\\ 0&0&0&0&0&0&0&1\\ 1&0&0&0&0&0&0&0\\ 0&1&0&0&0&0&0&0\\ 0&0&1&0&0&0&0&0\\ 0&0&0&1&0&0&0&0 \end{pmatrix}, \tag{14.136}$$

то существенные координаты операторов $\eta^\Lambda{}_{KL}$ в некотором базисе будут такими

$$\begin{array}{llll}
\eta^2{}_{12} = \frac{1}{\sqrt{2}}, & \eta^2{}_{34} = \frac{1}{\sqrt{2}}, & \eta^5{}_{12} = -\frac{i}{\sqrt{2}}, & \eta^5{}_{34} = \frac{i}{\sqrt{2}}, \\
\eta^2{}_{78} = \frac{1}{\sqrt{2}}, & \eta^2{}_{56} = \frac{1}{\sqrt{2}}, & \eta^5{}_{78} = -\frac{i}{\sqrt{2}}, & \eta^5{}_{56} = \frac{i}{\sqrt{2}}, \\
\eta^1{}_{14} = -\frac{i}{\sqrt{2}}, & \eta^1{}_{23} = \frac{i}{\sqrt{2}}, & \eta^8{}_{14} = -\frac{1}{\sqrt{2}}, & \eta^8{}_{23} = -\frac{1}{\sqrt{2}}, \\
\eta^1{}_{67} = -\frac{i}{\sqrt{2}}, & \eta^1{}_{58} = \frac{i}{\sqrt{2}}, & \eta^8{}_{67} = -\frac{1}{\sqrt{2}}, & \eta^8{}_{58} = -\frac{1}{\sqrt{2}}, \\
\eta^7{}_{13} = -\frac{i}{\sqrt{2}}, & \eta^7{}_{24} = -\frac{i}{\sqrt{2}}, & \eta^6{}_{13} = \frac{1}{\sqrt{2}}, & \eta^6{}_{24} = -\frac{1}{\sqrt{2}}, \\
\eta^7{}_{68} = \frac{i}{\sqrt{2}}, & \eta^7{}_{57} = \frac{i}{\sqrt{2}}, & \eta^6{}_{68} = -\frac{1}{\sqrt{2}}, & \eta^6{}_{57} = \frac{1}{\sqrt{2}}, \\
\eta^4{}_{15} = \frac{1}{\sqrt{2}}, & \eta^4{}_{51} = \frac{1}{\sqrt{2}}, & \eta^3{}_{15} = -\frac{i}{\sqrt{2}}, & \eta^3{}_{51} = \frac{i}{\sqrt{2}}, \\
\eta^4{}_{26} = \frac{1}{\sqrt{2}}, & \eta^4{}_{62} = \frac{1}{\sqrt{2}}, & \eta^3{}_{26} = -\frac{i}{\sqrt{2}}, & \eta^3{}_{62} = \frac{i}{\sqrt{2}}, \\
\eta^4{}_{37} = \frac{1}{\sqrt{2}}, & \eta^4{}_{73} = \frac{1}{\sqrt{2}}, & \eta^3{}_{37} = -\frac{i}{\sqrt{2}}, & \eta^3{}_{73} = \frac{i}{\sqrt{2}}, \\
\eta^4{}_{48} = \frac{1}{\sqrt{2}}, & \eta^4{}_{84} = \frac{1}{\sqrt{2}}, & \eta^3{}_{48} = -\frac{i}{\sqrt{2}}, & \eta^3{}_{84} = \frac{i}{\sqrt{2}},
\end{array} \tag{14.137}$$

а существенные координаты операторов $\eta_\Lambda{}^{KL}$ такими

$$\begin{array}{llll}
\eta_2{}^{12} = \frac{1}{\sqrt{2}}, & \eta_2{}^{34} = \frac{1}{\sqrt{2}}, & \eta_5{}^{12} = \frac{i}{\sqrt{2}}, & \eta_5{}^{34} = -\frac{i}{\sqrt{2}}, \\
\eta_2{}^{78} = \frac{1}{\sqrt{2}}, & \eta_2{}^{56} = \frac{1}{\sqrt{2}}, & \eta_5{}^{78} = \frac{i}{\sqrt{2}}, & \eta_5{}^{56} = -\frac{i}{\sqrt{2}}, \\
\eta_1{}^{14} = \frac{i}{\sqrt{2}}, & \eta_1{}^{23} = -\frac{i}{\sqrt{2}}, & \eta_8{}^{14} = -\frac{1}{\sqrt{2}}, & \eta_8{}^{23} = -\frac{1}{\sqrt{2}}, \\
\eta_1{}^{67} = \frac{i}{\sqrt{2}}, & \eta_1{}^{58} = -\frac{i}{\sqrt{2}}, & \eta_8{}^{67} = -\frac{1}{\sqrt{2}}, & \eta_8{}^{58} = -\frac{1}{\sqrt{2}}, \\
\eta_7{}^{13} = \frac{i}{\sqrt{2}}, & \eta_7{}^{24} = \frac{i}{\sqrt{2}}, & \eta_6{}^{13} = \frac{1}{\sqrt{2}}, & \eta_6{}^{24} = -\frac{1}{\sqrt{2}}, \\
\eta_7{}^{68} = -\frac{i}{\sqrt{2}}, & \eta_7{}^{57} = -\frac{i}{\sqrt{2}}, & \eta_6{}^{68} = -\frac{1}{\sqrt{2}}, & \eta_6{}^{57} = \frac{1}{\sqrt{2}}, \\
\eta_4{}^{15} = \frac{1}{\sqrt{2}}, & \eta_4{}^{51} = \frac{1}{\sqrt{2}}, & \eta_3{}^{15} = \frac{i}{\sqrt{2}}, & \eta_3{}^{51} = -\frac{i}{\sqrt{2}}, \\
\eta_4{}^{26} = \frac{1}{\sqrt{2}}, & \eta_4{}^{62} = \frac{1}{\sqrt{2}}, & \eta_3{}^{26} = \frac{i}{\sqrt{2}}, & \eta_3{}^{62} = -\frac{i}{\sqrt{2}}, \\
\eta_4{}^{37} = \frac{1}{\sqrt{2}}, & \eta_4{}^{73} = \frac{1}{\sqrt{2}}, & \eta_3{}^{37} = \frac{i}{\sqrt{2}}, & \eta_3{}^{73} = -\frac{i}{\sqrt{2}}, \\
\eta_4{}^{48} = \frac{1}{\sqrt{2}}, & \eta_4{}^{84} = \frac{1}{\sqrt{2}}, & \eta_3{}^{48} = \frac{i}{\sqrt{2}}, & \eta_3{}^{84} = -\frac{i}{\sqrt{2}},
\end{array} \tag{14.138}$$

что несколько отличается от базиса, в котором выше были определены однородные координаты $\mathbb{CR}^8$. Таким образом, биспинор $\hat{R}^{AB}$ определит прямолинейную образующую $\mathbb{CP}_1(\mathbb{C}Q_6)$, принадлежащую некоторой плоской образующей $\mathbb{CP}_3(\mathbb{C}Q_6)$, которая определит некоторую точку с координатами R^{AB} квадрики $\mathbb{C}\tilde{Q}_6$. Определим некоторый тензор $\hat{\hat{R}}^{AB}$

$$\hat{\hat{R}}^{AB} := \hat{R}^{AK}\hat{P}_K{}^B, \quad \hat{P}_K{}^B := \begin{pmatrix} i\delta_m{}^k & 0 \\ 0 & -pf(r)i\delta^n{}_r \end{pmatrix}. \tag{14.139}$$

Тензор $\hat{\hat{R}}^{AB}$ по-прежнему будет представлять прямолинейную образующую $\mathbb{CP}_1(\mathbb{C}Q_6)$, принадлежащую некоторой плоской образующей $\mathbb{CP}_3(\mathbb{C}Q_6)$. Применим операторы $\eta^\Lambda{}_{KL}$ к тензору $\hat{\hat{R}}^{AB}$ и получим

$$\eta^\Lambda{}_{KL}\hat{\hat{R}}^{KL} = \eta^\Lambda{}_{KL}T_mX^m \begin{pmatrix} r^{an} & -\frac{1}{2}i\delta^a{}_k r^\gamma r_\gamma \\ -i\delta_c{}^n & r_{ck} \end{pmatrix} = \eta^\Lambda{}_{KL}R^{KL}. \tag{14.140}$$

Таким образом, операторы $\eta^{\Lambda}{}_{KL}$ осуществляют факторизацию прямолинейных образующих $\mathbb{CP}_1(\mathbb{C}Q_6)$ по принадлежности к одной плоской образующей $\mathbb{CP}_3(\mathbb{C}Q_6)$, и это определяет точку квадрики $\mathbb{C}\tilde{Q}_6$. В однородных же координатах тензор R^{KL} определит координаты точки R пространства $\mathbb{CR}^8$.

Выясним какому семейству принадлежат рассматриваемые выше образующие $\mathbb{CP}_3(\mathbb{C}Q_6)$. Для этого рассмотрим условия

$$\begin{gathered}\varepsilon_{AB}(X_I)^A(X_J)^B = 0,\\ X^{ABCD} := \varepsilon^{IJKL}(X_I)^A(X_J)^B(X_K)^C(X_L)^D,\end{gathered} \tag{14.141}$$

где ε^{IJKL} - квадривектор, кососимметричный по всем индексам. Кроме того рассмотрим 8-вектор $e_{ABCDKLMN}$, тоже кососимметричный по всем индексам. Тогда, если в условии

$$\frac{1}{24}e_{ABCDKLMN}X^{ABCD} = \rho\varepsilon_{KR}\varepsilon_{LT}\varepsilon_{MU}\varepsilon_{NV}X^{RTUV}, \quad \rho^2 = 1 \tag{14.142}$$

$\rho = 1$, то будем говорить, что плоские образующие $\mathbb{CP}_3(\mathbb{C}Q_6)$ принадлежат I семейству, а если $\rho = -1$, то - II семейству. В нашем случае

$$(X_I)^A = ((X_I)^a, (Y_I)_b), \tag{14.143}$$

и тогда

$$\varepsilon^{IJKL}(X_I)^1(X_J)^2(X_K)^3(X_L)^4 = \rho\varepsilon^{IJKL}(X_I)^1(X_J)^2(X_K)^3(X_L)^4. \tag{14.144}$$

Откуда $\rho = 1$. Это означает, что наши образующие необходимо принадлежат I семейству. Кроме того, существует тензор $P_{\Lambda}{}^{B} := \eta_{\Lambda}{}^{AB}X_A$ для $X_A = (1, 0, 0, 0, 1, 0, 0, 0)$

$$\| P_{\Lambda}{}^{B} \| = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 & 0 & 0 & i & 0 & 0 & 0 & -i\\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0\\ -i & 0 & 0 & 0 & i & 0 & 0 & 0\\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0\\ 0 & i & 0 & 0 & 0 & -i & 0 & 0\\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0\\ 0 & 0 & i & 0 & 0 & 0 & -i & 0\\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & -1 \end{pmatrix}, \quad det \| P_{\Lambda}{}^{B} \| = -1, \tag{14.145}$$

$$\varepsilon_{AB}P_{\Lambda}{}^{A}P_{\Psi}{}^{B} = g_{\Lambda\Psi}, \quad (X_I)^A = P_{\Lambda}{}^{A}(r_I)^{\Lambda}.$$

Тогда из

$$\begin{gathered} g_{\Lambda\Psi}(r_I)^{\Lambda}(r_J)^{\Psi}=0, \\ X^{ABCD}={P_{\Lambda}}^{A}{P_{\Psi}}^{B}{P_{\Phi}}^{C}{P_{\Theta}}^{D}r^{\Lambda\Psi\Phi\Theta}, \end{gathered} \tag{14.146}$$

следует, что $\rho=1$, и

$$\tfrac{1}{24}e_{ABCDKLMN}X^{ABCD}=\varepsilon_{KR}\varepsilon_{LT}\varepsilon_{MU}\varepsilon_{NV}X^{RTUV},$$

$$\begin{gathered} \tfrac{1}{24}e_{ABCDKLMN}{P_{\Lambda}}^{A}{P_{\Psi}}^{B}{P_{\Phi}}^{C}{P_{\Theta}}^{D}r^{\Lambda\Psi\Phi\Theta}= \\ =\varepsilon_{KA}\varepsilon_{LB}\varepsilon_{MC}\varepsilon_{ND}{P_{\Lambda_1}}^{A}{P_{\Psi_1}}^{B}{P_{\Phi_1}}^{C}{P_{\Theta_1}}^{D}r^{\Lambda_1\Psi_1\Phi_1\Theta_1}, \end{gathered}$$

$$\begin{gathered} \tfrac{1}{24}e_{ABCDKLMN}{P_{\Lambda}}^{A}{P_{\Psi}}^{B}{P_{\Phi}}^{C}{P_{\Theta}}^{D}{P_{\Lambda_1}}^{K}{P_{\Psi_1}}^{L}{P_{\Phi_1}}^{M}{P_{\Theta_1}}^{N}r^{\Lambda\Psi\Phi\Theta}= \\ =g_{\Lambda\Lambda_1}g_{\Psi\Psi_1}g_{\Phi\Phi_1}g_{\Theta\Theta_1}r^{\Lambda_1\Psi_1\Phi_1\Theta_1}, \end{gathered} \tag{14.147}$$

$$\tfrac{1}{24}det\parallel {P_{\Omega}}^{A}\parallel e_{\Lambda\Psi\Phi\Theta\Lambda_1\Psi_1\Phi_1\Theta_1}r^{\Lambda_1\Psi_1\Phi_1\Theta_1}=g_{\Lambda\Lambda_1}g_{\Psi\Psi_1}g_{\Phi\Phi_1}g_{\Theta\Theta_1}r^{\Lambda_1\Psi_1\Phi_1\Theta_1},$$

$$\tfrac{1}{24}e_{\Lambda\Psi\Phi\Theta\Lambda_1\Psi_1\Phi_1\Theta_1}r^{\Lambda_1\Psi_1\Phi_1\Theta_1}=g_{\Lambda\Lambda_1}g_{\Psi\Psi_1}g_{\Phi\Phi_1}g_{\Theta\Theta_1}r^{\Lambda_1\Psi_1\Phi_1\Theta_1},$$

Отсюда видно, что и образующие $\mathbb{CP}_3(\mathbb{C}\tilde{Q}_6)$ необходимо принадлежат I семейству. Для того, чтобы получить образующие II семейства, необходимо подействовать элементарным ортогональным отражением (с определителем, равным -1) на оператор ${P_{\Lambda}}^{K}$.

14.5.5 Соответствие $\mathbb{C}\tilde{Q}_6\longmapsto\mathbb{C}Q_6$

> *В этом параграфе рассказывается о том, как построить обратное соответствие для квадрик, удовлетворяющие принципу тройственности Картана. Вывод всех результатов этого параграфа сделан на основе [6].*

Применяя операторы ${\eta_{\Lambda}}^{KL}$ к $g_{\Lambda\Psi}(r_I)^{\Lambda}(r_J)^{\Psi}=0$, получим

$$\begin{gathered} (R_I)^{AB}(R_J)_{AB}=0\Leftrightarrow((R_I)^{AB}-(R_J)^{AB})((R_K)_{AB}-(R_L)_{AB})=0\Leftrightarrow \\ ((R_I)^{ab}-(R_J)^{ab})((R_K)_{ab}-(R_L)_{ab})=0. \end{gathered} \tag{14.148}$$

Здесь I,J, как обычно, номера базисных точек. Этим определится система

$$\left\{\begin{array}{lcl} i(R_1)^{ab}Y_b & = & X^a, \\ i(R_2)^{ab}Y_b & = & X^a, \\ i(R_3)^{ab}Y_b & = & X^a, \\ i(R_4)^{ab}Y_b & = & X^a, \end{array}\right. \Leftrightarrow \left\{\begin{array}{lcl} i((R_1)^{ab}-(R_2)^{ab})Y_b & = & 0, \\ i((R_1)^{ab}-(R_3)^{ab})Y_b & = & 0, \\ i((R_3)^{ab}-(R_4)^{ab})Y_b & = & 0, \\ i(R_1)^{ab}Y_b & = & X^a. \end{array}\right. \tag{14.149}$$

Далее будем рассматривать только правую систему. Она строится следующим образом. Всегда существует такой ковектор Y_a, который обнуляет 3 различных простых бивектора. Это утверждение сводится к существованию ковектора, ортогонального данным трем векторам, поскольку каждый из простых бивекторов раскладывается по формуле $R^{ab} = 2P^{[a}Q^{b]}$. По четвертому же уравнению определится некоторый вектор X^a. С другой стороны, на основании того, что все $(R_I)^{ab}$ имеют вид $R^{ab}_{\text{частное}} + \lambda_1 S^{[a}X^{b]} + \lambda_2 X^{[a}Z^{b]} + \lambda_3 S^{[a}Z^{b]}$ при фиксированных $\lambda_1, \lambda_2, \lambda_3$, верно равенство

$$((R_I)^{ab} - (R_J)^{ab})((R_K)_{ab} - (R_L)_{ab}) = 0. \tag{14.150}$$

1. Итак, пусть нам известно последнее уравнение системы (14.149)

$$iR_1{}^{ab}Y_b = X^a, \tag{14.151}$$

то мы имеем 4 уравнения, которые все будут существенными. Поскольку у нас 8 неизвестных (X^a, Y_b) при фиксированных $(R_I)^{ab}$, то точка квадрики $\mathbb{C}\tilde{Q}_6$ определит 3-мерную плоскую образующую $\mathbb{CP}_3(\mathbb{C}Q_6)$.

2. Если нам известны все уравнения системы (14.149) с условиями

$$((R_1)^{ab} - (R_2)^{ab})((R_1)_{ab} - (R_2)_{ab}) = 0, \quad ((R_1)^{ab} - (R_3)^{ab})((R_1)_{ab} - (R_3)_{ab}) = 0,$$

$$((R_3)^{ab} - (R_4)^{ab})((R_3)_{ab} - (R_4)_{ab}) = 0,$$

$$(R_1)^{ab}(R_2)_{ab} = 0, \quad (R_1)^{ab}(R_3)_{ab} = 0, \quad (R_1)^{ab}(R_4)_{ab} = 0,$$

$$(R_2)^{ab}(R_3)_{ab} = 0, \quad (R_2)^{ab}(R_4)_{ab} = 0, \quad (R_3)^{ab}(R_4)_{ab} = 0, \tag{14.152}$$

то из 16 уравнений, существенными будут 7 (8 неизвестных и 9 условий связи). Поэтому образующей $\mathbb{CP}_3(\mathbb{C}\tilde{Q}_6)$ будет соответствовать точка квадрики $\mathbb{C}Q_6$.

3. Если нам известны 3 уравнения системы (14.149)

$$\begin{cases} i((R_1)^{ab} - (R_2)^{ab})Y_b &= 0, \\ i((R_1)^{ab} - (R_3)^{ab})Y_b &= 0, \\ i(R_1)^{ab}Y_b &= X^a \end{cases} \tag{14.153}$$

с условиями

$$((R_1)^{ab} - (R_2)^{ab})((R_1)_{ab} - (R_2)_{ab}) = 0, \quad ((R_1)^{ab} - (R_3)^{ab})((R_1)_{ab} - (R_3)_{ab}) = 0,$$

$$(R_1)^{ab}(R_2)_{ab} = 0, \quad (R_1)^{ab}(R_3)_{ab} = 0, \quad (R_2)^{ab}(R_3)_{ab} = 0, \tag{14.154}$$

то из 12 уравнений, существенными будут тоже 7 уравнений (8 неизвестных и 5 условий связи). Это означает, что образующей $\mathbb{CP}_2(\mathbb{C}\tilde{Q}_6)$ будет соответствовать точка квадрики $\mathbb{C}Q_6$. При этом многообразие образующих $\mathbb{CP}_2(\mathbb{C}\tilde{Q}_6)$, принадлежащих одной образующей $\mathbb{CP}_3(\mathbb{C}\tilde{Q}_6)$, определит единственную точку квадрики $\mathbb{C}Q_6$.

4. Если нам известны только 2 уравнения системы (14.149)

$$\begin{cases} i((R_1)^{ab} - (R_2)^{ab})Y_b &= 0, \\ i(R_1)^{ab}Y_b &= X^a \end{cases} \tag{14.155}$$

с условиями

$$((R_1)^{ab} - (R_2^{ab})((R_1)_{ab} - (R_2)_{ab}) = 0, \quad (R_1)^{ab}(R_2)_{ab} = 0, \tag{14.156}$$

то из 8 уравнений, существенными будет только 6 уравнений (8 неизвестных и 2 условия связи). Поэтому образующей $\mathbb{CP}_1(\mathbb{C}\tilde{Q}_6)$ будет соответствовать прямолинейная образующая $\mathbb{CP}_1(\mathbb{C}Q_6)$. При этом многообразие образующих $\mathbb{CP}_1(\mathbb{C}\tilde{Q}_6)$, принадлежащих одной образующей $\mathbb{CP}_3(\mathbb{C}\tilde{Q}_6)$, определит пучок прямых $\mathbb{CP}_1(\mathbb{C}Q_6)$, принадлежащий квадрике $\mathbb{C}Q_6$. Центр пучка определится системой (14.149).

14.5.6 О майорановских спинорах.

В этом параграфе рассказывается о том, что такое условие майорантности спинора с точки зрения инволюции в спинорном пространстве. Вывод всех результатов этого параграфа сделан на основе [22], [21].

В качестве иллюстрации рассмотрим символы Инфельда - Ван дер Вардена [22, т. 1, с. 161]. Пусть уравнение Клиффорда задано на векторном пространстве $\mathbb{C}^4$ и в некотором специальном базисе имеет вид

$$\eta_\Lambda \sigma_\Psi + \eta_\Psi \sigma_\Lambda = \delta_{\Lambda\Psi}, \quad (\Lambda, \Psi, ... = \overline{0,3}). \tag{14.157}$$

Если задано действительное вложение ${H_i}^\Lambda : \mathbb{R}^4_{(1,3)} \subset \mathbb{C}^4$, то этим определена инволюция ${S_A}^{A'} := \varepsilon^{A'B'} S_{AB'}$ согласно следствию 8.4, случай II).

$${S_\Lambda}^{\Psi'} {\bar{\eta}_{\Psi'}}{}^{A'B'} = {\eta_\Lambda}^{BA} {S_A}^{A'} {S_B}^{B'}, \quad (A, A', ... = \overline{1,2}). \tag{14.158}$$

Это позволит определить символы Инфельда - Ван дер Вардена следующим образом

$$\begin{gathered} {g_i}^{AA'} := {H_i}^\Lambda {\eta_\Lambda}^{AB} {S_B}^{A'}, \\ \overline{{g_i}^{AD'}} = {\bar{g}_i}{}^{A'D} := {\bar{H}_i}{}^{\Lambda'} {\bar{\eta}_{\Lambda'}}{}^{A'B'} {\bar{S}_{B'}}{}^{D} = {\bar{H}_i}{}^{\Lambda'} {\bar{S}_{\Lambda'}}{}^{\Lambda} {\eta_\Lambda}^{DC} {S_C}^{A'} = {H_i}^\Lambda {\eta_\Lambda}^{DC} {S_C}^{A'} = {g_i}^{DA'}, \\ {\bar{g}_i}{}^{A'D} = {g_i}^{DA'}, \quad (i, j, ... = \overline{0,3}). \end{gathered} \tag{14.159}$$

Этим уравнением инволюция ${S_B}^{A'}$ вводится в определение символов ${g_i}^{AA'}$. Такую операцию позволяет осуществить наличие метрического кососимметрического тензора ε_{AB}, с помощью которого можно поднимать и опускать одиночные индексы. Построим теперь операторы Дирака

$$\gamma_i = \begin{pmatrix} 0 & (g^T)_{iAA'} \\ {g_i}^{BB'} & 0 \end{pmatrix}. \tag{14.160}$$

В некотором специальном базисе они будут иметь вид

$$\gamma_0 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad \gamma_1 = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix},$$
$$\gamma_2 = \begin{pmatrix} 0 & 0 & 0 & -i \\ 0 & 0 & i & 0 \\ 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \end{pmatrix}, \quad \gamma_3 = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}, \tag{14.161}$$

Соответственно, инволюция ${S_A}^{A'}$ будет генерировать инволюцию S комплексных операторов Дирака $\bar{\gamma}_i = S\gamma_i S$, и в указанном специальном базисе выполнено $||S|| = ||\gamma^2||$. Таким образом, можно определить майорановский спинор как

$$\psi^i = \gamma^2 \bar{\psi}^i, \tag{14.162}$$

что соответствует изложению [21, с. 115, Приложение E]. При переходе к пространству $\mathbb{R}^6_{(2,4)}$ условие майорантности будет таким ($A, A', ... = \overline{1,4}$, $a, a', ... = \overline{1,2}$)

$$\bar{X}^{A'} = (\bar{X}^{a'}, \bar{Y}_{b'}) = ({\bar{S}^{a'}}_a X^a, {\bar{S}_{b'}}^{\ b} Y_b) = \begin{pmatrix} 0 & {\bar{S}^{a'}}_a \\ {\bar{S}_{b'}}^{\ b} & 0 \end{pmatrix} \begin{pmatrix} Y_b \\ X^a \end{pmatrix} = \bar{S}^{A'A} X_A. \tag{14.163}$$

Однако, на таком спинорном пространстве согласно (8.19) и (8.19′) не существует метрических тензоров, кроме $S_{AA'}$, способных поднимать и опускать одиночные индексы. Таким же будет условие майорантности и для пространства $\mathbb{R}^6$. В этом случае воспользуемся системой ($a, a', ... = \overline{1,4}$)

$$\begin{cases} iR^{ab} Y_b = X^a, \\ -iR_{ab} Z^b = T_a. \end{cases} \tag{14.164}$$

Если наше пространство было бы комплексным $\mathbb{C}^6$, тогда бы эта система устанавливала взаимно-однозначное соответствие между прямолинейными образующими соответствующих квадрик. Свобода представления диктовалась бы однородным решением системы, которое было бы изотропным. Но, из-за действительного вложения $\mathbb{R}^6 \subset \mathbb{C}^6$ изотропные векторы представлены только нулевым вектором. Поэтому возможно взаимно однозначное соответствие между точками соответствующих квадрик

$$R^{ab} = \frac{-i}{Y^k Y_k}(Y_c X_d \varepsilon^{abcd} + 2X^{[a}Y^{b]}), \ Z^b := Y^b = S^{bb'}\bar{Y}_{b'}, \quad T_a := X_a = S_{aa'}\bar{X}^{a'}, \quad X^a Y_a = 0. \tag{14.165}$$

$$X^a X_a + Y^a Y_a(-\frac{1}{4}R^{ab}R_{ab}) = 0. \tag{14.166}$$

(Очевидно, что $R^{ab}R_{ab} > 0$, $X^a X_a = S_{aa'}X^a\bar{X}^{a'} > 0$, $Y^a Y_a = S_{aa'}Y^a\bar{Y}^{a'} > 0$ для ненулевых векторов.) Последнее условие можно переписать следующим образом

$$(X^a, Y_b)(X_a, \frac{-X^d X_d}{Y^c Y_c}Y^b) = 0. \tag{14.167}$$

Таким образом 1-мерной прямолинейной образующей, проходящей через точки (X^a, Y_b) и $(Y^a\frac{-X^d X_d}{Y^c Y_c}, X_b)$ ставится в соответствие ортогональная плоскость $(X_a, \frac{-X^d X_d}{Y^c Y_c}Y_b)$, которая высекает на квадрике $(X^a Y_a = 0)$ 2-мерную плоскую образующую. Совокупность 1-мерной и 2-мерной ортогональных образующих определит 3-мерную плоскую образующую, которая однозначно определит точку на квадрике с координатами R^{ab}. В этом и состоит геометрический смысл майорантности при n=6 для вложения $\mathbb{R}_6 \subset \mathbb{C}\mathbb{R}_6$. Кстати говоря, R^{ab} представляет собой изотропный вектор изотропного конуса пространства $R^8_{(1,7)}$. Соответствующая квадрика высекается сечением этого конуса.

Example 14.1. *Итак, пусть* $Y^k Y_k = 2$, $X^a Y_a = 0$*, тогда в специальном базисе возможно представление*

$$\begin{gathered} iR^{12} = -i\ \overline{R^{34}} = X^{[1}Y^{2]} + \overline{X^{[3}Y^{4]}}, \quad iR^{13} = i\ \overline{R^{24}} = X^{[1}Y^{3]} - \overline{X^{[2}Y^{4]}}, \\ iR^{14} = -i\ \overline{R^{23}} = X^{[1}Y^{4]} + \overline{X^{[2}Y^{3]}}. \end{gathered} \tag{14.168}$$

14.5.7 Теорема о двух квадриках.

В этом параграфе рассказывается о том, как можно интерпретировать принцип тройственности Картана на основании предыдущих результатов и свести их воедино. Вывод всех результатов этого параграфа сделан на основе [6].

Таким образом доказана теорема

Теорема 14.3. *(принцип тройственности для двух B- цилиндров).*
В проективном пространстве $\mathbb{CP}_7$ *существуют две квадрики (два B - цилиндра), обладающие следующими общими свойствами*

1. *Плоская образующая* $\mathbb{CP}_3$ *одной квадрики взаимно однозначно определит точку* R *другой.*

2. *Плоская образующая* $\mathbb{CP}_2$ *одной квадрики однозначно определит точку* R *другой. Но точке* R *можно сопоставить многообразие плоских образующих* $\mathbb{CP}_2$*, принадлежащих одной плоской образующей* $\mathbb{CP}_3$ *второй квадрики.*

3. *Прямолинейная образующая* $\mathbb{CP}_1$ *одной квадрики взаимооднозначно определит прямолинейную образующую* $\mathbb{CP}_1$ *из другой. Причем все прямолинейные образующие, принадлежащие одной плоской образующей* $\mathbb{CP}_3$ *первой квадрики, определят пучок с центром в точке* R*, принадлежащий второй квадрике.*

Эта теорема на самом деле является обобщением соответствия Кляйна. Докажем это.

Доказательство. Рассмотрим на квадрике $\mathbb{C}Q_6$ только те образующие, которые имеют вид $X^A = (0, Y_b)$. Многообразие таких образующих диффеоморфно $\mathbb{CP}_3$. При этом каждой такой образующей можно поставить в соответствие точку квадрики $\mathbb{C}Q_4 \subset \mathbb{C}\tilde{Q}_6$. Тогда $R^{ab}Y_b = 0$. До конца доказательства положим $\mathbf{A}, \mathbf{B}, \mathbf{A}', \mathbf{B}', ... = \overline{1,2}$. Кроме того, рассмотрим спинорное представление твисторов согласно [22, формулы (6.1.24) и (6.2.18)]

$$Y_b = (\pi_{\mathbf{B}}, \bar{\omega}^{\mathbf{B}'}),$$

$$R^{ab} = const \begin{pmatrix} -\frac{1}{2}\varepsilon^{\mathbf{AB}} r_c r^c & iR^{\mathbf{A}}{}_{\mathbf{B'}} \\ -i\bar{R}_{\mathbf{A}'}{}^{\mathbf{B}} & \bar{\varepsilon}_{\mathbf{A}'\mathbf{B}'} \end{pmatrix}. \tag{14.169}$$

Поэтому уравнение $R^{ab}Y_b = 0$ перепишется в виде системы двух уравнений

$$\begin{cases} -\frac{1}{2}\varepsilon^{\mathbf{AB}} r_c r^c \pi_{\mathbf{B}} + iR^{\mathbf{A}}{}_{\mathbf{B}'}\bar{\omega}^{\mathbf{B}'} &= 0, \\ -i\bar{R}_{\mathbf{A}'}{}^{\mathbf{B}}\pi_{\mathbf{B}} + \bar{\varepsilon}_{\mathbf{A}'\mathbf{B}'}\bar{\omega}^{\mathbf{B}'} &= 0, \end{cases} \tag{14.170}$$

из которых будет существенным только одно

$$ir^{\mathbf{AA}'}\bar{\pi}_{\mathbf{A}'} = \omega^{\mathbf{A}}. \tag{14.171}$$

Здесь мы воспользовались операцией сопряжения и метрическими спинорами $\varepsilon_{\mathbf{A'B'}}, \varepsilon_{\mathbf{AB}}$, с помощью которых подымаются и опускаются спинорные индексы и которые при сопряжении переходят друг в друга. Этим определится система

$$\begin{cases} ir^{\mathbf{AA}'}\bar{\pi}_{\mathbf{A}'} &= \bar{\omega}^{\mathbf{A}'}, \\ ir^{\mathbf{AA}'}\bar{\eta}_{\mathbf{A}'} &= \xi^{\mathbf{A}}, \end{cases} \quad Y_b = (\pi_{\mathbf{B}}, \bar{\omega}^{\mathbf{B}'}), \quad T_b = (\eta_{\mathbf{B}}, \bar{\xi}^{\mathbf{B}'}). \tag{14.172}$$

Эта система совпадает с системой [22, формула (6.2.14)], которая, в свою очередь, приводит к соответствию Кляйна. □

Следует в заключении отметить, что из этой теоремы следует принцип тройственности Картана: существует 3 диффеоморфных многообразия - многообразие точек квадрики и 2 многообразия плоских образующих I и II семейств. Это действительно так, поскольку две построенные квадрики можно отождествить, например, с помощью тензора $P_\Lambda{}^L$. При этом многообразие точек квадрики будет диффеоморфно многообразию плоских образующих одного из двух семейства. Кроме того, поскольку принцип тройственности Картана выполнен, то операторы $\eta^i{}_{KL}$ для вложения $\mathbb{R}^8 \subset \mathbb{C}^8$ определят алгебру октав, поскольку удовлетворяют уравнению Клиффорда. Это утверждение основано на результатах, приведенных в монографии [22, т. 2, с. 543-544], где рассматриваются структурные константы этой алгебры.

14.6 Явное построение аналогов операторов Ли при n=4.

В этом параграфе рассказывается о том, как можно в некотором базисе получить представление операторов P, с помощью которых можно построить спинорный аналог операторов Ли. Вывод всех результатов этого параграфа сделан на основе [9].

Пусть на пространстве $\mathbb{R}^8_{(4,4)}$ задана метрика

$$dS^2 = -dU^2 + dS^2 + dV^2 - dW^2 + dX^2 - dY^2 - dT^2 + dZ^2, \tag{14.173}$$

тогда в некотором базисе связующие операторы $\eta_\Lambda{}^{KL}$ будут иметь вид

$$\begin{array}{llll}
\eta_3{}^{12} = \frac{1}{\sqrt{2}}, & \eta_3{}^{34} = \frac{1}{\sqrt{2}}, & \eta_4{}^{12} = \frac{1}{\sqrt{2}}, & \eta_4{}^{34} = -\frac{1}{\sqrt{2}}, \\
\eta_3{}^{78} = \frac{1}{\sqrt{2}}, & \eta_3{}^{56} = \frac{1}{\sqrt{2}}, & \eta_4{}^{78} = \frac{1}{\sqrt{2}}, & \eta_4{}^{56} = -\frac{1}{\sqrt{2}}, \\
\eta_7{}^{14} = \frac{1}{\sqrt{2}}, & \eta_7{}^{23} = -\frac{1}{\sqrt{2}}, & \eta_8{}^{14} = \frac{1}{\sqrt{2}}, & \eta_8{}^{23} = \frac{1}{\sqrt{2}}, \\
\eta_7{}^{67} = \frac{1}{\sqrt{2}}, & \eta_7{}^{58} = -\frac{1}{\sqrt{2}}, & \eta_8{}^{67} = \frac{1}{\sqrt{2}}, & \eta_8{}^{58} = \frac{1}{\sqrt{2}}, \\
\eta_6{}^{13} = \frac{1}{\sqrt{2}}, & \eta_6{}^{24} = \frac{1}{\sqrt{2}}, & \eta_5{}^{13} = \frac{1}{\sqrt{2}}, & \eta_5{}^{24} = -\frac{1}{\sqrt{2}}, \\
\eta_6{}^{68} = -\frac{1}{\sqrt{2}}, & \eta_6{}^{57} = -\frac{1}{\sqrt{2}}, & \eta_5{}^{68} = -\frac{1}{\sqrt{2}}, & \eta_5{}^{57} = \frac{1}{\sqrt{2}}, \\
\eta_2{}^{15} = \frac{1}{\sqrt{2}}, & \eta_2{}^{51} = \frac{1}{\sqrt{2}}, & \eta_1{}^{15} = -\frac{1}{\sqrt{2}}, & \eta_1{}^{51} = \frac{1}{\sqrt{2}}, \\
\eta_2{}^{26} = \frac{1}{\sqrt{2}}, & \eta_2{}^{62} = \frac{1}{\sqrt{2}}, & \eta_1{}^{26} = -\frac{1}{\sqrt{2}}, & \eta_1{}^{62} = \frac{1}{\sqrt{2}}, \\
\eta_2{}^{37} = \frac{1}{\sqrt{2}}, & \eta_2{}^{73} = \frac{1}{\sqrt{2}}, & \eta_1{}^{37} = -\frac{1}{\sqrt{2}}, & \eta_1{}^{73} = \frac{1}{\sqrt{2}}, \\
\eta_2{}^{48} = \frac{1}{\sqrt{2}}, & \eta_2{}^{84} = \frac{1}{\sqrt{2}}, & \eta_1{}^{48} = -\frac{1}{\sqrt{2}}, & \eta_1{}^{84} = \frac{1}{\sqrt{2}},
\end{array} \tag{14.174}$$

а существенные координаты операторов $\eta^{\Lambda}{}_{KL}$ будут такими

$$\begin{array}{llll}
\eta^{3}{}_{12}=\frac{1}{\sqrt{2}}, & \eta^{3}{}_{34}=\frac{1}{\sqrt{2}}, & \eta^{4}{}_{12}=\frac{1}{\sqrt{2}}, & \eta^{4}{}_{34}=-\frac{1}{\sqrt{2}},\\
\eta^{3}{}_{78}=\frac{1}{\sqrt{2}}, & \eta^{3}{}_{56}=\frac{1}{\sqrt{2}}, & \eta^{4}{}_{78}=\frac{1}{\sqrt{2}}, & \eta^{4}{}_{56}=-\frac{1}{\sqrt{2}},\\
\eta^{7}{}_{14}=\frac{1}{\sqrt{2}}, & \eta^{7}{}_{23}=-\frac{1}{\sqrt{2}}, & \eta^{8}{}_{14}=\frac{1}{\sqrt{2}}, & \eta^{8}{}_{23}=\frac{1}{\sqrt{2}},\\
\eta^{7}{}_{67}=\frac{1}{\sqrt{2}}, & \eta^{7}{}_{58}=-\frac{1}{\sqrt{2}}, & \eta^{8}{}_{67}=\frac{1}{\sqrt{2}}, & \eta^{8}{}_{58}=\frac{1}{\sqrt{2}},\\
\eta^{6}{}_{13}=\frac{1}{\sqrt{2}}, & \eta^{6}{}_{24}=\frac{1}{\sqrt{2}}, & \eta^{5}{}_{13}=\frac{1}{\sqrt{2}}, & \eta^{5}{}_{24}=-\frac{1}{\sqrt{2}},\\
\eta^{6}{}_{68}=-\frac{1}{\sqrt{2}}, & \eta^{6}{}_{57}=-\frac{1}{\sqrt{2}}, & \eta^{5}{}_{68}=-\frac{1}{\sqrt{2}}, & \eta^{5}{}_{57}=\frac{1}{\sqrt{2}},\\
\eta^{2}{}_{15}=\frac{1}{\sqrt{2}}, & \eta^{2}{}_{51}=\frac{1}{\sqrt{2}}, & \eta^{1}{}_{15}=-\frac{1}{\sqrt{2}}, & \eta^{1}{}_{51}=\frac{1}{\sqrt{2}},\\
\eta^{2}{}_{26}=\frac{1}{\sqrt{2}}, & \eta^{2}{}_{62}=\frac{1}{\sqrt{2}}, & \eta^{1}{}_{26}=-\frac{1}{\sqrt{2}}, & \eta^{1}{}_{62}=\frac{1}{\sqrt{2}},\\
\eta^{2}{}_{37}=\frac{1}{\sqrt{2}}, & \eta^{2}{}_{73}=\frac{1}{\sqrt{2}}, & \eta^{1}{}_{37}=-\frac{1}{\sqrt{2}}, & \eta^{1}{}_{73}=\frac{1}{\sqrt{2}},\\
\eta^{2}{}_{48}=\frac{1}{\sqrt{2}}, & \eta^{2}{}_{84}=\frac{1}{\sqrt{2}}, & \eta^{1}{}_{48}=-\frac{1}{\sqrt{2}}, & \eta^{1}{}_{84}=\frac{1}{\sqrt{2}}.
\end{array} \tag{14.175}$$

Поэтому в том же базисе

$$\begin{array}{cc}
\sqrt{2}\parallel P_{\tilde{A}}{}^{\tilde{\tilde{B}}}\parallel= & \sqrt{2}\parallel P^{\tilde{A}}{}_{\tilde{\tilde{B}}}\parallel= \\
\begin{pmatrix}
1 & 0 & 0 & 0 & -1 & 0 & 0 & 0\\
1 & 0 & 0 & 0 & 1 & 0 & 0 & 0\\
0 & 1 & 0 & 0 & 0 & 1 & 0 & 0\\
0 & 1 & 0 & 0 & 0 & -1 & 0 & 0\\
0 & 0 & 1 & 0 & 0 & 0 & 1 & 0\\
0 & 0 & 1 & 0 & 0 & 0 & -1 & 0\\
0 & 0 & 0 & 1 & 0 & 0 & 0 & -1\\
0 & 0 & 0 & 1 & 0 & 0 & 0 & 1
\end{pmatrix}, &
\begin{pmatrix}
1 & 0 & 0 & 0 & -1 & 0 & 0 & 0\\
1 & 0 & 0 & 0 & 1 & 0 & 0 & 0\\
0 & 1 & 0 & 0 & 0 & 1 & 0 & 0\\
0 & 1 & 0 & 0 & 0 & -1 & 0 & 0\\
0 & 0 & 1 & 0 & 0 & 0 & 1 & 0\\
0 & 0 & 1 & 0 & 0 & 0 & -1 & 0\\
0 & 0 & 0 & 1 & 0 & 0 & 0 & -1\\
0 & 0 & 0 & 1 & 0 & 0 & 0 & 1
\end{pmatrix},
\end{array} \tag{14.176}$$

Следовательно, алгоритм 8.1 примет вид

Алгоритм 14.1.

1. На первом шаге используем в качестве операторов (6.24) и (6.30) операторы

$$
\begin{aligned}
&\underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} e^{i\frac{\pi}{8}}\delta^{a}{}_{\tilde{\tilde{a}}} & 0 & \pm ie^{i\frac{\pi}{8}}\varepsilon^{a\tilde{\tilde{p}}} & 0 \\ 0 & e^{-i\frac{\pi}{8}}\delta_{c}{}^{\tilde{\tilde{c}}} & 0 & \mp e^{-i\frac{\pi}{8}}\varepsilon_{c\tilde{\tilde{q}}} \end{pmatrix}}_{:=(\tilde{\tilde{m}}_{1_{\pm}})^{A}{}_{\tilde{\tilde{A}}}} \cdot \\
&\cdot \underbrace{\begin{pmatrix} \alpha_1\varepsilon^{\tilde{\tilde{a}}\tilde{\tilde{b}}} & \eta_{\alpha}{}^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{d}}} & \gamma_1\delta^{\tilde{\tilde{a}}}{}_{\tilde{\tilde{k}}} & 0 \\ -(\eta^{T})_{\alpha\tilde{\tilde{c}}}{}^{\tilde{\tilde{b}}} & \beta_1\varepsilon_{\tilde{\tilde{c}}\tilde{\tilde{d}}} & 0 & \gamma_1\delta_{\tilde{\tilde{c}}}{}^{\tilde{\tilde{l}}} \\ \delta_1\delta_{\tilde{\tilde{p}}}{}^{\tilde{\tilde{b}}} & 0 & \beta_1\varepsilon_{\tilde{\tilde{p}}\tilde{\tilde{k}}} & \eta_{\alpha\tilde{\tilde{p}}}{}^{\tilde{\tilde{l}}} \\ 0 & \delta_1\delta^{\tilde{\tilde{q}}}{}_{\tilde{\tilde{d}}} & -(\eta^{T})_{\alpha}{}^{\tilde{\tilde{q}}}{}_{\tilde{\tilde{k}}} & \alpha_1\varepsilon^{\tilde{\tilde{q}}\tilde{\tilde{l}}} \end{pmatrix}}_{\eta_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}} \underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} 0 & e^{i\frac{\pi}{8}}\varepsilon_{\tilde{\tilde{b}}d} \\ -e^{-i\frac{\pi}{8}}\varepsilon^{\tilde{\tilde{d}}b} & 0 \\ 0 & \pm e^{i\frac{\pi}{8}}\delta^{\tilde{\tilde{k}}}{}_{d} \\ \mp e^{-i\frac{\pi}{8}}i\delta_{\tilde{\tilde{l}}}{}^{b} & 0 \end{pmatrix}}_{:=(\tilde{m}^{T}_{1_{\pm}})_{\tilde{\tilde{B}}}{}^{B}} = \\
&= \underbrace{\begin{pmatrix} \eta_{\alpha}{}^{ab} & \underbrace{\frac{e^{i\frac{\pi}{4}}}{2}\left((-\alpha_1 - i\beta_1) \pm (\gamma_1 - i\delta_1)\right)}_{:=(\phi_{1_{\pm}})}\delta^{a}{}_{d} \\ \underbrace{\frac{ie^{-i\frac{\pi}{4}}}{2}\left((-\alpha_1 - i\beta_1) \mp (\gamma_1 - i\delta_1)\right)}_{:=(\psi_{1_{\pm}})}\delta_{c}{}^{b} & -(\eta^{T})_{\alpha cd} \end{pmatrix}}_{:=\eta_{1_{\pm}\tilde{\Lambda}}{}^{AB}},
\end{aligned}
\tag{14.177}
$$

$$\alpha_1 = \tfrac{1}{2}(\eta_3 + \eta_4), \quad \beta_1 = \tfrac{1}{2}(\eta_3 - \eta_4), \quad \gamma_1 = \tfrac{1}{2}(\eta_2 - \eta_1), \quad \delta_1 = \tfrac{1}{2}(\eta_2 + \eta_1).$$

Здесь $\varepsilon^{\tilde{\tilde{a}}\tilde{\tilde{b}}}$ *- кососимметрический метрический спинор для связующих операторов* $\eta_{\alpha\tilde{\tilde{p}}}{}^{\tilde{\tilde{l}}}$ *(*$a, b, ..., \tilde{\tilde{a}}, \tilde{\tilde{b}}, ... = \overline{1,2},\ \tilde{\Lambda}, ..., \tilde{\tilde{A}}, \tilde{\tilde{B}}, ... = \overline{1,8},\ A, B, ... = \overline{1,4}$*).*

2. На втором шаге используем в качестве операторов (6.24) и (6.30) операторы

$$\begin{aligned}&\underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} e^{i\frac{\pi}{8}} & \pm i e^{i\frac{\pi}{8}} & 0 & 0 \\ 0 & 0 & e^{-i\frac{\pi}{8}} & \pm e^{-i\frac{\pi}{8}} \end{pmatrix}}_{:=(\tilde{\tilde{m}}_{2_\pm})^{A}{}_{\tilde{\tilde{A}}}} \cdot \\ &\cdot \underbrace{\begin{pmatrix} \alpha_2 & \gamma_2 & \phi_{1_\pm} & 0 \\ \delta_2 & \beta_2 & 0 & \phi_{1_\pm} \\ \psi_{1_\pm} & 0 & -\beta_2 & \gamma_2 \\ 0 & \psi_{1_\pm} & \delta_2 & -\alpha_2 \end{pmatrix}}_{(\eta_{1_\pm})_{\tilde{\Lambda}}{}^{\tilde{\tilde{A}}\tilde{\tilde{B}}}} \underbrace{\frac{1}{\sqrt{2}}\begin{pmatrix} e^{i\frac{\pi}{8}} & 0 \\ \pm e^{i\frac{\pi}{8}} & 0 \\ 0 & e^{-i\frac{\pi}{8}} \\ 0 & \mp i e^{-i\frac{\pi}{8}} \end{pmatrix}}_{:=(\tilde{m}^{T}_{2_\pm})_{\tilde{\tilde{B}}}{}^{B}} = \\ &= \underbrace{\begin{pmatrix} \underbrace{\frac{e^{i\frac{\pi}{4}}}{2}((\alpha_2 + i\beta_2) \pm (\gamma_2 + i\delta_2))}_{:=\phi_{2_\pm}} & \underbrace{\frac{e^{i\frac{\pi}{4}}}{2}((-\alpha_1 - i\beta_1) \pm (\gamma_1 - i\delta_1))}_{:=\phi_{1_\pm}} \\ \underbrace{\frac{ie^{-i\frac{\pi}{4}}}{2}((-\alpha_1 - i\beta_1) \mp (\gamma_1 - i\delta_1))}_{:=\psi_{1_\pm}} & \underbrace{\frac{ie^{-i\frac{\pi}{4}}}{2}((\alpha_2 + i\beta_2) \mp (\gamma_2 + i\delta_2))}_{:=\psi_{2_\pm}} \end{pmatrix}}_{:=(\eta_{1,2,3,4})_{\tilde{\Lambda}}{}^{AB}}, \\ &\alpha_2 = \tfrac{1}{2}(\eta_7 + \eta_8), \quad \beta_2 = \tfrac{1}{2}(\eta_7 - \eta_8), \quad \gamma_2 = \tfrac{1}{2}(-\eta_6 - \eta_5), \quad \delta_2 = \tfrac{1}{2}(\eta_6 - \eta_5).\end{aligned} \tag{14.178}$$

($\tilde{\Lambda}, ... = \overline{1,8}$, $\tilde{\tilde{A}}, \tilde{\tilde{B}}, ... = \overline{1,4}$, $A, B, ... = \overline{1,2}$).

3. Для касательного пространства операторы ${m_\Lambda}^{\tilde{\Lambda}}$ будут иметь вид

$$\begin{aligned} {m_\Lambda}^{\tilde{\Lambda}} &:= \tfrac{1}{\sqrt{2}} \begin{pmatrix} -i & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -i & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & i & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & i & 1 \end{pmatrix}, \\ {m^\Lambda}_{\tilde{\Lambda}} &:= \tfrac{1}{\sqrt{2}} \begin{pmatrix} i & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & i & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -i & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -i & 1 \end{pmatrix}. \end{aligned} \tag{14.179}$$

Тогда операторы

$$\underbrace{\begin{pmatrix} \tfrac{1}{2}(\underbrace{\frac{1}{\sqrt{2}}(\eta_8 + i\eta_7)}_{:=(\eta^{\mathbb{C}})_4} \mp \underbrace{\frac{1}{\sqrt{2}}(\eta_6 + i\eta_5)}_{:=(\eta^{\mathbb{C}})_3}) & \tfrac{1}{2}(\underbrace{\frac{1}{\sqrt{2}}(-\eta_4 - i\eta_3)}_{:=(\eta^{\mathbb{C}})_2} \pm \underbrace{\frac{1}{\sqrt{2}}(\eta_2 - i\eta_1)}_{:=(\eta^{\mathbb{C}})_1}) \\ \tfrac{1}{2}(\underbrace{\frac{1}{\sqrt{2}}(-\eta_4 - i\eta_3)}_{:=(\eta^{\mathbb{C}})_2} \mp \underbrace{\frac{1}{\sqrt{2}}(\eta_2 - i\eta_1)}_{:=(\eta^{\mathbb{C}})_1}) & \tfrac{1}{2}(\underbrace{\frac{1}{\sqrt{2}}(\eta_8 + i\eta_7)}_{:=(\eta^{\mathbb{C}})_4} \pm \underbrace{\frac{1}{\sqrt{2}}(\eta_6 + i\eta_5)}_{:=(\eta^{\mathbb{C}})_3}) \end{pmatrix}}_{:=(\eta_{1,2,3,4})_{\Lambda}{}^{AB}} \tag{14.180}$$

будут связующими операторами для комплексной реализации при n=4. Таким образом *построен переход от вещественной реализации связующих операторов при n=8 к их комплексным аналогам при n=4.*

Таким образом, операторы $2((\tilde{M}_K^*)^T)_{\tilde{\tilde{A}}}{}^A$ согласно (6.33) и алгоритму 8.1 будут иметь вид

$$\begin{pmatrix} 0 & 0 & 0 & e^{i\frac{\pi}{8}} \\ 0 & 0 & -e^{i\frac{\pi}{8}} & 0 \\ 0 & -e^{-i\frac{\pi}{8}} & 0 & 0 \\ e^{-i\frac{\pi}{8}} & 0 & 0 & 0 \\ 0 & 0 & p_1 e^{i\frac{\pi}{8}} & 0 \\ 0 & 0 & 0 & p_1 e^{i\frac{\pi}{8}} \\ -p_1 i e^{-i\frac{\pi}{8}} & 0 & 0 & 0 \\ 0 & -p_1 i e^{-i\frac{\pi}{8}} & 0 & 0 \end{pmatrix} \begin{pmatrix} e^{i\frac{\pi}{8}} & 0 \\ p_2 e^{i\frac{\pi}{8}} & 0 \\ 0 & e^{-i\frac{\pi}{8}} \\ 0 & -p_2 i e^{-i\frac{\pi}{8}} \end{pmatrix} = \begin{pmatrix} 0 & -p_2 i \\ 0 & -1 \\ -p_2 & 0 \\ 1 & 0 \\ 0 & p_1 \\ 0 & -p_1 p_2 i \\ -p_1 i & 0 \\ -p_1 p_2 i & 0 \end{pmatrix}. \tag{14.181}$$

При этом p_1, p_2 принимают два значения ±1, обеспечивая 4 различных варианта операторов $\tilde{M}_K^*$. Осталось применить эти операторы и операторы $m_\Lambda{}^{\tilde{A}}$, $\bar{m}_{\Lambda'}{}^{\tilde{A}}$ к операторам $P_{\tilde{A}}{}^{\tilde{\tilde{A}}}$

$$(P_K^*)_\Lambda{}^A := \frac{1+i}{4}\begin{pmatrix} 0 & p_1 - p_2 \\ 0 & 1 - p_1 p_2 \\ p_1 - p_2 & 0 \\ 1 - p_1 p_2 & 0 \end{pmatrix}, \quad (\bar{P}_K^*)_{\Lambda'}{}^A := \frac{1-i}{4}\begin{pmatrix} 0 & p_1 + p_2 \\ 0 & 1 + p_1 p_2 \\ -p_1 - p_2 & 0 \\ 1 + p_1 p_2 & 0 \end{pmatrix}, \tag{14.182}$$

$$(P_K)^\Lambda{}_A := \frac{1-i}{4}\begin{pmatrix} 0 & p_1 - p_2 \\ 0 & 1 - p_1 p_2 \\ p_1 - p_2 & 0 \\ 1 - p_1 p_2 & 0 \end{pmatrix}, \quad (\bar{P}_K)^{\Lambda'}{}_A := \frac{1+i}{4}\begin{pmatrix} 0 & p_1 + p_2 \\ 0 & 1 + p_1 p_2 \\ -p_1 - p_2 & 0 \\ 1 + p_1 p_2 & 0 \end{pmatrix}. \tag{14.183}$$

Очевидно, что из 4 вариантов для $K = \overline{1,4}$, существенными будут только два $(P_2^*)_\Lambda{}^A$, $(P_3^*)_\Lambda{}^A$ и $(\bar{P}_1^*)_{\Lambda'}{}^A$, $(\bar{P}_4^*)_{\Lambda'}{}^A$ $(\Lambda, ... = \overline{1,4})$. При условии постоянства таких операторов при обычном и ковариантном дифференцировании коэффициенты связности $\Gamma_{\tilde{A}\Psi}{}^\Phi$ распадаются на 8 существенных компонент

$$\Gamma_{\tilde{A}\Psi}{}^\Phi \to \begin{cases} (\Gamma_K)_{\Lambda A}{}^B := \Gamma_{\Lambda\Psi}{}^\Phi (P_K)^\Psi{}_A (P_K^*)_\Phi{}^B, \\ (\Gamma_K)_{\Lambda A'}{}^{B'} := \Gamma_{\Lambda\Psi}{}^\Phi (P_K)^\Psi{}_{A'} (P_K^*)_\Phi{}^{B'}, \\ (\bar{\Gamma}_K)_{\Lambda' A'}{}^{B'} := \bar{\Gamma}_{\Lambda'\Psi'}{}^{\Phi'} (\bar{P}_K)^{\Psi'}{}_{A'} (\bar{P}_K^*)_{\Phi'}{}^{B'}, \\ (\bar{\Gamma}_K)_{\Lambda' A}{}^B := \bar{\Gamma}_{\Lambda'\Psi'}{}^{\Phi'} (\bar{P}_K)^{\Psi'}{}_A (\bar{P}_K^*)_{\Phi'}{}^B. \end{cases} \tag{14.184}$$

В свою очередь, коэффициенты ${\Gamma_{\tilde{\varLambda}\Psi}}^{\Phi}$ определяются из уравнения $\nabla_{\tilde{\varLambda}} {m_{\Psi}}^{\tilde{\varPsi}} = 0$. Таким образом теоремы 11.2, 11.3 остаются верными и в этом случае. В случае вещественного вложения с помощью операторов ${H_i}^{\Lambda}$ при условии ковариантного и обычного постоянства операторов ${(P_2^*)_i}^A$, ${(P_3^*)_i}^A$ и ${(P_1^*)_i}^A$, ${(P_4^*)_i}^A$ коэффициенты связности ${\Gamma_{\tilde{\varLambda}i}}^j$ распадаются на 4 существенные компоненты

$$
{\Gamma_{\tilde{\varLambda}i}}^j \to \begin{cases} {(\Gamma_K)_{kA}}^B := {\Gamma_{ki}}^j {(P_K)^i}_A {(P_K^*)_j}^B, \\ {(\bar{\Gamma}_K)_{kA'}}^{B'} := {\Gamma_{ki}}^j {(\bar{P}_K)^i}_{A'} {(\bar{P}_K^*)_j}^{B'}. \end{cases} \tag{14.185}
$$

Однако, условие действительности ограничивает выбор спинора $X^{\tilde{\tilde{A}}}$ при определении операторов ${P_{\tilde{\varLambda}}}^{\tilde{\tilde{A}}}$

$$
X^{\tilde{\tilde{A}}} = {\tilde{S}_{\tilde{\tilde{B}}}}^{\tilde{\tilde{A}}} X^{\tilde{\tilde{B}}}, \quad X^{\tilde{\tilde{A}}} = \tilde{\tilde{S}}^{\tilde{\tilde{B}}\tilde{\tilde{A}}} X_{\tilde{\tilde{B}}}, \tag{14.186}
$$

где ${\tilde{S}_{\tilde{\tilde{B}}}}^{\tilde{\tilde{A}}}$ ($S^{\tilde{\tilde{B}}\tilde{\tilde{A}}}$) есть действительная реализация комплексной инволюции ${\tilde{S}_B}^{A'}$ ($\tilde{S}^{BA'}$), которая является образом инволюции ${S_{\Lambda}}^{\Psi'}$ в касательном пространстве согласно следствию 8.4. Необходимо добавить, что существует несобственное ортогональное преобразование

$$
(-\eta_{\Lambda}\eta^{\Psi} + {\delta_{\Lambda}}^{\Psi}){\eta_{\Psi}}^{AB} = {\eta_{\Lambda}}^{BA}, \tag{14.187}
$$

осуществляющее транспонирование связующих операторов (14.84).

15 Благодарности

Эта статья является ответами на часто задаваемые вопросы. Поэтому автор благодарен всем, кто эти вопросы верно сформулировал, что дало возможность найти на них ответ. Отдельная благодарность К. Шарнхорсту за копии статей, без которого этой публикации не было.

16 Приложение

16.1 Доказательство формул о метрических тензорах и инволюции

Необходимо доказать тождества (8.19).

Случай I)., метрический тензор $\varepsilon_{AB} = -\varepsilon_{BA}$.

$$\begin{gathered}
{\eta_\Lambda}^{A_1B_1}\varepsilon_{A_1A_{(n/2+q+1)}}\varepsilon_{B_1B_{(n/2+q+1)}} = \\
= {\eta_\Lambda}^{A_1B_1}(\varepsilon_{(n/2+q)})_{A_{(n/2+q)}A_{(n/2+q+1)}}...(\varepsilon_1)_{A_1A_2}(\varepsilon_2)^{A_3A_2} \\
(\varepsilon_{(n/2+q)})_{B_{(n/2+q)}B_{(n/2+q+1)}}...(\varepsilon_1)_{B_1B_2}(\varepsilon_2)^{B_3B_2} = \\
= (-\eta_{\Lambda B_2A_2} + (\eta_1)_\Lambda\varepsilon_{1A_2B_2})(\varepsilon_{(n/2+q)})_{A_{(n/2+q)}A_{(n/2+q+1)}}...(\varepsilon_2)^{A_3A_2} \\
(\varepsilon_{(n/2+q)})_{B_{(n/2+q)}B_{(n/2+q+1)}}...(\varepsilon_2)^{B_3B_2} = \\
= -(-{\eta_\Lambda}^{A_3B_3} + (\eta_2)_\Lambda{\varepsilon_2}^{A_3B_3})(\varepsilon_{(n/2+q)})_{A_{(n/2+q)}A_{(n/2+q+1)}}\cdots \\
(\varepsilon_{(n/2+q)})_{B_{(n/2+q)}B_{(n/2+q+1)}}\cdots + (\eta_1)_\Lambda(\varepsilon_1)_{A_{(n/2+q+1)}B_{(n/2+q+1)}} = \\
= {\eta_\Lambda}^{A_3B_3}(\varepsilon_{(n/2+q)})_{A_{(n/2+q)}A_{(n/2+q+1)}}...(\varepsilon_{(n/2+q)})_{B_{(n/2+q)}B_{(n/2+q+1)}}...+ \\
+\sum_{Q=1}^{2}(\eta_Q)_\Lambda(\varepsilon_Q)_{A_{(n/2+q+1)}B_{(n/2+q+1)}} = \\
= -\eta_{\Lambda B_{(n/2+q+1)}A_{(n/2+q+1)}} + \sum_{Q=1}^{n/2+q}(\eta_Q)_\Lambda(\varepsilon_Q)_{A_{(n/2+q+1)}B_{(n/2+q+1)}} = \\
= \eta_{\Lambda A_{(n/2+q+1)}B_{(n/2+q+1)}}.
\end{gathered}\tag{16.1}$$

Случай I)., метрический тензор $\tilde{\varepsilon}_{AB} = \tilde{\varepsilon}_{BA}$.

$$\begin{gathered}
{\eta_\Lambda}^{A_1B_1}\tilde{\varepsilon}_{A_1A_{(n/2+\tilde{q}+1)}}\tilde{\varepsilon}_{B_1B_{(n/2+\tilde{q}+1)}} = \\
= {\eta_\Lambda}^{A_1B_1}(\varepsilon_{\widetilde{(n/2+\tilde{q})}})_{A_{(n/2+\tilde{q})}A_{(n/2+\tilde{q}+1)}}...(\varepsilon_{\tilde{1}})_{A_1A_2}(\varepsilon_{\tilde{2}})^{A_3A_2} \\
(\varepsilon_{\widetilde{(n/2+\tilde{q})}})_{B_{(n/2+\tilde{q})}B_{(n/2+\tilde{q}+1)}}...(\varepsilon_{\tilde{1}})_{B_1B_2}(\varepsilon_{\tilde{2}})^{B_3B_2} = \\
= (\eta_{\Lambda B_2A_2} + (\eta_{\tilde{1}})_\Lambda\varepsilon_{\tilde{1}A_2B_2})(\varepsilon_{\widetilde{(n/2+\tilde{q})}})_{A_{(n/2+\tilde{q})}A_{(n/2+\tilde{q}+1)}}...(\varepsilon_{\tilde{2}})^{A_3A_2} \\
(\varepsilon_{\widetilde{(n/2+\tilde{q})}})_{B_{(n/2+\tilde{q})}B_{(n/2+\tilde{q}+1)}}...(\varepsilon_{\tilde{2}})^{B_3B_2} = \\
= ({\eta_\Lambda}^{A_3B_3} + (\eta_{\tilde{2}})_\Lambda{\varepsilon_{\tilde{2}}}^{B_3A_3})(\varepsilon_{\widetilde{(n/2+\tilde{q})}})_{A_{(n/2+\tilde{q})}A_{(n/2+\tilde{q}+1)}}...(\varepsilon_{\widetilde{(n/2+\tilde{q})}})_{B_{(n/2+\tilde{q})}B_{(n/2+\tilde{q}+1)}}...+ \\
+(\eta_{\tilde{1}})_\Lambda(\varepsilon_{\tilde{1}})_{A_{(n/2+\tilde{q}+1)}B_{(n/2+\tilde{q}+1)}} = \\
= {\eta_\Lambda}^{A_3B_3}(\varepsilon_{\widetilde{(n/2+\tilde{q})}})_{A_{(n/2+\tilde{q})}A_{(n/2+\tilde{q}+1)}}...(\varepsilon_{\widetilde{(n/2+\tilde{q})}})_{B_{(n/2+\tilde{q})}B_{(n/2+\tilde{q}+1)}}...+ \\
+\sum_{\tilde{Q}=1}^{2}(\eta_{\tilde{Q}})_\Lambda(\varepsilon_{\tilde{Q}})_{A_{(n/2+\tilde{q}+1)}B_{(n/2+\tilde{q}+1)}} = \\
= \eta_{\Lambda B_{(n/2+\tilde{q}+1)}A_{(n/2+\tilde{q}+1)}} + \sum_{\tilde{Q}=1}^{n/2+\tilde{q}}(\eta_{\tilde{Q}})_\Lambda(\varepsilon_{\tilde{Q}})_{A_{(n/2+\tilde{q}+1)}B_{(n/2+\tilde{q}+1)}} = \\
= \eta_{\Lambda A_{(n/2+\tilde{q}+1)}B_{(n/2+\tilde{q}+1)}}.
\end{gathered}\tag{16.2}$$

Случай II)., инволюция ${E_A}^B {E_B}^C = -{\delta_A}^C$.

$$\begin{gathered}
{\eta_\Lambda}^{A_1 B_1} {E_{A_1}}^{A_{(n/2+q)}} {E_{B_1}}^{B_{(n/2+q+1)}} = \\
= {\eta_\Lambda}^{A_1 B_1} \big(\varepsilon_{(n/2+q)}\big)^{A_{(n/2+q)} A_{(n/2+q+1)}} \ldots \big(\varepsilon_1\big)_{A_1 A_2} \big(\varepsilon_2\big)^{A_3 A_2} \\
\big(\varepsilon_{(n/2+q)}\big)^{B_{(n/2+q)} B_{(n/2+q+1)}} \ldots \big(\varepsilon_1\big)_{B_1 B_2} \big(\varepsilon_2\big)^{B_3 B_2} = \\
= \big(-\eta_{\Lambda B_2 A_2} + (\eta_1)_\Lambda \varepsilon_{1 A_2 B_2}\big) \big(\varepsilon_{(n/2+q)}\big)^{A_{(n/2+q)} A_{(n/2+q+1)}} \ldots \big(\varepsilon_2\big)^{A_3 A_2} \\
\big(\varepsilon_{(n/2+q)}\big)^{B_{(n/2+q)} B_{(n/2+q+1)}} \ldots \big(\varepsilon_2\big)^{B_3 B_2} = \\
= -\big(-{\eta_\Lambda}^{A_3 B_3} + (\eta_2)_\Lambda {\varepsilon_2}^{A_3 B_3}\big) \big(\varepsilon_{(n/2+q)}\big)^{A_{(n/2+q)} A_{(n/2+q+1)}} \ldots \big(\varepsilon_{(n/2+q)}\big)^{B_{(n/2+q)} B_{(n/2+q+1)}} \ldots - \\
-(\eta_1)_\Lambda \big(\varepsilon_1\big)^{A_{(n/2+q+1)} B_{(n/2+q+1)}} = \\
= {\eta_\Lambda}^{A_3 B_3} \big(\varepsilon_{(n/2+q)}\big)^{A_{(n/2+q)} A_{(n/2+q+1)}} \ldots \big(\varepsilon_{(n/2+q)}\big)^{B_{(n/2+q)} B_{(n/2+q+1)}} \ldots - \\
- \sum_{Q=1}^{2} (\eta_Q)_\Lambda \big(\varepsilon_Q\big)^{A_{(n/2+q+1)} B_{(n/2+q+1)}} = \\
= {\eta_\Lambda}^{A_{(n/2+q+1)} B_{(n/2+q+1)}} - \sum_{Q=1}^{n/2+q} (\eta_Q)_\Lambda \big(\varepsilon_Q\big)^{A_{(n/2+q+1)} B_{(n/2+q+1)}} = \\
= -{\eta_\Lambda}^{B_{(n/2+q+1)} A_{(n/2+q+1)}}.
\end{gathered} \tag{16.3}$$

Случай II)., инволюция ${\tilde{E}_A}^B {\tilde{E}_B}^C = -{\delta_A}^C$.

$$\begin{gathered}
{\eta_\Lambda}^{A_1 B_1} {\tilde{E}_{A_1}}^{A_{(n/2+\tilde{q}+1)}} {\tilde{E}_{B_1}}^{B_{(n/2+\tilde{q}+1)}} = \\
= {\eta_\Lambda}^{A_1 B_1} \big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)^{A_{(n/2+\tilde{q}+1)} A_{(n/2+\tilde{q})}} \ldots \big(\varepsilon_{\tilde{1}}\big)_{A_1 A_2} \big(\varepsilon_{\tilde{2}}\big)^{A_3 A_2} \\
\big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)^{B_{(n/2+\tilde{q}+1)} B_{(n/2+\tilde{q})}} \ldots \big(\varepsilon_{\tilde{1}}\big)_{B_1 B_2} \big(\varepsilon_{\tilde{2}}\big)^{B_3 B_2} = \\
= \big(\eta_{\Lambda B_2 A_2} + (\eta_{\tilde{1}})_\Lambda \varepsilon_{\tilde{1} A_2 B_2}\big) \big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)^{A_{(n/2+\tilde{q}+1)} A_{(n/2+\tilde{q})}} \ldots \big(\varepsilon_{\tilde{2}}\big)^{A_3 A_2} \\
\big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)^{B_{(n/2+\tilde{q}+1)} B_{(n/2+\tilde{q})}} \ldots \big(\varepsilon_{\tilde{2}}\big)^{B_3 B_2} = \\
= \big({\eta_\Lambda}^{A_3 B_3} + (\eta_{\tilde{2}})_\Lambda {\varepsilon_{\tilde{2}}}^{B_3 A_3}\big) \big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)^{A_{(n/2+\tilde{q}+1)} A_{(n/2+\tilde{q})}} \ldots \big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)^{B_{(n/2+\tilde{q}+1)} B_{(n/2+\tilde{q})}} \ldots + \\
+(\eta_{\tilde{1}})_\Lambda \big(\varepsilon_{\tilde{1}}\big)_{A_{(n/2+\tilde{q}+1)} B_{(n/2+\tilde{q}+1)}} = \\
= {\eta_\Lambda}^{A_3 B_3} \big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)^{A_{(n/2+\tilde{q}+1)} A_{(n/2+\tilde{q})}} \ldots \big(\varepsilon_{\widetilde{(n/2+\tilde{q})}}\big)^{B_{(n/2+\tilde{q}+1)} B_{(n/2+\tilde{q})}} \ldots + \\
+ \sum_{\tilde{Q}=1}^{2} (\eta_{\tilde{Q}})_\Lambda \big(\varepsilon_{\tilde{Q}}\big)^{A_{(n/2+\tilde{q}+1)} B_{(n/2+\tilde{q}+1)}} = \\
= {\eta_\Lambda}^{A_{(n/2+\tilde{q}+1)} B_{(n/2+\tilde{q}+1)}} + \sum_{\tilde{Q}=1}^{n/2+\tilde{q}} (\eta_{\tilde{Q}})_\Lambda \big(\varepsilon_{\tilde{Q}}\big)^{B_{(n/2+\tilde{q}+1)} A_{(n/2+\tilde{q}+1)}} = \\
= {\eta_\Lambda}^{B_{(n/2+\tilde{q}+1)} A_{(n/2+\tilde{q}+1)}}.
\end{gathered} \tag{16.4}$$

16.2 Доказательство формул о связующих операторах

Необходимо доказать тождество (8.27)

$$\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}\eta^\Theta{}_{CB}=\frac{N}{4}(g_{\Lambda\Psi}g^{\Omega\Theta}+\delta_\Lambda{}^\Theta\delta_\Psi{}^\Omega-\delta_\Psi{}^\Theta\delta_\Lambda{}^\Omega). \tag{16.5}$$

1. Первый шаг.

$$\boxed{\boxed{\begin{aligned}&\eta_\Lambda{}^{AB}\eta_\Psi{}^{CD}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC}=\tfrac{N}{4}((\eta_Q)_\Psi(\eta_P)_\Lambda+(\eta_Q)_\Lambda(\eta_P)_\Psi),\quad P\neq Q.\\&\eta_{\Lambda AD}(\varepsilon_P)^{AB}(\varepsilon_Q)_{BC}\eta_\Psi{}^{CD}=\tfrac{N}{4}((\eta_Q)_\Psi(\eta_P)_\Lambda-(\eta_Q)_\Lambda(\eta_P)_\Psi),\quad P\neq Q.\end{aligned}}} \tag{16.6}$$

Доказательство.

$$\begin{gathered}\eta_\Lambda{}^{AB}\eta_\Psi{}^{CD}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC}=(\eta_\Lambda{}^{BA}+\sum_{\tilde P}(\varepsilon_{\tilde P})^{AB}(\eta_{\tilde P})_\Lambda)\eta_\Psi{}^{CD}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC}=\\=\eta_\Lambda{}^{BA}(\eta_\Psi{}^{DC}+\sum_{\tilde P}(\varepsilon_{\tilde P})^{CD}(\eta_{\tilde P})_\Psi)(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC}+\\+\sum_{\tilde P}(\varepsilon_{\tilde P})^{AB}(\eta_{\tilde P})_\Lambda\eta_\Psi{}^{CD}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC}=\eta_\Lambda{}^{BA}\eta_\Psi{}^{DC}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC}+\\+\sum_{\tilde P}((\varepsilon_{\tilde P})^{AB}(\eta_{\tilde P})_\Lambda\eta_\Psi{}^{CD}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC}+(\varepsilon_{\tilde P})^{CD}(\eta_{\tilde P})_\Psi\eta_\Lambda{}^{BA}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC}).\end{gathered} \tag{16.7}$$

(a) Первый шаг.

$$\begin{gathered}\eta_\Lambda{}^{BA}\eta_\Psi{}^{DC}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC}=\tfrac{N}{2}(\eta_P)_\Lambda(\eta_Q)_\Psi-(\varepsilon_P)^{BA}\eta_\Psi{}^{DC}\eta_{\Lambda AD}(\varepsilon_Q)_{BC}=\\=\tfrac{N}{2}(\eta_P)_\Lambda(\eta_Q)_\Psi+(\varepsilon_Q)^{BA}\eta_\Psi{}^{DC}\eta_{\Lambda AD}(\varepsilon_P)_{BC}=\\=\tfrac{N}{2}((\eta_P)_\Lambda(\eta_Q)_\Psi+(\eta_Q)_\Lambda(\eta_P)_\Psi)-\eta_\Lambda{}^{BA}\eta_\Psi{}^{DC}(\varepsilon_Q)_{AD}(\varepsilon_P)_{BC}.\end{gathered} \tag{16.8}$$

$$\boxed{\begin{gathered}\eta_\Lambda{}^{BA}\eta_\Psi{}^{DC}(\varepsilon_P)_{AD}(\varepsilon_Q)_{BC}=\\\tfrac{N}{2}((\eta_P)_\Lambda(\eta_Q)_\Psi+(\eta_Q)_\Lambda(\eta_P)_\Psi)-\eta_\Lambda{}^{BA}\eta_\Psi{}^{DC}(\varepsilon_Q)_{AD}(\varepsilon_P)_{BC}.\end{gathered}} \tag{16.9}$$

(b) Второй шаг.

$$\begin{gathered}(\varepsilon_{\tilde P})^{AD}(\varepsilon_P)_{AB}(\varepsilon_{\tilde Q})^{CB}(\varepsilon_Q)_{CD}\overset{\tilde P\neq\tilde Q}{=}-(\varepsilon_{\tilde Q})^{AD}(\varepsilon_P)_{AB}(\varepsilon_{\tilde P})^{CB}(\varepsilon_Q)_{CD}=\\-(\varepsilon_{\tilde Q})^{BC}(\varepsilon_P)_{BA}(\varepsilon_{\tilde P})^{DA}(\varepsilon_Q)_{DC}.\end{gathered} \tag{16.10}$$

$$\boxed{(\varepsilon_{\tilde P})^{AD}(\varepsilon_P)_{AB}(\varepsilon_{\tilde Q})^{CB}(\varepsilon_Q)_{CD}\overset{\tilde P\neq\tilde Q}{=}0.} \tag{16.11}$$

(c) Третий шаг.

$$\begin{gathered}\eta_\Lambda{}^{AD}(\varepsilon_P)_{AB}(\varepsilon_{\tilde P})^{CB}(\varepsilon_Q)_{CD}=-\eta_{\Lambda KL}(\varepsilon_{\tilde Q})^{KD}(\varepsilon_{\tilde Q})^{AL}(\varepsilon_P)_{AB}(\varepsilon_{\tilde P})^{CB}(\varepsilon_Q)_{CD}=\\=\eta_{\Lambda KL}(\varepsilon_{\tilde Q})^{KD}(\varepsilon_P)^{AL}(\varepsilon_{\tilde P})_{AB}(\varepsilon_Q)^{CB}(\varepsilon_{\tilde Q})_{CD}=\eta_{\Lambda KL}(\varepsilon_P)^{AL}(\varepsilon_{\tilde P})_{AB}(\varepsilon_Q)^{KB}=\\=-\eta_\Lambda{}^{AD}(\varepsilon_P)_{KD}(\varepsilon_{\tilde P})_{AB}(\varepsilon_Q)^{KB}=-\eta_\Lambda{}^{AD}(\varepsilon_Q)_{CD}(\varepsilon_P)_{AB}(\varepsilon_{\tilde P})^{CB}.\end{gathered} \tag{16.12}$$

$$\boxed{\eta_\Lambda{}^{AD}(\varepsilon_P)_{AB}(\varepsilon_{\tilde P})^{CB}(\varepsilon_Q)_{CD}=0.} \tag{16.13}$$

□

2. Второй шаг.

$$\boxed{\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}(\varepsilon_Q)_{BC}=\frac{N}{4}(g_{\Lambda\Psi}(\eta_Q)^\Omega+(\eta_Q)^\Lambda\delta_\Psi{}^\Omega-(\eta_Q)_\Psi\delta_\Lambda{}^\Omega).}\tag{16.14}$$

Доказательство.

$$\begin{gathered}
\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}(\varepsilon_Q)_{BC}= \\
=\eta_\Lambda{}^{AB}\eta_{\Psi AD}(-\eta^\Omega{}_{KL}(\varepsilon_P)^{KD}(\varepsilon_P)^{CL}+(\eta_P)^\Omega(\varepsilon_P)^{CD})(\varepsilon_Q)_{BC}= \\
=\eta_\Lambda{}^{AB}(-\eta_\Psi{}^{KM}(\varepsilon_P)_{MA}+(\eta_P)_\Psi\delta_A{}^K)(\varepsilon_Q)_{BC}(\varepsilon_P)^{CL}\eta^\Omega{}_{KL}+ \\
+(\eta_P)^\Omega\frac{N}{4}((\eta_P)_\Psi(\eta_Q)_\Lambda-(\eta_P)_\Lambda(\eta_Q)_\Psi)= \\
=(-\eta_{\Lambda NM}(\varepsilon_P)^{NB}+(\eta_P)_\Lambda\delta_M{}^B)\eta_\Psi{}^{KM}(\varepsilon_Q)_{BC}(\varepsilon_P)^{CL}\eta^\Omega{}_{KL}- \\
-\frac{N}{4}(\eta_P)_\Psi((\eta_Q)_\Lambda(\eta_P)^\Omega-(\eta_Q)^\Omega(\eta_P)_\Lambda)+\frac{N}{4}(\eta_P)^\Omega((\eta_P)_\Psi(\eta_Q)_\Lambda+(\eta_P)_\Lambda(\eta_Q)_\Psi)= \\
=\eta_{\Lambda NM}\eta_\Psi{}^{KM}\eta^\Omega{}_{KL}(\varepsilon_Q)^{LN}+\frac{N}{4}(\eta_P)_\Lambda(\underbrace{(\eta_Q)_\Psi(\eta_P)^\Omega}_{1}-\underbrace{(\eta_P)_\Psi(\eta_Q)^\Omega}_{2})- \\
-\frac{N}{4}(\eta_P)_\Psi(\underbrace{(\eta_Q)_\Lambda(\eta_P)^\Omega}_{3}-\underbrace{(\eta_Q)^\Omega(\eta_P)_\Lambda}_{2})+\frac{N}{4}(\eta_P)^\Omega(\underbrace{(\eta_P)_\Psi(\eta_Q)_\Lambda}_{3}-\underbrace{(\eta_P)_\Lambda(\eta_Q)_\Psi}_{1})= \\
=(-\eta_\Lambda{}^{AL}(\varepsilon_Q)_{AM}+(\eta_Q)_\Lambda\delta_M{}^L)\eta_\Psi{}^{KM}\eta^\Omega{}_{KL}= \\
=-\eta_\Lambda{}^{AL}(-\eta_{\Psi AD}(\varepsilon_Q)^{DK}+(\eta_Q)_\Psi\delta_A{}^K)\eta^\Omega{}_{KL}+\frac{N}{2}(\eta_Q)_\Lambda\delta_\Psi{}^\Omega= \\
=\eta_\Lambda{}^{AL}\eta_{\Psi AD}(-\eta^{\Omega CD}(\varepsilon_Q)_{CL}+(\eta_Q)^\Omega\delta_L{}^D)+\frac{N}{2}((\eta_Q)_\Lambda\delta_\Psi{}^\Omega-(\eta_Q)_\Psi\delta_\Lambda{}^\Omega)= \\
=-\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}(\varepsilon_Q)_{CB}+\frac{N}{2}(g_{\Lambda\Psi}(\eta_Q)^\Omega+(\eta_Q)_\Lambda\delta_\Psi{}^\Omega-(\eta_Q)_\Psi\delta_\Lambda{}^\Omega).
\end{gathered}\tag{16.15}$$

$$\boxed{\eta_\Lambda{}^{AB}\eta^\Omega{}_{CB}\eta_\Psi{}^{CD}(\varepsilon_Q)_{AD}=\frac{N}{4}(-g_{\Lambda\Psi}(\eta_Q)^\Omega+(\eta_Q)_\Lambda\delta_\Psi{}^\Omega+(\eta_Q)_\Psi\delta_\Lambda{}^\Omega).}\tag{16.16}$$

$\square$

3. Третий шаг.

$$\boxed{\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}\eta^\Theta{}_{CB}=\frac{N}{4}(g_{\Lambda\Psi}g^{\Omega\Theta}+\delta_\Lambda{}^\Theta\delta_\Psi{}^\Omega-\delta_\Psi{}^\Theta\delta_\Lambda{}^\Omega).}\tag{16.17}$$

Доказательство.

$$\begin{gathered}
\eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}\eta^\Theta{}_{CB} = \\
= \eta_\Lambda{}^{AB}\eta_{\Psi AD}\eta^{\Omega CD}(-\eta^{\Theta KL}(\varepsilon_Q)_{KB}(\varepsilon_Q)_{LC} + (\eta_Q)^\Theta(\varepsilon_Q)_{BC}) = \\
= -\eta_\Lambda{}^{AB}\eta_{\Psi AD}(-\eta^\Omega{}_{ML}(\varepsilon_Q)^{MD} + (\eta_Q)^\Omega\delta_L{}^D)(\varepsilon_Q)_{KB}\eta^{\Theta KL} + \\
+(\eta_Q)^\Theta\tfrac{N}{4}(g_{\Lambda\Psi}(\eta_Q)^\Omega + (\eta_Q)_\Lambda\delta_\Psi{}^\Omega - (\eta_Q)_\Psi\delta_\Lambda{}^\Omega) = \\
= \eta_\Lambda{}^{AB}(-\eta_\Psi{}^{MN}(\varepsilon_Q)_{NA} + (\eta_Q)_\Psi\delta_A{}^M)\eta^\Omega{}_{ML}\eta^{\Theta KL}(\varepsilon_Q)_{KB} + \\
+\tfrac{N}{4}(\eta_Q)^\Theta(g_{\Lambda\Psi}(\eta_Q)^\Omega + (\eta_Q)_\Lambda\delta_\Psi{}^\Omega - (\eta_Q)_\Psi\delta_\Lambda{}^\Omega) - \\
-\tfrac{N}{4}(\eta_Q)^\Omega(g_{\Lambda\Psi}(\eta_Q)^\Theta + (\eta_Q)_\Lambda\delta_\Psi{}^\Theta - (\eta_Q)_\Psi\delta_\Lambda{}^\Theta) = \\
= -(-\eta_{\Lambda SN}(\varepsilon_Q)^{SB} + (\eta_Q)^\Lambda\delta_N{}^B)\eta_\Psi{}^{MN}\eta^\Omega{}_{ML}\eta^{\Theta KL}(\varepsilon_Q)_{KB} + \tfrac{N}{4}(\eta_Q)^\Theta(g_{\Lambda\Psi}(\eta_Q)^\Omega + \\
+(\eta_Q)_\Lambda\delta_\Psi{}^\Omega - (\eta_Q)_\Psi\delta_\Lambda{}^\Omega) - \tfrac{N}{4}(\eta_Q)^\Omega(g_{\Lambda\Psi}(\eta_Q)^\Theta + (\eta_Q)_\Lambda\delta_\Psi{}^\Theta - (\eta_Q)_\Psi\delta_\Lambda{}^\Theta) + \\
+\tfrac{N}{4}(\eta_Q)_\Psi(\delta_\Lambda{}^\Omega(\eta_Q)^\Theta + (\eta_Q)_\Lambda g^{\Omega\Theta} - (\eta_Q)^\Omega\delta_\Lambda{}^\Theta) = \eta_{\Lambda SN}\eta_\Psi{}^{MN}\eta^\Omega{}_{ML}\eta^{\Theta SL} + \\
+\tfrac{N}{4}(\eta_Q)^\Theta(\underbrace{g_{\Lambda\Psi}(\eta_Q)^\Omega}_{1} + \underbrace{(\eta_Q)_\Lambda\delta_\Psi{}^\Omega}_{2} - \underbrace{(\eta_Q)_\Psi\delta_\Lambda{}^\Omega)}_{3} - \tfrac{N}{4}(\eta_Q)^\Omega(\underbrace{g_{\Lambda\Psi}(\eta_Q)^\Theta}_{1} + \underbrace{(\eta_Q)_\Lambda\delta_\Psi{}^\Theta}_{4} - \\
-\underbrace{(\eta_Q)_\Psi\delta_\Lambda{}^\Theta)}_{5} + \tfrac{N}{4}(\eta_Q)_\Psi(\underbrace{\delta_\Lambda{}^\Omega(\eta_Q)^\Theta}_{3} + \underbrace{(\eta_Q)_\Lambda g^{\Omega\Theta}}_{6} - \underbrace{(\eta_Q)^\Omega\delta_\Lambda{}^\Theta)}_{5} - \tfrac{N}{4}(\eta_Q)_\Lambda(\underbrace{\delta_\Psi{}^\Omega(\eta_Q)^\Theta}_{2} + \\
+\underbrace{(\eta_Q)_\Psi g^{\Omega\Theta}}_{6} - \underbrace{(\eta_Q)^\Omega\delta_\Psi{}^\Theta)}_{4} = \\
= -\eta_{\Psi AD}\eta^{\Omega CD}\eta^\Theta{}_{CB}\eta^{\Lambda AB} + \tfrac{N}{2}(g_{\Lambda\Psi}g^{\Omega\Theta} + \delta_\Lambda{}^\Theta\delta_\Psi{}^\Omega - \delta_\Psi{}^\Theta\delta_\Lambda{}^\Omega).
\end{gathered} \tag{16.18}$$

□

16.3 Доказательство формул об инфинитезимальных преобразованиях

Необходимо доказать тождество (10.5).

1. Случай 1).

$$\begin{gathered}
S_\Lambda{}^\Psi\eta_\Psi{}^{AB} = -\eta_\Lambda{}^{CD}S_C{}^A E_D{}^K S_K{}^M E_M{}^B, \\
T_\Lambda{}^\Psi\eta_\Psi{}^{AB} = \eta_\Lambda{}^{CB}T_C{}^A - \eta_\Lambda{}^{AD}E_D{}^K T_K{}^M E_M{}^B, \\
(\eta_I)^\Lambda T_\Lambda{}^\Psi\eta_\Psi{}^{AB} = (\varepsilon_I)^{CB}T_C{}^A - (\varepsilon_I)^{AD}E_D{}^K T_K{}^M E_M{}^B, \\
E_D{}^K T_K{}^M E_M{}^B = (\varepsilon_I)^{CB}T_C{}^A(\varepsilon_I)_{AD} - (\eta_I)^\Lambda T_\Lambda{}^\Psi\eta_\Psi{}^{AB}(\varepsilon_I)_{AD}, \\
T_\Lambda{}^\Psi\eta_\Psi{}^{AB} = \eta_\Lambda{}^{CB}T_C{}^A - \eta_\Lambda{}^{AD}(\varepsilon_I)^{KB}T_K{}^M(\varepsilon_I)_{MD} + \eta_\Lambda{}^{AD}(\varepsilon_I)_{MD}(\eta_I)^\Omega T_\Omega{}^\Phi\eta_\Phi{}^{MB}, \\
T_C{}^A = -E_C{}^B T_B{}^D E_D{}^A.
\end{gathered} \tag{16.19}$$

2. Случай 2).

$$\begin{gathered}
S_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \eta_\Lambda{}^{CD} S_C{}^A (\eta_I)^\Omega S_\Omega{}^\Phi \eta_\Phi{}^{KB} \varepsilon_{KL} S_X{}^L \varepsilon^{MX} (\varepsilon_I)_{MD}, \\
T_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \eta_\Lambda{}^{CB} T_C{}^A + \eta_\Lambda{}^{AD} (\eta_I)^\Omega T_\Omega{}^\Phi \eta_\Psi{}^{KB} (\varepsilon_I)_{KD} + \eta_\Lambda{}^{AD} (\varepsilon_I)^{KB} \varepsilon_{KL} T_X{}^L \varepsilon^{MX} (\varepsilon_I)_{MD}, \\
T_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \eta_\Lambda{}^{CB} T_C{}^A - \eta_\Lambda{}^{AD} (\varepsilon_I)^{KB} T_K{}^M (\varepsilon_I)_{MD} + \eta_\Lambda{}^{AD} (\varepsilon_I)_{MD} (\eta_I)^\Omega T_\Omega{}^\Phi \eta_\Phi{}^{MB}, \\
T_A{}^K = -T_C{}^D \varepsilon_{AD} \varepsilon^{KC}.
\end{gathered} \tag{16.20}$$

Подставим в это уравнение $T_C{}^A = \frac{1}{2} T^{\Theta\Phi} \eta_\Phi{}^{AB} \eta_{\Theta CB}$ и получим (16.21)

$$\begin{gathered}
T_\Lambda{}^\Psi \eta_\Psi{}^{AB} = \tfrac{1}{2}(\eta_\Lambda{}^{CB} T^{\Theta\Phi} \eta_\Phi{}^{AD} \eta_{\Theta CD} - \eta_\Lambda{}^{AD} (\varepsilon_I)^{KB} T^{\Theta\Phi} \eta_\Phi{}^{MN} \eta_{\Theta KN} (\varepsilon_I)^{MD}) + \\
+ \eta_\Lambda{}^{AD} (\varepsilon_I)_{MD} (\eta_I)^\Omega T_\Omega{}^\Phi \eta_\Phi{}^{MB} = \tfrac{1}{2}(T_\Lambda{}^\Phi \eta_\Phi{}^{AB} - \eta_{\Lambda CD} T^{\Theta\Phi} \eta_\Phi{}^{AD} \eta_\Theta{}^{CB} - \\
- \eta_\Lambda{}^{AD} (\eta_I)_\Theta T^{\Theta\Phi} \eta_\Phi{}^{KB} (\varepsilon_I)_{MD} + \eta_\Lambda{}^{AD} T^{\Theta\Phi} \eta_\Theta{}^{LB} (\varepsilon_I)_{LN} \eta_\Phi{}^{MN} (\varepsilon_I)_{MD}) + \\
+ \eta_\Lambda{}^{AD} (\varepsilon_I)_{MD} (\eta_I)^\Omega T_\Omega{}^\Phi \eta_\Phi{}^{MB} = \tfrac{1}{2}(T_\Lambda{}^\Phi \eta_\Phi{}^{AB} - T^\Theta{}_\Lambda \eta_\Theta{}^{AB} + T^{\Theta\Phi} \eta_{\Phi CD} \eta_\Lambda{}^{AD} \eta_\Theta{}^{CB} - \\
- \eta_\Lambda{}^{AB} (\eta_I)_\Theta T^{\Theta\Phi} \eta_\Phi{}^{MB} (\varepsilon_I)_{MD} - \eta_\Lambda{}^{AD} T^{\Theta\Phi} \eta_{\Phi LD} \eta_\Theta{}^{LB} + \\
+ \eta_\Lambda{}^{AD} T^{\Theta\Phi} \eta_\Theta{}^{LB} (\eta_I)_\Phi (\varepsilon_I)_{LD}) + \eta_\Lambda{}^{AB} (\varepsilon_I)_{MD} (\eta_I)^\Omega T_\Omega{}^\Phi \eta_\Phi{}^{MB} = T_\Lambda{}^\Psi \eta_\Psi{}^{AB}.
\end{gathered} \tag{16.22}$$

16.4 Доказательство тождества о скобке Ли

Необходимо доказать тождество (11.70).

$$\begin{gathered}
(L_\pm)_{[x,y]} (Y_K)^A = (x^\Theta \partial_\Theta y^\Omega - y^\Theta \partial_\Theta x^\Omega) \partial_\Omega (Y_K)^A - \sum_J^{2N} (Y_J)^B (P_J)^\Lambda{}_B (P_K^*)_\Theta{}^A \partial_\Lambda (x^\Xi \partial_\Xi y^\Theta - y^\Xi \partial_\Xi x^\Theta) = \\
= (x^\Theta \partial_\Theta y^\Omega - y^\Theta \partial_\Theta x^\Omega) \partial_\Omega (Y_K)^A - \sum_J^{2N} (Y_J)^B (P_J)^\Lambda{}_B (P_K^*)_\Theta{}^A (\partial_\Lambda x^\Xi \partial_\Xi y^\Theta - \partial_\Lambda y^\Xi \partial_\Xi x^\Theta) - \\
- \sum_J^{2N} (Y_J)^B (P_J)^\Lambda{}_B (P_K^*)_\Theta{}^A (x^\Xi \partial_\Lambda \partial_\Xi y^\Theta - y^\Xi \partial_\Lambda \partial_\Xi x^\Theta), \\
((L_\pm)_x (L_\pm)_y - (L_\pm)_y (L_\pm)_x)(Y)^A = x^\Phi \partial_\Phi (y^\Omega \partial_\Omega (Y_K)^A - \sum_J^{2N} (Y_J)^B (P_J)^\Lambda{}_B (P_K^*)_\Theta{}^A \partial_\Lambda y^\Theta) - \\
- (y^\Omega \partial_\Omega (Y_K)^C - \sum_{I,J}^{2N} (Y_J)^B (P_J)^\Lambda{}_B (P_I^*)_\Theta{}^C \partial_\Lambda y^\Theta)(P_I)^\Phi{}_C (P_K^*)_\Xi{}^A \partial_\Phi x^\Xi - y^\Phi \partial_\Phi (x^\Omega \partial_\Omega (Y_K)^A - \\
- \sum_J^{2N} (Y_J)^B (P_J)^\Lambda{}_B (P_K^*)_\Theta{}^A \partial_\Lambda x^\Theta) + (x^\Omega \partial_\Omega (Y_K)^C - \sum_{I,J}^{2N} (Y_J)^B (P_J)^\Lambda{}_B (P_I^*)_\Theta{}^C \partial_\Lambda x^\Theta)(P_I)^\Phi{}_C (P_K^*)_\Xi{}^A \partial_\Phi y^\Xi = \\
= x^\Phi \partial_\Phi y^\Omega \partial_\Omega (Y_K)^A + x^\Phi y^\Omega \partial_\Phi \partial_\Omega (Y_K)^A - x^\Phi \sum_J^{2N} \partial_\Phi (Y_J)^B (P_J)^\Lambda{}_B (P_K^*)_\Theta{}^A \partial_\Lambda y^\Theta - \\
- x^\Theta \sum_J^{2N} (Y_J)^B (P_J)^\Lambda{}_B (P_K^*)_\Theta{}^A \partial_\Phi \partial_\Lambda y^\Theta - y^\Omega \sum_J^{2N} \partial_\Omega (Y_J)^C (P_J)^\Phi{}_C (P_K^*)_\Xi{}^A \partial_\Phi x^\Xi + \\
+ \sum_{I,J}^{2N} (Y_J)^B (P_J)^\Lambda{}_B (P_I^*)_\Theta{}^C \partial_\Lambda y^\Theta (P_I)^\Phi{}_C (P_K^*)_\Xi{}^A \partial_\Phi x^\Xi - \\
- y^\Phi \partial_\Phi x^\Omega \partial_\Omega (Y_K)^A - y^\Phi x^\Omega \partial_\Phi \partial_\Omega (Y_K)^A + y^\Phi \sum_J^{2N} \partial_\Phi (Y_J)^B (P_J)^\Lambda{}_B (P_K^*)_\Theta{}^A \partial_\Lambda x^\Theta + \\
+ y^\Phi \sum_J^{2N} (Y_J)^B (P_J)^\Lambda{}_B (P_K^*)_\Theta{}^A \partial_\Phi \partial_\Lambda x^\Theta + x^\Omega \sum_J^{2N} \partial_\Omega (Y_J)^C (P_J)^\Phi{}_C (P_K^*)_\Xi{}^A \partial_\Phi y^\Xi - \\
- \sum_{I,J}^{2N} (Y_J)^B (P_J)^\Lambda{}_B (P_I^*)_\Theta{}^C \partial_\Lambda x^\Theta (P_I)^\Phi{}_C (P_K^*)_\Xi{}^A \partial_\Phi y^\Xi.
\end{gathered} \tag{16.23}$$

16.5 Доказательство формул о тензоре кривизны

Необходимо доказать тождество (12.7).

Тензор кривизны в спинорном расслоении вычисляется по формуле

$$(\mathcal{R}_K)_{\Lambda\Psi C}{}^{A} = 2(\partial_{[\Lambda}(\Gamma_K)_{\Psi]C}{}^{A} + \Gamma_{[\Lambda|L|}{}^{A}(\Gamma_K)_{\Psi]C}{}^{L}). \tag{16.24}$$

Тогда $\partial_\Lambda(\varepsilon_K)_{AB}{}^{CD} = 0$ при выполнении условия (11.33) означает

$$\partial_{[\Lambda}(\Gamma_K)_{\Psi]A}{}^{C} = \underbrace{-\frac{2}{n-4}((\varepsilon_K)_{AB}{}^{CD} - \frac{n-2}{N}\delta_A{}^{C}\delta_B{}^{D})}_{:=(\tilde{\varepsilon}_K)_{AB}{}^{CD}}\partial_{[\Lambda}(\Gamma_K)_{\Psi]D}{}^{B}. \tag{16.25}$$

В свою очередь, $\nabla_\Lambda(\varepsilon_K)_{AB}{}^{CD} = 0$ при выполнении условия (11.33) означает

$$\begin{gathered}
(\Gamma_K)_{\Lambda A}{}^{L}(\tilde{\varepsilon}_K)_{LB}{}^{CD} + (\Gamma_K)_{\Lambda B}{}^{L}(\tilde{\varepsilon}_K)_{AL}{}^{CD} = (\Gamma_K)_{\Lambda L}{}^{C}(\tilde{\varepsilon}_K)_{AB}{}^{LD} + (\Gamma_K)_{\Lambda L}{}^{D}(\tilde{\varepsilon}_K)_{AB}{}^{CL},\\
(\Gamma_K)_{\Lambda A}{}^{L}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{LB}{}^{CD} + (\Gamma_K)_{\Lambda B}{}^{L}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{AL}{}^{CD} =\\
= (\Gamma_K)_{\Lambda L}{}^{C}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{AB}{}^{LD} + (\Gamma_K)_{\Lambda L}{}^{D}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{AB}{}^{CL},\\
(\Gamma_K)_{\Lambda A}{}^{L}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{LB}{}^{CD} + (\Gamma_K)_{\Lambda B}{}^{L}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{AL}{}^{CD} =\\
= (\Gamma_K)_{\Lambda C}{}^{A}(\Gamma_K)_{\Psi A}{}^{L}(\tilde{\varepsilon}_K)_{LB}{}^{CD} + (\Gamma_K)_{\Lambda L}{}^{D}(\Gamma_K)_{\Psi C}{}^{A}(\tilde{\varepsilon}_K)_{AB}{}^{CL},\\
(\Gamma_K)_{[\Lambda|A|}{}^{L}(\Gamma_K)_{\Psi]C}{}^{A}(\tilde{\varepsilon}_K)_{LB}{}^{CD} = (\Gamma_K)_{[\Lambda|L|}{}^{D}(\Gamma_K)_{\Psi]B}{}^{L}.
\end{gathered} \tag{16.26}$$

Откуда

$$(\mathcal{R}_K)_{\Lambda\Psi C}{}^{M} = (\mathcal{R}_K)_{\Lambda\Psi L}{}^{D}\tilde{\varepsilon}_{CD}{}^{ML}. \tag{16.27}$$

16.6 Доказательство вспомогательных тождеств для теоремы о спинорных аналогах тензора кривизны

Необходимо доказать тождество (12.9).

$$\boxed{\boxed{\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_2} = \tfrac{N}{2}g_{\Lambda_1[\Lambda_4}g_{\Lambda_3]\Lambda_2}.}} \tag{16.28}$$

Доказательство.

$$\begin{gathered}
\eta_{\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2A_1A_3} = (-\eta_{\Lambda_1B_1B_2}(\varepsilon_I)^{A_1B_2}(\varepsilon_I)^{B_1A_2} + (\eta_I)_{\Lambda_1}(\varepsilon_I)^{A_1A_2})\eta_{\Lambda_2A_1A_3} =\\
= -\eta_{\Lambda_1B_1B_2}(\varepsilon_I)^{B_1A_2}(-\eta_{\Lambda_2}{}^{A_1B_2}(\varepsilon_I)_{A_1A_3} + (\eta_I)_{\Lambda_2}\delta_{A_3}{}^{B_2}) + (\eta_I)_{\Lambda_1}(\varepsilon_I)^{A_1A_2}\eta_{\Lambda_2A_1A_3} =\\
= \eta_{\Lambda_1B_1B_2}\eta_{\Lambda_2}{}^{A_1B_2}(\varepsilon_I)^{B_1A_2}(\varepsilon_I)_{A_1A_3} - (\eta_I)_{\Lambda_2}(\varepsilon_I)^{B_1A_2}\eta_{\Lambda_1B_1A_3} + (\eta_I)_{\Lambda_1}(\varepsilon_I)^{A_1A_2}\eta_{\Lambda_2A_1A_3}.
\end{gathered} \tag{16.29}$$

Альтернируем по Λ_1, Λ_2 и суммируем по I

$$\begin{gathered}
\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3} = \tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2}{}_{A_1A_3}, \\
\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_2} = \\
= \eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\big(\tfrac{2}{n-4}\varepsilon^{B_1A_2}{}_{A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_2}\big) = \\
= \eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1B_3|}\eta_{\Lambda_4]}{}^{B_1B_3} = \\
= -\eta_{[\Lambda_1}{}^{A_1B_2}\eta_{\Lambda_2]A_1B_3}\eta_{[\Lambda_3}{}^{B_1B_3}\eta_{\Lambda_4]B_1B_2} + \tfrac{N}{2}\big(g_{\Lambda_1[\Lambda_4}g_{\Lambda_3]\Lambda_2} - g_{\Lambda_1[\Lambda_3}g_{\Lambda_4]\Lambda_2}\big),
\end{gathered} \tag{16.30}$$

откуда и следует требуемое. □

Необходимо доказать тождество (12.10).

$$\boxed{\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} = Ng_{[\Lambda_3|[\Lambda_2}g_{\Lambda_1]|[\Lambda_6}g_{\Lambda_5]|\Lambda_4]}\,.} \tag{16.31}$$

Доказательство.

$$\begin{gathered}
\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} = \\
= \tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2}{}_{A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} = \\
= -\tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2A_4A_3}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} + \\
+ \tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3}{}^{B_1A_2}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} = \\
= \tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2}{}_{A_4A_5}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} + \\
+ \tfrac{4}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3}{}^{B_1A_2}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} = \\
= \eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4B_3|}\eta_{\Lambda_6]}{}^{B_1B_3} - \\
- \tfrac{4}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_5|}\eta_{\Lambda_4]}{}^{A_6A_5}\eta_{[\Lambda_5|A_6A_2|}\eta_{\Lambda_6]}{}^{B_1A_2} - \\
- \tfrac{8}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_5|}g_{\Lambda_4][\Lambda_6}\eta_{\Lambda_5]}{}^{B_1A_5} = \\
\left|\begin{array}{c}
\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4B_3|}\eta_{\Lambda_6]}{}^{B_1B_3} = \\
= 2\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4B_3|}g_{\Lambda_6][\Lambda_1}\eta_{\Lambda_2]}{}^{A_1B_3} + \\
+ \eta_{[\Lambda_1|A_4B_3|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5}{}^{B_1B_3}\eta_{\Lambda_6]B_1B_2} = \\
= 2\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4B_3|}g_{\Lambda_6][\Lambda_1}\eta_{\Lambda_2]}{}^{A_1B_3} + \\
+ 2\eta_{[\ \Lambda_2}{}^{A_1B_2}g_{\Lambda_1][\Lambda_4}\eta_{\Lambda_3]A_1A_3}\eta_{[\Lambda_5}{}^{B_1A_3}\eta_{\Lambda_6]B_1B_2} + \\
+ \eta_{[\Lambda_1|A_1A_3|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4B_3}\eta_{[\Lambda_5}{}^{B_1B_3}\eta_{\Lambda_6]B_1B_2}
\end{array}\right| \\
= \big(-1 + \tfrac{4}{n-4}\big)\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_2} + \\
+ \big(\tfrac{8}{n-4} - 4\big)\tfrac{N}{2}g_{[\Lambda_3|[\Lambda_2}g_{\Lambda_1]|[\Lambda_5}g_{\Lambda_6]|\Lambda_4]}\,.
\end{gathered} \tag{16.32}$$

откуда и следует требуемое. □

Необходимо доказать тождество (12.11).

$$\boxed{\begin{aligned}&\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2}=\\&=\tfrac{N}{4}(g_{\Lambda_1[\Lambda_8}g_{\Lambda_7]\Lambda_2}g_{\Lambda_3[\Lambda_6}g_{\Lambda_5]\Lambda_4}-g_{\Lambda_1[\Lambda_6}g_{\Lambda_5]\Lambda_2}g_{\Lambda_4[\Lambda_8}g_{\Lambda_7]\Lambda_3}+\\&+g_{\Lambda_1[\Lambda_4}g_{\Lambda_3]\Lambda_2}g_{\Lambda_5[\Lambda_8}g_{\Lambda_7]\Lambda_6})+N(g_{[\Lambda_3|[\Lambda_2}g_{\Lambda_1][\Lambda_8}g_{\Lambda_7][\Lambda_6}g_{\Lambda_5]|\Lambda_4]}+\\&+g_{[\Lambda_5|[\Lambda_4}g_{\Lambda_3][\Lambda_7}g_{\Lambda_8][\Lambda_2}g_{\Lambda_1]|\Lambda_6]}+g_{[\Lambda_3|[\Lambda_7}g_{\Lambda_8][\Lambda_5}g_{\Lambda_6][\Lambda_1}g_{\Lambda_2]|\Lambda_4]}).\end{aligned}} \tag{16.33}$$

Доказательство.

1. Первый шаг.

$$\begin{aligned}&\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2}=\\
&=\tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2}{}_{A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2}=\\
&=-\tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2A_4A_3}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2}+\\
&+\tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3}{}^{B_1A_2}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2}=\\
&=\tfrac{2}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\varepsilon^{B_1A_2}{}_{A_4A_5}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2}+\\
&+\tfrac{4}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3}{}^{B_1A_2}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2}=\\
&=\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4A_5|}\eta_{\Lambda_6]}{}^{A_6A_5}\eta_{[\Lambda_7|A_6A_7|}\eta_{\Lambda_8]}{}^{B_1A_7}+\\
&+\tfrac{4}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3}{}^{B_1A_2}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2}+\\
&+\tfrac{4}{n-4}\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5}{}^{B_1A_2}\eta_{\Lambda_6]A_4A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2}=\\
&=(1-\tfrac{4}{n-4})\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4A_5|}\eta_{\Lambda_6]}{}^{A_6A_5}\eta_{[\Lambda_7|A_6A_7|}\eta_{\Lambda_8]}{}^{B_1A_7}+\\
&+\tfrac{4}{n-4}\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2}+\\
&+\tfrac{8}{n-4}\tfrac{N}{2}(\tfrac{1}{2}g_{\Lambda_1[\Lambda_3}g_{\Lambda_4]\Lambda_2}g_{\Lambda_5[\Lambda_8}g_{\Lambda_7]\Lambda_6}+g_{[\Lambda_3|[\Lambda_1}g_{\Lambda_2][\Lambda_8}g_{\Lambda_7][\Lambda_6}g_{\Lambda_5]|\Lambda_4]}-\\
&-g_{[\Lambda_3|[\Lambda_1}g_{\Lambda_2][\Lambda_5}g_{\Lambda_6][\Lambda_7}g_{\Lambda_8]|\Lambda_4]}+\tfrac{1}{2}g_{\Lambda_5[\Lambda_8}g_{\Lambda_7]\Lambda_6}g_{\Lambda_1[\Lambda_4}g_{\Lambda_3]\Lambda_2}+\\
&+g_{[\Lambda_5|[\Lambda_8}g_{\Lambda_7][\Lambda_1}g_{\Lambda_2][\Lambda_3}g_{\Lambda_4]|\Lambda_6]}-g_{[\Lambda_5|[\Lambda_8}g_{\Lambda_7][\Lambda_4}g_{\Lambda_3][\Lambda_2}g_{\Lambda_1]|\Lambda_6]}),\end{aligned} \tag{16.34}$$

$$\begin{aligned}&\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_1A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2}=\\
&=\eta_{[\Lambda_1|B_1B_2|}\eta_{\Lambda_2]}{}^{A_1B_2}\eta_{[\Lambda_3|A_1A_3|}\eta_{\Lambda_4]}{}^{A_4A_3}\eta_{[\Lambda_5|A_4A_5|}\eta_{\Lambda_6]}{}^{A_6A_5}\eta_{[\Lambda_7|A_6A_7|}\eta_{\Lambda_8]}{}^{B_1A_7}\end{aligned} \tag{16.35}$$

2. Второй шаг.

$$\begin{gathered}
\eta_{[\Lambda_1}{}^{A_1A_2}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_7}{}^{A_8A_7}\eta_{\Lambda_8]A_8A_2} = \\
= \eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_1}{}^{A_8A_7}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_7|A_8A_2|}\eta_{\Lambda_8]}{}^{A_1A_2}+ \\
+2\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_2|A_1A_3|}g_{\Lambda_1][\Lambda_8}\eta_{\Lambda_7]}{}^{A_1A_7} = \\
= \eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_1}{}^{A_6A_5}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_5|A_6A_7|}\eta_{\Lambda_6]}{}^{A_8A_7}\eta_{[\Lambda_7|A_8A_2|}\eta_{\Lambda_8]}{}^{A_1A_2}+ \\
+2(\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_2|A_1A_3|}g_{\Lambda_1][\Lambda_8}\eta_{\Lambda_7]}{}^{A_1A_7}+ \\
+\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_2|A_1A_3|}g_{\Lambda_1][\Lambda_6}\eta_{\Lambda_5]}{}^{A_6A_5}\eta_{[\Lambda_7|A_6A_2|}\eta_{\Lambda_8]}{}^{A_1A_2}) = \\
= -\eta_{[\Lambda_1|A_1A_3|}\eta_{\Lambda_2]}{}^{A_4A_3}\eta_{[\Lambda_3|A_4A_5|}\eta_{\Lambda_4]}{}^{A_6A_5}\eta_{[\Lambda_5|A_6A_7|}\eta_{\Lambda_6]}{}^{A_8A_7}\eta_{[\Lambda_7|A_8A_2|}\eta_{\Lambda_8]}{}^{A_1A_2}+ \\
+2(\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_5}{}^{A_6A_5}\eta_{\Lambda_6]A_6A_7}\eta_{[\Lambda_2|A_1A_3|}g_{\Lambda_1][\Lambda_8}\eta_{\Lambda_7]}{}^{A_1A_7}+ \\
+\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_2|A_6A_2|}g_{\Lambda_1][\Lambda_6}\eta_{\Lambda_5]}{}^{A_6A_5}\eta_{[\Lambda_8|A_1A_3|}\eta_{\Lambda_7]}{}^{A_1A_2}+ \\
+2\eta_{[\Lambda_3}{}^{A_4A_3}\eta_{\Lambda_4]A_4A_5}\eta_{[\Lambda_7|A_6A_3|}g_{\Lambda_8][\Lambda_2}g_{\Lambda_1][\Lambda_6}\eta_{\Lambda_5]}{}^{A_6A_5}+ \\
+\eta_{[\Lambda_3}{}^{A_4A_3}g_{\Lambda_4][\Lambda_1}\eta_{\Lambda_2]A_1A_3}\eta_{[\Lambda_5|A_4A_7|}\eta_{\Lambda_6]}{}^{A_8A_7}\eta_{[\Lambda_7|A_8A_2|}\eta_{\Lambda_8]}{}^{A_1A_2}) = \\
= 2(\tfrac{N}{4}(g_{\Lambda_1[\Lambda_8}g_{\Lambda_7]\Lambda_2}g_{\Lambda_3[\Lambda_6}g_{\Lambda_5]\Lambda_4} - g_{\Lambda_1[\Lambda_6}g_{\Lambda_5]\Lambda_2}g_{\Lambda_4[\Lambda_8}g_{\Lambda_7]\Lambda_3}+ \\
+g_{\Lambda_1[\Lambda_4}g_{\Lambda_3]\Lambda_2}g_{\Lambda_5[\Lambda_8}g_{\Lambda_7]\Lambda_6}) + \tfrac{N}{2}(g_{[\Lambda_3|[\Lambda_2}g_{\Lambda_1][\Lambda_8}g_{\Lambda_7][\Lambda_6}g_{\Lambda_5]|\Lambda_4]}- \\
-g_{[\Lambda_3|[\Lambda_7}g_{\Lambda_8][\Lambda_1}g_{\Lambda_2][\Lambda_6}g_{\Lambda_5]|\Lambda_4]} + 2g_{[\Lambda_5|[\Lambda_4}g_{\Lambda_3][\Lambda_7}g_{\Lambda_8][\Lambda_2}g_{\Lambda_1]|\Lambda_6]}+ \\
+g_{[\Lambda_2|[\Lambda_8}g_{\Lambda_7][\Lambda_6}g_{\Lambda_5][\Lambda_3}g_{\Lambda_4]|\Lambda_1]} - g_{[\Lambda_3|[\Lambda_8}g_{\Lambda_7][\Lambda_6}g_{\Lambda_5][\Lambda_2}g_{\Lambda_1]|\Lambda_4]}- \\
-g_{[\Lambda_3|[\Lambda_7}g_{\Lambda_8][\Lambda_2}g_{\Lambda_1][\Lambda_6}g_{\Lambda_5]|\Lambda_4]} + g_{[\Lambda_3|[\Lambda_7}g_{\Lambda_8][\Lambda_5}g_{\Lambda_6][\Lambda_1}g_{\Lambda_2]|\Lambda_4]}))- \\
-\eta_{[\Lambda_1|A_1A_3|}\eta_{\Lambda_2]}{}^{A_4A_3}\eta_{[\Lambda_3|A_4A_5|}\eta_{\Lambda_4]}{}^{A_6A_5}\eta_{[\Lambda_5|A_6A_7|}\eta_{\Lambda_6]}{}^{A_8A_7}\eta_{[\Lambda_7|A_8A_2|}\eta_{\Lambda_8]}{}^{A_1A_2},
\end{gathered} \tag{16.36}$$

откуда и следует требуемое. □

Замечание 16.1. *Очевидно, что данные выкладки не применимы для* $n<8$*, поскольку* $\frac{8}{n-4}-4$ *не будет иметь смысла в этом случае.*

16.7 Доказательство формул о твисторном уравнении.

Необходимо доказать (13.4).

$$\begin{gathered}
\nabla_\Psi X^C = \tfrac{2}{n}\eta_\Psi{}^{CD}\nabla_{AD}X^A, \\
\nabla_\Phi\nabla_\Psi X^C = \tfrac{2}{n}\eta_\Psi{}^{CD}\nabla_\Phi\nabla_{AD}X^A = \tfrac{2}{n}(\eta_\Psi{}^{CD}\nabla_{AD}\nabla_\Phi X^A - \eta_\Psi{}^{CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^A X^K) = \\
= (\tfrac{2}{n})^2\eta_\Psi{}^{CD}\nabla_{AD}\eta_\Phi{}^{AK}\nabla_{LK}X^L - \tfrac{2}{n}\eta_\Psi{}^{CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^A X^K = \\
= (\tfrac{2}{n})^2(-\eta_{\Psi AD}\nabla^{CD}\eta_\Phi{}^{AK}\nabla_{LK}X^L + \nabla_\Psi\eta_\Phi{}^{CK}\nabla_{LK}X^L) - \tfrac{2}{n}\eta_\Psi{}^{CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^A X^K = \\
= -(\tfrac{2}{n})^2\eta_{\Psi AD}\eta_\Phi{}^{AK}\nabla^{CD}\nabla_{LK}X^L + \tfrac{2}{n}\nabla_\Psi\nabla_\Phi X^C - \tfrac{2}{n}\eta_\Psi{}^{CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^A X^K.
\end{gathered} \tag{16.37}$$

$$\begin{gathered}
(1-\tfrac{2}{n})\nabla_\Phi\nabla_\Psi X^C+\tfrac{2}{n}R_{\Phi\Psi K}{}^C X^K+\tfrac{2}{n}\eta_\Psi{}^{CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^A X^K=\\
=-(\tfrac{2}{n})^2\eta_{\Psi AD}\eta_\Phi{}^{AK}\nabla^{CD}\nabla_{LK}X^L,\\
(1-\tfrac{2}{n})\nabla_{(\Phi}\nabla_{\Psi)}X^C+\tfrac{2}{n}\eta_{(\Psi}{}^{CD}\eta^\Omega{}_{|AD}R_{\Omega|\Phi)K}{}^A X^K=\\
=-(\tfrac{2}{n})^2\tfrac{1}{2}g_{\Psi\Phi}\nabla^{CK}\nabla_{LK}X^L=-\tfrac{1}{n}g_{\Psi\Phi}\nabla^\Omega\nabla_\Omega X^C\\
(1-\tfrac{2}{n})\nabla^\Theta\nabla_\Theta X^C+\tfrac{2}{n}\eta^{\Phi CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^A X^K=-\nabla^\Theta\nabla_\Theta X^C,\\
\nabla^\Theta\nabla_\Theta X^C=-\tfrac{1}{n-1}\eta^{\Phi CD}\eta^\Omega{}_{AD}R_{\Omega\Phi K}{}^A X^K.
\end{gathered}\tag{16.38}$$

$$\begin{gathered}
(1-\tfrac{2}{n})\nabla_{(\Phi}\nabla_{\Psi)}X^C+\tfrac{2}{n}\eta_{(\Psi}{}^{CD}\eta^\Omega{}_{|AD}R_{\Omega|\Phi)K}{}^A X^K=\tfrac{1}{n}g_{\Psi\Phi}\tfrac{1}{n-1}\eta^{\Theta CD}\eta^\Omega{}_{AD}R_{\Omega\Theta K}{}^A X^K\\
\nabla_{(\Phi}\nabla_{\Psi)}X^C=\tfrac{1}{(n-1)(n-2)}g_{\Psi\Phi}\eta^{\Theta CD}\eta^\Omega{}_{AD}R_{\Omega\Theta K}{}^A X^K-\tfrac{2}{n-2}\eta_{(\Psi}{}^{CD}\eta^\Omega{}_{|AD}R_{\Omega|\Phi)K}{}^A X^K,\\
2\nabla_\Phi\nabla_\Psi X^C=\tfrac{2}{n}\eta_\Psi{}^{CD}\eta^\Gamma{}_{AD}(R_{\Phi\Gamma K}{}^A X^K+\tfrac{2}{(n-1)(n-2)}g_{\Phi\Gamma}\eta^{\Theta AL}\eta^\Omega{}_{SL}R_{\Omega\Theta K}{}^S X^K-\\
-\tfrac{2}{n-2}\eta_\Phi{}^{AL}\eta^\Omega{}_{SL}R_{\Omega\Gamma K}{}^S X^K-\tfrac{2}{n-2}\eta_\Gamma{}^{AL}\eta^\Omega{}_{SL}R_{\Omega\Phi K}{}^S X^K)=\\
=\tfrac{2}{n}((-1-\tfrac{n}{n-2})\eta_\Psi{}^{CD}\eta^\Omega{}_{SD}R_{\Omega\Phi K}{}^S X^K+\tfrac{2}{(n-1)(n-2)}\eta_\Psi{}^{CD}\eta_{\Phi AD}\eta^{\Theta AL}\eta^\Omega{}_{SL}R_{\Omega\Theta K}{}^S X^K-\\
-\tfrac{2}{n-2}\eta_\Psi{}^{CD}\eta_\Phi{}^{AL}\eta^\Omega{}_{SL}\eta^\Theta{}_{AD}R_{\Omega\Theta K}{}^S X^K)=\\
=-\tfrac{4}{n-2}\eta_\Psi{}^{CD}\eta^\Omega{}_{SD}R_{\Omega\Phi K}{}^S X^K+\tfrac{4}{(n-1)(n-2)}\eta_\Psi{}^{CD}\eta_{\Phi AD}\eta^{\Theta AR}\eta^\Omega{}_{SR}R_{\Omega\Theta K}{}^S X^K.
\end{gathered}\tag{16.39}$$

Таким образом условия интегрируемости примут вид

$$R_{\Phi\Psi K}{}^C X^K=-\tfrac{4}{n-2}\eta_{[\Psi}{}^{CD}\eta^\Omega{}_{|SD}R_{\Omega|\Phi]K}{}^S X^K+\tfrac{4}{(n-1)(n-2)}\eta_{[\Psi}{}^{CD}\eta_{\Phi]AD}\eta^{\Theta AL}\eta^\Omega{}_{SL}R_{\Omega\Theta K}{}^S X^K.\tag{16.40}$$

Будем считать, что данное условие выполнено для любого X^K. Воспользуемся тождеством Бианки $2R^{\Delta[\Phi\Psi]\Lambda}=R^{\Psi\Phi\Delta\Lambda}$

$$\begin{gathered}
(R_{\Phi\Psi K}{}^C=-\tfrac{4}{n-2}\eta_{[\Psi}{}^{CD}\eta^\Omega{}_{|SD}R_{\Omega|\Phi]K}{}^S+\tfrac{4}{(n-1)(n-2)}\eta_{[\Psi}{}^{CD}\eta_{\Phi]AD}\eta^{\Theta AL}\eta^\Omega{}_{SL}R_{\Omega\Theta K}{}^S)\times\\
\times\tfrac{4}{N}\eta^{[\Lambda|KM|}\eta^{\Delta]}{}_{CM},
\end{gathered}$$

$$\begin{gathered}
R^{\Phi\Psi\Lambda\Delta}=-\tfrac{4}{n-2}\tfrac{4}{N}\eta^{[\Psi|CD}\eta^\Omega{}_{SD}\eta^{[\Sigma|SL|}\eta^{\Theta]}{}_{KL}\eta^{[\Lambda|KM|}\eta^{\Delta]|}{}_{CM}\tfrac{1}{2}R_\Omega{}^{\Phi]}{}_{\Sigma\Theta}+\\
+\tfrac{4}{(n-1)(n-2)}\tfrac{4}{N}\eta^{[\Psi|CD|}\eta^{\Phi]}{}_{AD}\eta^{[\Theta|AL|}\eta^{\Omega]}{}_{SL}\eta^{[\Sigma|SN|}\eta^{\Xi]}{}_{KN}\eta^{[\Lambda|KM|}\eta^{\Delta]}{}_{CM}\tfrac{1}{2}R_{\Omega\Theta\Sigma\Xi}=\\
=-\tfrac{8}{n-2}(\tfrac{1}{2}g^{[\Psi|[\Delta}g^{\Lambda]\Theta}g^{\Sigma\Omega|}R_\Omega{}^{\Phi]}{}_{\Sigma\Theta}-\tfrac{1}{2}g^{\Omega[\Delta}g^{\Lambda]\Theta}g^{\Sigma[\Psi}R_\Omega{}^{\Phi]}{}_{\Sigma\Theta}+\tfrac{1}{4}g^{[\Psi|\Omega}g^{\Sigma[\Delta}g^{\Lambda]\Theta|}R_\Omega{}^{\Phi]}{}_{\Sigma\Theta})+\\
+\tfrac{8}{(n-1)(n-2)}(\tfrac{1}{4}g^{\Psi[\Delta}g^{\Lambda]\Phi}g^{\Theta\Xi}g^{\Sigma\Omega}R_{\Omega\Theta\Sigma\Xi}-\tfrac{1}{4}g^{\Psi\Xi}g^{\Sigma\Phi}g^{\Omega\Delta}g^{\Lambda\Theta}R_{\Omega\Theta\Sigma\Xi}+\tfrac{1}{4}g^{\Psi\Omega}g^{\Theta\Phi}g^{\Sigma\Delta}g^{\Lambda\Xi}R_{\Omega\Theta\Sigma\Xi}+\\
+g^{\Theta[\Phi}g^{\Psi][\Delta}g^{\Lambda]\Xi}g^{\Sigma\Omega}R_{\Omega\Theta\Sigma\Xi}+g^{\Theta[\Lambda}g^{\Delta][\Phi}g^{\Psi]\Xi}g^{\Sigma\Omega}R_{\Omega\Theta\Sigma\Xi}+g^{\Theta[\Lambda}g^{\Delta]\Sigma}g^{\Xi[\Psi}g^{\Phi]\Omega}R_{\Omega\Theta\Sigma\Xi})=\\
=-\tfrac{4}{n-2}g^{[\Delta|[\Psi}R^{\Phi]|\Lambda]}+\tfrac{4}{n-2}R^{[\Delta|[\Phi\Psi]|\Lambda]}-\tfrac{2}{n-2}R^{\Psi\Phi\Delta\Lambda}+\\
+\tfrac{2}{(n-1)(n-2)}Rg^{\Psi[\Delta}g^{\Lambda]\Phi}-\tfrac{2}{(n-1)(n-2)}R^{\Delta\Lambda\Phi\Psi}+\tfrac{2}{(n-1)(n-2)}R^{\Psi\Phi\Delta\Lambda}+\\
+\tfrac{8}{(n-1)(n-2)}R^{[\Phi|[\Lambda}g^{\Delta]|\Psi]}+\tfrac{8}{(n-1)(n-2)}g^{[\Phi|[\Delta}R^{\Lambda]|\Psi]}+\tfrac{8}{(n-1)(n-2)}R^{[\Phi|[\Lambda\Delta]|\Psi]}.
\end{gathered}\tag{16.41}$$

Окончательно

$$R^{\Phi\Psi\Lambda\Delta}-\frac{4}{n-2}R^{[\Lambda|[\Phi}g^{\Psi]|\Delta]}+\frac{2}{(n-1)(n-2)}Rg^{[\Delta|[\Psi}g^{\Phi]|\Lambda]}=C^{\Phi\Psi\Lambda\Delta}=0.\tag{16.42}$$

16.8 Доказательство тождеств для n=6

16.8.1 Доказательство основных тождеств для n=6

Необходимо доказать (14.8).

$$A_{\alpha\beta d}{}^{c} = \eta_{[\alpha}{}^{ca}\eta_{\beta]da}, \tag{16.43}$$

$$\begin{gathered}
A_{\alpha\beta d}{}^{c}A_{\gamma}{}^{\beta}{}_{r}{}^{s} = \tfrac{1}{4}(\eta_{\alpha}{}^{ca}\eta_{\beta ad} - \eta_{\alpha ad}\eta_{\beta}{}^{ca})(\eta_{\gamma}{}^{sa_1}\eta^{\beta}{}_{a_1r} - \eta_{\gamma a_1r}\eta^{\beta sa_1}) = \\
= \tfrac{1}{4}(\eta_{\alpha}{}^{ca}\eta_{\gamma}{}^{sa_1}\varepsilon_{a_1rad} - \eta_{\alpha ad}\eta_{\gamma}{}^{sa_1}\varepsilon_{a_1r}{}^{ca} - \eta_{\alpha}{}^{ca}\eta_{\gamma a_1r}\varepsilon_{ad}{}^{sa_1} + \eta_{\alpha ad}\eta_{\gamma a_1r}\varepsilon^{casa_1}) = \\
= \tfrac{1}{4}(\tfrac{1}{2}\eta_{\alpha}{}^{ca}\eta_{\gamma kl}\varepsilon^{klsa_1}\varepsilon_{a_1rad} - \eta_{\alpha ad}\eta_{\gamma}{}^{sa_1}(\delta_{a_1}{}^{c}\delta_{r}{}^{a} - \delta_{a_1}{}^{a}\delta_{r}{}^{c}) - \\
- \eta_{\alpha}{}^{ca}\eta_{\gamma a_1r}(\delta_{a}{}^{s}\delta_{d}{}^{a_1} - \delta_{a}{}^{a_1}\delta_{d}{}^{s}) + \tfrac{1}{2}\eta_{\alpha}{}^{kl}\eta_{\gamma a_1r}\varepsilon_{klad}\varepsilon^{casa_1}) = \\
= \tfrac{1}{4}(-3\eta_{\alpha}{}^{ca}\eta_{\gamma[ra}\delta_{d]}{}^{s} - \eta_{\alpha rd}\eta_{\gamma}{}^{sc} + \eta_{\alpha ad}\eta_{\gamma}{}^{sa}\delta_{r}{}^{c} - \\
- \eta_{\alpha}{}^{cs}\eta_{\gamma dr} + \eta_{\alpha}{}^{ca}\eta_{\gamma ar}\delta_{d}{}^{s} - 3\eta_{\alpha}{}^{[cs}\delta_{d}{}^{a_1]}\eta_{\gamma a_1r} = \\
= \tfrac{1}{4}(-\eta_{\alpha}{}^{ck}\eta_{\gamma rk}\delta_{d}{}^{s} + \eta_{\alpha}{}^{cs}\eta_{\gamma rd} - \eta_{\alpha}{}^{ck}\eta_{\gamma kd}\delta_{r}{}^{s} - \eta_{\alpha rd}\eta_{\gamma}{}^{sc} + \eta_{\alpha ad}\eta_{\gamma}{}^{sa}\delta_{r}{}^{c} - \\
- \eta_{\alpha}{}^{cs}\eta_{\gamma dr} + \eta_{\alpha}{}^{ca}\eta_{\gamma ar}\delta_{d}{}^{s} - \eta_{\alpha}{}^{cs}\eta_{\gamma dr} + \eta_{\alpha}{}^{ck}\eta_{\gamma kr}\delta_{d}{}^{s} - \eta_{\alpha}{}^{sk}\eta_{\gamma kr}\delta_{d}{}^{c}) = \\
= \tfrac{1}{4}(3\eta_{\alpha}{}^{cs}\eta_{\gamma rd} + 3\eta_{\alpha}{}^{ck}\eta_{\gamma kr}\delta_{d}{}^{s} + \eta_{\alpha}{}^{ck}\eta_{\gamma dk}\delta_{r}{}^{s} + \eta_{\alpha dr}\eta_{\gamma}{}^{sc} + \eta_{\alpha kd}\eta_{\gamma}{}^{sk}\delta_{r}{}^{c} + \eta_{\alpha}{}^{sk}\eta_{\gamma rk}\delta_{d}{}^{c}) =
\end{gathered} \tag{16.44}$$

$$\left|\begin{gathered}
\eta_{\alpha dr}\eta_{\gamma}{}^{sc} + \eta_{\alpha kd}\eta_{\gamma}{}^{sk}\delta_{r}{}^{c} = \tfrac{1}{4}(\eta_{\alpha}{}^{mm_1}\eta_{\gamma ll_1}\varepsilon_{mm_1dr}\varepsilon^{ll_1sc} + \eta_{\alpha}{}^{mm_1}\eta_{\gamma ll_1}\varepsilon_{mm_1kd}\varepsilon^{ll_1sk}\delta_{r}{}^{c}) = \\
= 6\eta_{\alpha}{}^{mm_1}\eta_{\gamma ll_1}\delta_{[m}{}^{l}\delta_{m_1}{}^{l_1}\delta_{d}{}^{s}\delta_{r]}{}^{c} - \tfrac{3}{2}\eta_{\alpha}{}^{mm_1}\eta_{\gamma ll_1}\delta_{[m}{}^{l}\delta_{m_1}{}^{l_1}\delta_{d]}{}^{s}\delta_{r}{}^{c} = \\
\\
= -g_{\alpha\gamma}\delta_{r}{}^{s}\delta_{d}{}^{c} + \eta_{\alpha}{}^{kc}\eta_{\gamma kd}\delta_{r}{}^{s} + \eta_{\alpha}{}^{ck}\eta_{\gamma kr}\delta_{d}{}^{s} + \eta_{\alpha}{}^{ks}\eta_{\gamma kr}\delta_{d}{}^{c} + \eta_{\alpha}{}^{sc}\eta_{\gamma dr}.
\end{gathered}\right| \tag{16.45}$$

$$= \eta_{\alpha}{}^{cs}\eta_{\gamma rd} + \eta_{\alpha}{}^{ck}\eta_{\gamma kr}\delta_{d}{}^{s} + \tfrac{1}{2}(\eta_{\alpha}{}^{ck}\eta_{\gamma dk}\delta_{r}{}^{s} + \eta_{\alpha}{}^{sk}\eta_{\gamma rk}\delta_{d}{}^{c}) - \tfrac{1}{4}g_{\alpha\gamma}\delta_{r}{}^{s}\delta_{d}{}^{c}. \tag{16.46}$$

Свернем с $g^{\alpha\gamma}$

$$\begin{gathered}
A_{\alpha\beta d}{}^{c}A^{\alpha\beta}{}_{r}{}^{s} = \varepsilon_{rd}{}^{cs} + \varepsilon_{kr}{}^{ck}\delta_{d}{}^{s} + \tfrac{1}{2}(\varepsilon^{ck}{}_{dk}\delta_{r}{}^{s} + \varepsilon^{sk}{}_{rk}\delta_{d}{}^{c}) - \tfrac{3}{2}\delta_{r}{}^{s}\delta_{d}{}^{c} = \\
= \delta_{r}{}^{c}\delta_{d}{}^{s} - \delta_{r}{}^{s}\delta_{d}{}^{c} - 3\delta_{r}{}^{c}\delta_{d}{}^{s} + \tfrac{3}{2}\delta_{d}{}^{c}\delta_{r}{}^{s} + \tfrac{3}{2}\delta_{d}{}^{c}\delta_{r}{}^{s} - \tfrac{3}{2}\delta_{r}{}^{s}\delta_{d}{}^{c} = \tfrac{1}{2}\delta_{r}{}^{s}\delta_{d}{}^{c} - 2\delta_{r}{}^{c}\delta_{d}{}^{s}.
\end{gathered} \tag{16.47}$$

$$\begin{gathered}
A_{\alpha\beta d}{}^{c}A^{\lambda\mu}{}_{c}{}^{d} = \tfrac{1}{2}\eta_{[\alpha}{}^{ca}\eta_{\beta]}{}^{nn_1}\varepsilon_{adnn_1} \cdot \tfrac{1}{2}\eta^{[\lambda}{}_{ll_1}\eta^{\mu]}{}_{a_1c}\varepsilon^{da_1ll_1} = \\
= \tfrac{1}{4}\eta_{[\alpha}{}^{ca}\eta_{\beta]}{}^{nn_1}\eta^{[\lambda}{}_{ll_1}\eta^{\mu]}{}_{a_1c} \cdot (-6\delta_{[a}{}^{a_1}\delta_{n}{}^{l}\delta_{n_1]}{}^{l_1}) = \\
= -\tfrac{1}{2}(\eta_{[\alpha}{}^{ca}\eta_{\beta]}{}^{nn_1}\eta^{[\lambda}{}_{nn_1}\eta^{\mu]}{}_{ac} - 2\eta_{[\alpha}{}^{ca}\eta_{\beta]}{}^{a_1k_1}\eta^{[\lambda}{}_{ak_1}\eta^{\mu]}{}_{a_1c}) = \\
= -2\delta_{[\alpha}{}^{\lambda}\delta_{\beta]}{}^{\mu} + (\eta_{\alpha}{}^{ca}\eta_{\beta}{}^{a_1k_1}\eta^{[\lambda}{}_{ak_1}\eta^{\mu]}{}_{a_1c} - \eta_{\beta}{}^{ca}\eta_{\alpha}{}^{a_1k_1}\eta^{[\lambda}{}_{ak_1}\eta^{\mu]}{}_{a_1c}) = \\
= 2\delta_{[\alpha}{}^{\mu}\delta_{\beta]}{}^{\lambda} + (\eta_{\alpha}{}^{ca}\eta_{\beta}{}^{a_1k_1}\eta^{[\lambda}{}_{ak_1}\eta^{\mu]}{}_{a_1c} - \eta_{\beta}{}^{a_1k_1}\eta_{\alpha}{}^{ca}\eta^{[\lambda}{}_{k_1a}\eta^{\mu]}{}_{ca_1}) = 2\delta_{[\alpha}{}^{\mu}\delta_{\beta]}{}^{\lambda}.
\end{gathered} \tag{16.48}$$

$$\begin{gathered}
A^{\alpha\beta}{}_{m}{}^{n}T_{\alpha\beta} = A^{\alpha\beta}{}_{m}{}^{n}A_{\alpha\beta k}{}^{l}T_{l}{}^{k} = \\
= (\tfrac{1}{2}\delta_{m}{}^{n}\delta_{k}{}^{l} - 2\delta_{k}{}^{n}\delta_{m}{}^{l})T_{l}{}^{k} = -2T_{m}{}^{n}, \quad T_{\alpha\beta} = -T_{\beta\alpha}.
\end{gathered} \tag{16.49}$$

16.8.2 Доказательство тождеств о 4-векторе

Необходимо доказать (14.9).

$$e_{\alpha\beta\gamma\delta} = e_{[\alpha\beta\gamma\delta]} = A_{\alpha\beta b}{}^{a} A_{\gamma\delta d}{}^{c} e_{a}{}^{b}{}_{c}{}^{d}, \quad e_{k}{}^{k}{}_{c}{}^{d} = e_{a}{}^{b}{}_{k}{}^{k} = 0, \quad e_{a}{}^{b}{}_{c}{}^{d} = e_{c}{}^{d}{}_{a}{}^{b}. \tag{16.50}$$

Свертка с $A^{\alpha\beta}{}_{m}{}^{n} A^{\gamma\delta}{}_{r}{}^{s}$ даст

$$\begin{gathered}
A^{\alpha\beta}{}_{m}{}^{n} A^{\gamma\delta}{}_{r}{}^{s} A_{\alpha\beta b}{}^{a} A_{\gamma\delta d}{}^{c} e_{a}{}^{b}{}_{c}{}^{d} = \\
= (\tfrac{1}{2}\delta_{r}{}^{s}\delta_{d}{}^{c} - 2\delta_{r}{}^{c}\delta_{d}{}^{s})(\tfrac{1}{2}\delta_{b}{}^{a}\delta_{m}{}^{n} - 2\delta_{b}{}^{n}\delta_{m}{}^{a}) e_{a}{}^{b}{}_{c}{}^{d} = 4e_{m}{}^{n}{}_{r}{}^{s} = \\
= -A^{\beta\alpha}{}_{m}{}^{n} A^{\gamma\delta}{}_{r}{}^{s} A_{\gamma\alpha b}{}^{a} A_{\beta\delta d}{}^{c} e_{a}{}^{b}{}_{c}{}^{d} = \\
= -(\eta_{\gamma}{}^{an}\eta^{\beta}{}_{mb} + \eta_{\gamma}{}^{ak}\eta^{\beta}{}_{km}\delta_{b}{}^{n} + \tfrac{1}{2}\eta_{\gamma}{}^{ak}\eta^{\beta}{}_{bk}\delta_{m}{}^{n})\cdot \\
\cdot(\eta_{\beta}{}^{cs}\eta^{\gamma}{}_{rd} + \eta_{\beta}{}^{ct}\eta^{\gamma}{}_{tr}\delta_{d}{}^{s} + \tfrac{1}{2}\eta_{\beta}{}^{ct}\eta^{\gamma}{}_{dt}\delta_{r}{}^{s}) e_{a}{}^{b}{}_{c}{}^{d} = \\
= -(\varepsilon_{rd}{}^{an}\varepsilon_{mb}{}^{cs} + \varepsilon_{mb}{}^{ck}\varepsilon_{kr}{}^{an}\delta_{d}{}^{s} + \tfrac{1}{2}\varepsilon_{mb}{}^{ck}\varepsilon_{dk}{}^{an}\delta_{r}{}^{s} + \\
+\varepsilon_{rd}{}^{ak}\varepsilon_{km}{}^{cs}\delta_{b}{}^{n} + \varepsilon^{ak}{}_{tr}\varepsilon^{ct}{}_{km}\delta_{d}{}^{s}\delta_{b}{}^{n} + \tfrac{1}{2}\varepsilon^{ak}{}_{dt}\varepsilon^{ct}{}_{km}\delta_{r}{}^{s}\delta_{b}{}^{n} + \\
+\tfrac{1}{2}\varepsilon_{rd}{}^{ak}\varepsilon_{bk}{}^{cs}\delta_{m}{}^{n} + \tfrac{1}{2}\varepsilon_{bk}{}^{ct}\varepsilon_{tr}{}^{ak}\delta_{m}{}^{n}\delta_{d}{}^{s} + \tfrac{1}{4}\varepsilon_{bk}{}^{ct}\varepsilon_{dt}{}^{ak}\delta_{m}{}^{n}\delta_{r}{}^{s}) e_{a}{}^{b}{}_{c}{}^{d} = \\
= -(\delta_{r}{}^{a}\delta_{d}{}^{n}\delta_{m}{}^{c}\delta_{b}{}^{s} - \delta_{d}{}^{a}\delta_{r}{}^{n}\delta_{m}{}^{c}\delta_{b}{}^{s} + \delta_{d}{}^{a}\delta_{r}{}^{n}\delta_{b}{}^{c}\delta_{m}{}^{s} - \delta_{r}{}^{a}\delta_{d}{}^{n}\delta_{b}{}^{c}\delta_{m}{}^{s} + \\
+\delta_{m}{}^{c}\delta_{b}{}^{k}\delta_{k}{}^{a}\delta_{r}{}^{n}\delta_{d}{}^{s} - \delta_{m}{}^{k}\delta_{b}{}^{c}\delta_{k}{}^{a}\delta_{r}{}^{n}\delta_{d}{}^{s} + \delta_{m}{}^{k}\delta_{b}{}^{c}\delta_{k}{}^{n}\delta_{r}{}^{a}\delta_{d}{}^{s} - \delta_{m}{}^{c}\delta_{b}{}^{k}\delta_{k}{}^{n}\delta_{r}{}^{a}\delta_{d}{}^{s} + \\
+\tfrac{1}{2}\delta_{m}{}^{c}\delta_{b}{}^{k}\delta_{d}{}^{a}\delta_{k}{}^{n}\delta_{r}{}^{s} - \tfrac{1}{2}\delta_{m}{}^{k}\delta_{b}{}^{c}\delta_{d}{}^{a}\delta_{k}{}^{n}\delta_{r}{}^{s} + \tfrac{1}{2}\delta_{m}{}^{k}\delta_{b}{}^{c}\delta_{d}{}^{n}\delta_{k}{}^{a}\delta_{r}{}^{s} - \tfrac{1}{2}\delta_{m}{}^{c}\delta_{b}{}^{k}\delta_{d}{}^{n}\delta_{k}{}^{a}\delta_{r}{}^{s} + \\
+\delta_{r}{}^{a}\delta_{d}{}^{k}\delta_{k}{}^{c}\delta_{m}{}^{s}\delta_{b}{}^{n} - \delta_{r}{}^{k}\delta_{d}{}^{a}\delta_{k}{}^{c}\delta_{m}{}^{s}\delta_{b}{}^{n} + \delta_{r}{}^{k}\delta_{d}{}^{a}\delta_{k}{}^{s}\delta_{m}{}^{c}\delta_{b}{}^{n} - \delta_{r}{}^{a}\delta_{d}{}^{k}\delta_{k}{}^{s}\delta_{m}{}^{c}\delta_{b}{}^{n} + \\
+\delta_{m}{}^{a}\delta_{r}{}^{c}\delta_{d}{}^{s}\delta_{b}{}^{n} + 2\delta_{m}{}^{c}\delta_{r}{}^{a}\delta_{d}{}^{s}\delta_{b}{}^{n} - \tfrac{1}{2}\delta_{m}{}^{a}\delta_{d}{}^{c}\delta_{r}{}^{s}\delta_{b}{}^{n} - \delta_{m}{}^{c}\delta_{d}{}^{a}\delta_{r}{}^{s}\delta_{b}{}^{n} + \\
+\tfrac{1}{2}\delta_{r}{}^{a}\delta_{d}{}^{k}\delta_{b}{}^{c}\delta_{k}{}^{s}\delta_{m}{}^{n} - \tfrac{1}{2}\delta_{r}{}^{k}\delta_{d}{}^{a}\delta_{b}{}^{c}\delta_{k}{}^{s}\delta_{m}{}^{n} + \tfrac{1}{2}\delta_{r}{}^{k}\delta_{d}{}^{a}\delta_{b}{}^{s}\delta_{k}{}^{c}\delta_{m}{}^{n} - \tfrac{1}{2}\delta_{r}{}^{a}\delta_{d}{}^{k}\delta_{b}{}^{s}\delta_{k}{}^{c}\delta_{m}{}^{n} - \\
-\tfrac{1}{2}\delta_{b}{}^{a}\delta_{r}{}^{c}\delta_{m}{}^{n}\delta_{d}{}^{s} - \delta_{b}{}^{c}\delta_{r}{}^{a}\delta_{m}{}^{n}\delta_{d}{}^{s} + \tfrac{1}{4}\delta_{b}{}^{a}\delta_{d}{}^{c}\delta_{m}{}^{n}\delta_{r}{}^{s} + \tfrac{1}{2}\delta_{b}{}^{c}\delta_{d}{}^{a}\delta_{m}{}^{n}\delta_{r}{}^{s}) e_{a}{}^{b}{}_{c}{}^{d} = \\
= -(e_{r}{}^{s}{}_{m}{}^{n} - e_{t}{}^{s}{}_{m}{}^{t}\delta_{r}{}^{n} + e_{k}{}^{t}{}_{t}{}^{k}\delta_{r}{}^{n}\delta_{m}{}^{s} - e_{r}{}^{t}{}_{t}{}^{n}\delta_{m}{}^{s} - e_{m}{}^{t}{}_{t}{}^{s}\delta_{r}{}^{n} + e_{r}{}^{t}{}_{t}{}^{s}\delta_{m}{}^{n} - \\
-e_{r}{}^{n}{}_{m}{}^{s} + \tfrac{1}{2}e_{t}{}^{n}{}_{m}{}^{t}\delta_{r}{}^{s} - \tfrac{1}{2}e_{p}{}^{t}{}_{t}{}^{p}\delta_{m}{}^{n}\delta_{r}{}^{s} + \tfrac{1}{2}e_{m}{}^{t}{}_{t}{}^{n}\delta_{r}{}^{s} - e_{t}{}^{n}{}_{r}{}^{t}\delta_{m}{}^{s} + e_{t}{}^{n}{}_{m}{}^{t}\delta_{r}{}^{s} - \\
-e_{r}{}^{n}{}_{m}{}^{s} + e_{m}{}^{n}{}_{r}{}^{s} + 2e_{r}{}^{n}{}_{m}{}^{s} - e_{t}{}^{n}{}_{m}{}^{t}\delta_{r}{}^{s} + \tfrac{1}{2}e_{r}{}^{t}{}_{t}{}^{s}\delta_{m}{}^{n} - \\
-\tfrac{1}{2}e_{t}{}^{p}{}_{p}{}^{t}\delta_{r}{}^{s}\delta_{m}{}^{n} + \tfrac{1}{2}e_{t}{}^{s}{}_{r}{}^{t}\delta_{m}{}^{n} - e_{r}{}^{t}{}_{t}{}^{s}\delta_{m}{}^{n} + \tfrac{1}{2}e_{t}{}^{p}{}_{p}{}^{t}\delta_{m}{}^{n}\delta_{r}{}^{s}).
\end{gathered} \tag{16.51}$$

$$e_{m}{}^{s} := \tfrac{1}{3}e_{m}{}^{t}{}_{t}{}^{s} \quad \Rightarrow \quad 2e_{m}{}^{n}{}_{r}{}^{s} = 2(e_{m}{}^{s}\delta_{r}{}^{n} + e_{r}{}^{n}\delta_{m}{}^{s}) - (e_{r}{}^{s}\delta_{m}{}^{n} + e_{m}{}^{n}\delta_{r}{}^{s}). \tag{16.52}$$

16.8.3 Доказательство формул о 6-векторе

Необходимо доказать (14.10).

$$\begin{gathered}
e_{\alpha\beta\gamma\delta\rho\sigma} = e_{[\alpha\beta\gamma\delta\rho\sigma]} = A_{\alpha\beta b}{}^{a} A_{\gamma\delta d}{}^{c} A_{\rho\sigma s}{}^{r} e_{a}{}^{b}{}_{c}{}^{d}{}_{r}{}^{s}, \\
6e_{m}{}^{n}{}_{p}{}^{q}{}_{r}{}^{s} = 2(e_{m}{}^{s}{}_{p}{}^{q}\delta_{r}{}^{n} + e_{r}{}^{n}{}_{p}{}^{q}\delta_{m}{}^{s}) - (e_{r}{}^{s}{}_{p}{}^{q}\delta_{m}{}^{n} + e_{m}{}^{n}{}_{p}{}^{q}\delta_{r}{}^{s}) = \\
= 2(e_{m}{}^{q}{}_{r}{}^{s}\delta_{p}{}^{n} + e_{p}{}^{n}{}_{r}{}^{s}\delta_{m}{}^{q}) - (e_{p}{}^{q}{}_{r}{}^{s}\delta_{m}{}^{n} + e_{m}{}^{n}{}_{r}{}^{s}\delta_{p}{}^{q}), \quad e_{m}{}^{s}{}_{p}{}^{q} := e_{m}{}^{k}{}_{k}{}^{s}{}_{p}{}^{q} = e_{k}{}^{s}{}_{p}{}^{q}{}_{m}{}^{k}.
\end{gathered} \tag{16.53}$$

1. Первая свертка.

   Свертка с $\delta_q{}^p$ даст

$$0 = 6e_m{}^n{}_r{}^s{}_p{}^p = 2(e_m{}^n{}_r{}^s + e_m{}^n{}_r{}^s) - (e_p{}^p{}_r{}^s\delta_m{}^n + 4e_m{}^n{}_r{}^s) = -(e_p{}^p{}_r{}^s)\cdot\delta_m{}^n = 0,$$
$$e_p{}^p{}_r{}^s = e_r{}^s{}_p{}^p = 0. \tag{16.54}$$

2. Вторая свертка.

   Свертка с $\delta_n{}^r$ даст

$$8e_m{}^s{}_p{}^q + e_t{}^t{}_p{}^q\delta_m{}^s - e_m{}^s{}_p{}^q - e_m{}^s{}_p{}^q = 2(e_m{}^q{}_p{}^s + e_p{}^k{}_k{}^s\delta_m{}^q) - (e_p{}^q{}_m{}^s + e_m{}^k{}_k{}^s\delta_p{}^q),$$
$$6e_m{}^s{}_p{}^q + e_p{}^q{}_m{}^s - 2e_m{}^q{}_p{}^s = 2e_p{}^t{}_t{}^s\delta_m{}^q - e_m{}^k{}_k{}^s\delta_p{}^q. \tag{16.55}$$

1. Первая свертка.

   Свернем с $\delta_q{}^m$ и получим

$$6e_k{}^s{}_p{}^k + e_p{}^k{}_k{}^s = 8e_p{}^t{}_t{}^s - e_p{}^k{}_k{}^s, \quad e_p{}^t{}_t{}^q = e_t{}^q{}_p{}^t. \tag{16.56}$$

2. Вторая свертка.

   Свернем с $\delta_s{}^m$ и получим

$$10\tilde{e} := e_t{}^k{}_k{}^t, \quad -2e_k{}^q{}_p{}^k = 2e_p{}^t{}_t{}^q - e_t{}^k{}_k{}^t\delta_p{}^q, \quad e_p{}^t{}_t{}^q = \tfrac{5\tilde{e}}{2}\delta_p{}^q. \tag{16.57}$$

3. Третья свертка.

   Свернем с $\delta_s{}^p$ и получим

$$2e_m{}^k{}_k{}^q\delta_r{}^n + e_r{}^n{}_m{}^q - e_r{}^k{}_k{}^q\delta_m{}^n - e_m{}^n{}_r{}^q = 2e_m{}^q{}_r{}^n + 2e_k{}^n{}_r{}^k\delta_m{}^q - e_k{}^q{}_r{}^k\delta_m{}^n - e_m{}^n{}_r{}^q,$$
$$e_r{}^n{}_m{}^q = e_m{}^q{}_r{}^n. \tag{16.58}$$

$$\begin{gathered}
7e_m{}^s{}_p{}^q - 2e_m{}^q{}_p{}^s = 5\tilde{e}\delta_m{}^q\delta_p{}^s - \tfrac{5\tilde{e}}{2}\delta_m{}^s\delta_p{}^q, \quad 9e_m{}^{[s}{}_p{}^{q]} = \tfrac{15\tilde{e}}{2}\delta_m{}^{[q}\delta_p{}^{s]},\\
5e_m{}^s{}_p{}^q + 4e_m{}^{[s}{}_p{}^{q]} = 5\tilde{e}\delta_m{}^q\delta_p{}^s - \tfrac{5\tilde{e}}{2}\delta_m{}^s\delta_p{}^q, \quad 6e_m{}^s{}_p{}^q = \tilde{e}(4\delta_m{}^q\delta_p{}^s - \delta_m{}^s\delta_p{}^q),\\
e_m{}^n{}_r{}^s{}_p{}^q = \tilde{e}(2((4\delta_p{}^n\delta_r{}^q - \delta_p{}^q\delta_r{}^n)\delta_m{}^s + (4\delta_p{}^s\delta_m{}^q - \delta_p{}^q\delta_m{}^s)\delta_r{}^n)-\\
-((4\delta_p{}^n\delta_m{}^q - \delta_p{}^q\delta_m{}^n)\delta_r{}^s + (4\delta_p{}^s\delta_r{}^q - \delta_p{}^q\delta_r{}^s)\delta_m{}^n)).
\end{gathered} \tag{16.59}$$

Вычислим $\tilde{e}$

$$\begin{gathered}
720 = e^{\alpha\beta\gamma\delta\mu\nu}e_{\alpha\beta\gamma\delta\mu\nu} = A^{\alpha\beta}{}_{b_1}{}^{a_1}A_{\alpha\beta b}{}^{a}A^{\gamma\delta}{}_{d_1}{}^{c_1}A_{\gamma\delta d}{}^{c}A^{\mu\nu}{}_{n_1}{}^{m_1}A_{\mu\nu n}{}^{m},\\
e_a{}^b{}_c{}^d{}_m{}^n e_{a_1}{}^{b_1}{}_{c_1}{}^{d_1}{}_{m_1}{}^{n_1} = 8(\tfrac{1}{4}\delta_{b_1}{}^{a_1}\delta_b{}^a - \delta_{b_1}{}^a\delta_b{}^{a_1})(\tfrac{1}{4}\delta_{d_1}{}^{c_1}\delta_d{}^c - \delta_{d_1}{}^c\delta_d{}^{c_1})\cdot\\
\cdot(\tfrac{1}{4}\delta_{n_1}{}^{m_1}\delta_n{}^m - \delta_{n_1}{}^m\delta_n{}^{m_1})e_a{}^b{}_c{}^d{}_m{}^n e_{a_1}{}^{b_1}{}_{c_1}{}^{d_1}{}_{m_1}{}^{n_1} = -8e_a{}^b{}_c{}^d{}_m{}^n e_b{}^a{}_d{}^c{}_n{}^m =\\
= -8\tilde{e}^2(8\delta_m{}^b\delta_c{}^n\delta_a{}^d - 4\delta_m{}^n\delta_c{}^b\delta_a{}^d + 8\delta_m{}^d\delta_a{}^n\delta_c{}^b - 4\delta_m{}^b\delta_a{}^n\delta_c{}^d + 2\delta_m{}^n\delta_a{}^b\delta_c{}^d - 4\delta_m{}^d\delta_c{}^n\delta_a{}^b)\\
(8\delta_n{}^a\delta_d{}^m\delta_b{}^c - 4\delta_n{}^m\delta_d{}^a\delta_b{}^c + 8\delta_n{}^c\delta_b{}^m\delta_d{}^a - 4\delta_n{}^a\delta_b{}^m\delta_d{}^c + 2\delta_n{}^m\delta_b{}^a\delta_d{}^c - 4\delta_n{}^c\delta_d{}^m\delta_b{}^a) =\\
= -720\cdot 64\tilde{e}^2.
\end{gathered} \tag{16.60}$$

Откуда

$$\tilde{e} = \frac{1}{8}i. \tag{16.61}$$

16.8.4 Доказательство тождества Бианки

Тождество Бианки имеет вид

$$\begin{gathered} R_{\alpha\beta\gamma\delta} + R_{\alpha\gamma\delta\beta} + R_{\alpha\delta\beta\gamma} = 0 \\ (A_{\alpha\beta d}{}^{c} A_{\gamma\delta r}{}^{s} + A_{\alpha\gamma d}{}^{c} A_{\delta\beta r}{}^{s} + A_{\alpha\delta d}{}^{c} A_{\beta\gamma r}{}^{s}) R_{c}{}^{d}{}_{s}{}^{r} = 0. \end{gathered} \tag{16.62}$$

Свернем его с $A^{\alpha\beta}{}_{t}{}^{l} A^{\gamma\delta}{}_{m}{}^{n}$

$$\left| \begin{gathered} A_{\alpha\beta d}{}^{c} A_{\gamma\delta r}{}^{s} R_{c}{}^{d}{}_{s}{}^{r} A^{\alpha\beta}{}_{t}{}^{l} A^{\gamma\delta}{}_{m}{}^{n} R_{c}{}^{d}{}_{s}{}^{r} = \\ = (\tfrac{1}{2}\delta_{d}{}^{c}\delta_{t}{}^{l} - 2\delta_{d}{}^{l}\delta_{t}{}^{c})(\tfrac{1}{2}\delta_{r}{}^{s}\delta_{m}{}^{n} - 2\delta_{r}{}^{n}\delta_{m}{}^{s}) R_{c}{}^{d}{}_{s}{}^{r} = 4R_{t}{}^{l}{}_{m}{}^{n} \end{gathered} \right|$$

$$\left| \begin{gathered} (A_{\alpha\gamma d}{}^{c} A_{\delta\beta r}{}^{s} + A_{\alpha\delta d}{}^{c} A_{\beta\gamma r}{}^{s}) R_{c}{}^{d}{}_{s}{}^{r} A^{\alpha\beta}{}_{t}{}^{l} A^{\gamma\delta}{}_{m}{}^{n} = -2A_{\delta\beta r}{}^{s} A^{\delta\gamma}{}_{m}{}^{n} A_{\alpha\gamma d}{}^{c} A^{\alpha\beta}{}_{t}{}^{l} R_{c}{}^{d}{}_{s}{}^{r} = \\ = -2(\eta_{\beta}{}^{sn}\eta^{\gamma}{}_{mr} + \eta_{\beta}{}^{sk}\eta^{\gamma}{}_{km}\delta_{r}{}^{n} + \tfrac{1}{2}\eta_{\beta}{}^{sk}\eta^{\gamma}{}_{rk}\delta_{m}{}^{n})\cdot \\ \cdot(\eta_{\gamma}{}^{cl}\eta^{\beta}{}_{td} + \eta_{\gamma}{}^{ck_1}\eta^{\beta}{}_{k_1 t}\delta_{d}{}^{l} + \tfrac{1}{2}\eta_{\gamma}{}^{ck_1}\eta^{\beta}{}_{dk_1}\delta_{t}{}^{l}) R_{c}{}^{d}{}_{s}{}^{r} = \\ = -2(\varepsilon_{td}{}^{sn}\varepsilon_{mr}{}^{cl} + \varepsilon^{ck}{}_{mr}\varepsilon_{kt}{}^{sn}\delta_{d}{}^{l} + \tfrac{1}{2}\varepsilon_{mr}{}^{ck}\varepsilon_{dk}{}^{sn}\delta_{t}{}^{l} + \varepsilon_{td}{}^{sk}\varepsilon_{km}{}^{cl}\delta_{r}{}^{n} + \\ + \varepsilon^{ck_1}{}_{km}\varepsilon_{k_1 t}{}^{sk}\delta_{d}{}^{l}\delta_{r}{}^{n} + \tfrac{1}{2}\varepsilon^{ck_1}{}_{km}\varepsilon_{dk_1}{}^{sk}\delta_{t}{}^{l}\delta_{r}{}^{n} + \tfrac{1}{2}(\varepsilon^{cl}{}_{rk}\varepsilon_{td}{}^{sk}\delta_{m}{}^{n} + \varepsilon^{sk}{}_{k_1 t}\varepsilon^{ck_1}{}_{rk}\delta_{d}{}^{l}\delta_{m}{}^{n}) + \\ + \tfrac{1}{4}\varepsilon^{ck_1}{}_{rk}\varepsilon^{sk}{}_{dk_1}\delta_{t}{}^{l}\delta_{m}{}^{n}) R_{c}{}^{d}{}_{s}{}^{r} = -2(R_{m}{}^{n}{}_{t}{}^{l} + R_{k}{}^{r}{}_{r}{}^{k}\delta_{t}{}^{n}\delta_{m}{}^{l} - R_{m}{}^{k}{}_{k}{}^{l}\delta_{t}{}^{n} - R_{t}{}^{k}{}_{k}{}^{n}\delta_{m}{}^{l} - \\ - R_{m}{}^{l}{}_{t}{}^{n} - R_{k}{}^{l}{}_{m}{}^{k}\delta_{t}{}^{n} + R_{k}{}^{l}{}_{t}{}^{k}\delta_{m}{}^{n} + \tfrac{1}{2}(R_{m}{}^{k}{}_{k}{}^{n}\delta_{t}{}^{l} + R_{k}{}^{n}{}_{m}{}^{k}\delta_{t}{}^{l} - R_{k}{}^{r}{}_{r}{}^{k}\delta_{t}{}^{l}\delta_{m}{}^{n}) + \\ + R_{m}{}^{k}{}_{k}{}^{n}\delta_{t}{}^{l} - R_{m}{}^{l}{}_{t}{}^{n} - R_{t}{}^{k}{}_{k}{}^{n}\delta_{m}{}^{l} + 2R_{m}{}^{l}{}_{t}{}^{n} + R_{t}{}^{l}{}_{m}{}^{n} - \tfrac{1}{2}\cdot 2R_{m}{}^{k}{}_{k}{}^{n}\delta_{t}{}^{l} + \tfrac{1}{2}(R_{t}{}^{k}{}_{k}{}^{l}\delta_{m}{}^{n} + \\ + R_{k}{}^{l}{}_{t}{}^{k}\delta_{m}{}^{n} - R_{k}{}^{r}{}_{r}{}^{k}\delta_{m}{}^{n}\delta_{t}{}^{l}) - R_{t}{}^{k}{}_{k}{}^{l}\delta_{m}{}^{n} + \tfrac{1}{2}R_{k}{}^{r}{}_{r}{}^{k}\delta_{t}{}^{l}\delta_{m}{}^{n} = -2(2R_{m}{}^{n}{}_{t}{}^{l} - \\ - 2R_{k}{}^{l}{}_{m}{}^{k}\delta_{t}{}^{n} - 2R_{t}{}^{k}{}_{k}{}^{n}\delta_{m}{}^{l} + R_{k}{}^{l}{}_{t}{}^{k}\delta_{m}{}^{n} + R_{m}{}^{k}{}_{k}{}^{n}\delta_{t}{}^{l} + R_{k}{}^{r}{}_{r}{}^{k}\delta_{t}{}^{n}\delta_{m}{}^{l} - \tfrac{1}{2}R_{k}{}^{r}{}_{r}{}^{k}\delta_{m}{}^{n}\delta_{t}{}^{l}) \end{gathered} \right|$$

$$4R_{k}{}^{l}{}_{m}{}^{k}\delta_{t}{}^{n} + 4R_{t}{}^{k}{}_{k}{}^{n}\delta_{m}{}^{l} - 2R_{k}{}^{l}{}_{t}{}^{k}\delta_{m}{}^{n} - 2R_{m}{}^{k}{}_{k}{}^{n}\delta_{t}{}^{l} - 2R_{k}{}^{r}{}_{r}{}^{k}\delta_{t}{}^{n}\delta_{m}{}^{l} + R_{k}{}^{r}{}_{r}{}^{k}\delta_{m}{}^{n}\delta_{t}{}^{l} = 0. \tag{16.63}$$

Свертка с $\delta_{n}{}^{t}$ даст

$$\begin{gathered} 16R_{k}{}^{l}{}_{m}{}^{k} + 4R_{t}{}^{k}{}_{k}{}^{t}\delta_{m}{}^{l} - 2R_{k}{}^{l}{}_{m}{}^{k} - 2R_{m}{}^{k}{}_{k}{}^{l} - 8R_{k}{}^{r}{}_{r}{}^{k}\delta_{m}{}^{l} + R_{k}{}^{r}{}_{r}{}^{k}\delta_{m}{}^{l} = 0, \\ R_{k}{}^{l}{}_{m}{}^{k} = \tfrac{1}{4}R_{k}{}^{r}{}_{r}{}^{k}\delta_{m}{}^{l} = -\tfrac{1}{8}R\delta_{m}{}^{l}, \end{gathered} \tag{16.64}$$

что и завершает доказательство.

16.8.5 Доказательство тождеств, касающихся тензора Вейля.

Поскольку

$$\begin{gathered} T_{[\alpha\beta]} = \tfrac{1}{4}\eta_{\alpha}{}^{aa_1}\eta_{\beta}{}^{bb_1}\cdot\tfrac{3}{2}(T_{a[a_1 bb_1]} - T_{[bb_1 a]a_1}) = \\ = \tfrac{1}{4}\eta_{\alpha}{}^{aa_1}\eta_{\beta}{}^{bb_1}\cdot\tfrac{1}{2}(T_{ak}{}^{kd}\varepsilon_{a_1 dbb_1} - T^{kd}{}_{ka_1}\varepsilon_{adbb_1}) = \tfrac{1}{2}\eta_{\alpha}{}^{ca}\eta_{\beta}{}^{bk_1}\varepsilon_{adbk_1}\cdot \\ \cdot\tfrac{1}{4}(T^{kd}{}_{kc} - T_{kc}{}^{kd}) = A_{\alpha\beta d}{}^{c}\cdot\tfrac{1}{4}(T^{kd}{}_{kc} - T_{kc}{}^{kd}), \end{gathered} \tag{16.65}$$

то будет верным следующее соотношение

$$\begin{gathered}
R_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]} = A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\tfrac{1}{16}(P^{kdnr}\varepsilon_{kcns} - P_{kc}{}^{nr}\varepsilon^{kd}{}_{ns} + P_{kcns}\varepsilon^{kdnr} - P^{kd}{}_{ns}\varepsilon_{kc}{}^{nr}) = \\
= A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\tfrac{1}{16}(P^{kd}{}_{mm_1}\tfrac{1}{2}\varepsilon^{mm_1nr}\varepsilon_{kcns} + P_{kc}{}^{mm_1}\tfrac{1}{2}\varepsilon_{mm_1ns}\varepsilon^{kdnr} - \\
-P_{kc}{}^{nr}(\delta_n{}^{k}\delta_s{}^{d} - \delta_n{}^{d}\delta_s{}^{k}) - P^{kd}{}_{ns}(\delta_k{}^{n}\delta_c{}^{r} - \delta_k{}^{r}\delta_c{}^{n})) = \\
= A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\tfrac{1}{16}(P^{kd}{}_{kc}\delta_s{}^{r} - P^{kd}{}_{ks}\delta_c{}^{r} + P^{rd}{}_{cs} + \\
+P_{kc}{}^{kd}\delta_s{}^{r} - P_{kc}{}^{kr}\delta_s{}^{d} + P_{sc}{}^{dr} - P_{kc}{}^{kr}\delta_s{}^{d} + P_{sc}{}^{dr} - P^{kd}{}_{ks}\delta_c{}^{r} + P^{rd}{}_{cs}) = \\
= A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\tfrac{1}{4}(P_{sc}{}^{dr} - \tfrac{1}{2}P^{kd}{}_{ks}\delta_c{}^{r} + \tfrac{1}{2}P_{kc}{}^{kd}\delta_s{}^{r} - \tfrac{1}{2}P_{kc}{}^{kr}\delta_s{}^{d}) = \\
= A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\tfrac{1}{4}(P_{sc}{}^{dr} - \tfrac{1}{2}R\delta_s{}^{d}\delta_c{}^{r} + \tfrac{1}{4}R\delta_s{}^{r}\delta_c{}^{d}).
\end{gathered} \tag{16.66}$$

Аналогично

$$\begin{gathered}
g_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]} = A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\tfrac{1}{4}(\varepsilon_{sc}{}^{dr} - \tfrac{1}{2}\varepsilon^{kd}{}_{ks}\delta_c{}^{r} + \tfrac{1}{2}\varepsilon^{kd}{}_{kc}\delta_s{}^{r} - \tfrac{1}{2}\varepsilon_{kc}{}^{kr}\delta_s{}^{d}) = \\
= A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}\tfrac{1}{4}(\tfrac{1}{2}\delta_s{}^{r}\delta_c{}^{d} - 2\delta_s{}^{d}\delta_c{}^{r}).
\end{gathered} \tag{16.67}$$

Таким образом, из

$$C_{\alpha\beta}{}^{\gamma\delta} := R_{\alpha\beta}{}^{\gamma\delta} - R_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]} + 1/10Rg_{[\alpha}{}^{[\gamma}g_{\beta]}{}^{\delta]} \tag{16.68}$$

следует

$$\begin{gathered}
C_{\alpha\beta}{}^{\gamma\delta} = A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}(R_c{}^{d}{}_{s}{}^{r} - \tfrac{1}{4}P_{sc}{}^{dr} + \tfrac{1}{8}R\delta_s{}^{d}\delta_c{}^{r} - \tfrac{1}{16}R\delta_s{}^{r}\delta_c{}^{d} + \tfrac{1}{40}R(\tfrac{1}{2}\delta_s{}^{r}\delta_c{}^{d} - \\
-2\delta_s{}^{d}\delta_c{}^{r})) = A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}(R_{(c}{}^{d}{}_{s)}{}^{r} + R_{[c}{}^{k}{}_{|k|}{}^{[r}\delta_{s]}{}^{d]} + \tfrac{1}{40}R(3\delta_s{}^{d}\delta_c{}^{r} - 2\delta_s{}^{r}\delta_c{}^{d})).
\end{gathered} \tag{16.69}$$

Тогда

$$C_{\alpha\beta}{}^{\gamma\delta} = A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}(R_{(c}{}^{d}{}_{s)}{}^{r} + \tfrac{1}{80}R\delta_s{}^{d}\delta_c{}^{r}). \tag{16.70}$$

Свернем его с $A^{\alpha\beta}{}_{k}{}^{l}A_{\gamma\delta t}{}^{n}$

$$\begin{gathered}
4C_k{}^{l}{}_{t}{}^{n} := A^{\alpha\beta}{}_{k}{}^{l}A_{\gamma\delta t}{}^{n}C_{\alpha\beta}{}^{\gamma\delta} = A^{\alpha\beta}{}_{k}{}^{l}A_{\alpha\beta d}{}^{c}A^{\gamma\delta}{}_{r}{}^{s}A_{\gamma\delta t}{}^{n}(R_{(c}{}^{d}{}_{s)}{}^{r} + \tfrac{1}{80}R\delta_s{}^{d}\delta_c{}^{r}) = \\
= (\tfrac{1}{2}\delta_k{}^{l}\delta_d{}^{c} - 2\delta_k{}^{c}\delta_d{}^{l})(\tfrac{1}{2}\delta_r{}^{s}\delta_t{}^{n} - 2\delta_r{}^{n}\delta_t{}^{s})(\tfrac{1}{2}R_c{}^{d}{}_{s}{}^{r} + \tfrac{1}{2}R_s{}^{d}{}_{c}{}^{r} + \tfrac{1}{80}R\delta_s{}^{d}\delta_c{}^{r}) = \\
= (2R_k{}^{l}{}_{t}{}^{n} + \tfrac{1}{8}R_{k_1}{}^{r_1}{}_{r_1}{}^{k_1}\delta_k{}^{l}\delta_t{}^{n} - \tfrac{1}{2}R_{k_1}{}^{l}{}_{k}{}^{k_1}\delta_t{}^{n} - \tfrac{1}{2}R_t{}^{k_1}{}_{k_1}{}^{n}\delta_k{}^{l} + \\
+2R_t{}^{l}{}_{k}{}^{n} + \tfrac{1}{80}R(\delta_t{}^{n}\delta_k{}^{l} - \delta_t{}^{n}\delta_k{}^{l} - \delta_t{}^{n}\delta_k{}^{l} + 4\delta_t{}^{l}\delta_k{}^{n}) = \\
= (4R_{(k}{}^{l}{}_{t)}{}^{n} - \tfrac{1}{16}R\delta_k{}^{l}\delta_t{}^{n} + \tfrac{1}{16}R\delta_k{}^{l}\delta_t{}^{n} + \tfrac{1}{16}R\delta_t{}^{n}\delta_k{}^{l} + \tfrac{1}{80}R(4\delta_t{}^{l}\delta_k{}^{n} - \delta_t{}^{n}\delta_k{}^{l})) = \\
= 4(R_{(k}{}^{l}{}_{t)}{}^{n} + \tfrac{1}{40}R\delta_{(k}{}^{l}\delta_{t)}{}^{n}),
\end{gathered} \tag{16.71}$$

что завершает доказательство разложения.

16.8.6 Доказательство формул, связанных с метрикой, индуцированной в сечении конуса K_6

Пусть задано сечение конуса K_6

$$T^2 + V^2 - W^2 - X^2 - Y^2 - Z^2 = 0 \tag{16.72}$$

плоскостью V+W=1. Сделаем стереографическую проекцию полученного гиперболоида на плоскость (V=0,W=1).

$$\begin{gathered}
\tfrac{t}{T}=\tfrac{x}{X}=\tfrac{y}{Y}=\tfrac{z}{Z}=-\tfrac{\frac{1}{2}}{V-\frac{1}{2}},\quad x^2+y^2+z^2-t^2=\tfrac{1}{2V-1},\\
T=-t(2V-1),\quad X=-x(2V-1),\quad Y=-y(2V-1),\quad Z=-z(2V-1),\\
dT=-(dt\ (2V-1)+2tdV),\quad dX=-(dx\ (2V-1)+2xdV),\\
dY=-(dy\ (2V-1)+2ydV),\quad dZ=-(dz\ (2V-1)+2zdV),\\
dS^2=dT^2+dV^2-dW^2-dX^2-dY^2-dZ^2=|dV=-dW|=\\
=dT^2-dX^2-dY^2-dZ^2=(dt^2-dx^2-dy^2-dz^2)(2V-1)^2+\\
+4(tdt-xdx-ydy-zdz)dV(2V-1)+4(t^2-x^2-y^2-z^2)dV^2=\\
=|tdt-xdx-ydy-zdz=-\tfrac{1}{2}d(\tfrac{1}{2V-1})|=\tfrac{dt^2-dx^2-dy^2-dz^2}{(t^2-x^2-y^2-z^2)^2}.
\end{gathered}\tag{16.73}$$

Сделаем замену

$$\begin{gathered}
\varsigma=\tfrac{-iX+Y}{2V-1}=-y+ix,\quad \omega=-\tfrac{i(T+Z)}{2V-1}=i(t+z),\quad \eta=\tfrac{i(Z-T)}{2V-1}=i(t-z),\\
\varsigma\bar{\varsigma}+\eta\omega=\tfrac{1}{2V-1},\\
d\varsigma=(\tfrac{dY}{2V-1}-\tfrac{2YdV}{(2V-1)^2})-i(\tfrac{dX}{2V-1}-\tfrac{2XdV}{(2V-1)^2}),\quad d\bar{\varsigma}=(\tfrac{dY}{2V-1}-\tfrac{2YdV}{(2V-1)^2})+i(\tfrac{dX}{2V-1}-\tfrac{2XdV}{(2V-1)^2}),\\
d\omega=-i(\tfrac{dT}{2V-1}-\tfrac{2TdV}{(2V-1)^2}+\tfrac{dZ}{2V-1}-\tfrac{2ZdV}{(2V-1)^2}),\quad d\eta=-i(\tfrac{dT}{2V-1}-\tfrac{2TdV}{(2V-1)^2}-\tfrac{dZ}{2V-1}+\tfrac{2ZdV}{(2V-1)^2}),\\
d\varsigma d\bar{\varsigma}+d\omega d\eta=\tfrac{(dX^2+dY^2+dZ^2-dT^2)}{(2V-1)^2}-4\tfrac{dV}{(2V-1)^3}(XdX+YdY+ZdZ-TdT)+\\
+4\tfrac{dV^2}{(2V-1)^4}(X^2+Y^2+Z^2-T^2)=-\tfrac{dT^2-dX^2-dY^2-dZ^2}{(T^2-X^2-Y^2-Z^2)^2},\\
ds^2:=dT^2-dX^2-dY^2-dZ^2=-\tfrac{d\varsigma d\bar{\varsigma}+d\eta d\omega}{(\varsigma\bar{\varsigma}+\eta\omega)^2},\quad x^2+y^2+z^2-t^2=\tfrac{1}{2V-1}.
\end{gathered}\tag{16.74}$$

Поэтому есть резон положить

$$X:=\begin{pmatrix}\omega & \varsigma\\ -\bar{\varsigma} & \eta\end{pmatrix},\quad dX:=\begin{pmatrix}d\omega & d\varsigma\\ -d\bar{\varsigma} & d\eta\end{pmatrix},\quad \frac{\partial}{\partial X}:=\begin{pmatrix}\frac{\partial}{\partial\omega} & \frac{\partial}{\partial\varsigma}\\ -\frac{\partial}{\partial\bar{\varsigma}} & \frac{\partial}{\partial\eta}\end{pmatrix}.\tag{16.75}$$

Тогда

$$ds^2=-\frac{det(dX)}{(det(X))^2},\quad \bar{X}^T+X=0.\tag{16.76}$$

16.8.7 Доказательство формул о первом инварианте

Рассмотрим группу дробно-линейных преобразований L

$$\tilde{X}=(AX+B)(CX+D)^{-1},\quad S:=\begin{pmatrix}A & B\\ C & D\end{pmatrix},\quad detS=1.\tag{16.77}$$

Пусть имеется два последовательных преобразования

$$\tilde{X}=(AX+B)(CX+D)^{-1},\quad \tilde{\tilde{X}}=(\tilde{A}\tilde{X}+\tilde{B})(\tilde{C}\tilde{X}+\tilde{D})^{-1},\tag{16.78}$$

то

$$\begin{gathered}
\tilde{\tilde{X}} = (\tilde{\tilde{A}}X + \tilde{\tilde{B}})(\tilde{\tilde{C}}X + \tilde{\tilde{D}})^{-1} = \\
= (\tilde{A}(AX+B) + \tilde{B}(CX+D))(\tilde{C}(AX+B) + \tilde{D}(CX+D))^{-1} = \\
= ((\tilde{A}A + \tilde{B}C)X + \tilde{A}B + \tilde{B}D)((\tilde{C}A + \tilde{D}C)X + \tilde{C}B + \tilde{D}D)^{-1}, \\
S := \begin{pmatrix} A & B \\ C & D \end{pmatrix}, \quad \tilde{S} := \begin{pmatrix} \tilde{A} & \tilde{B} \\ \tilde{C} & \tilde{D} \end{pmatrix}, \quad \tilde{\tilde{S}} := \begin{pmatrix} \tilde{\tilde{A}} & \tilde{\tilde{B}} \\ \tilde{\tilde{C}} & \tilde{\tilde{D}} \end{pmatrix}, \quad \tilde{\tilde{S}} = \tilde{S}S.
\end{gathered} \tag{16.79}$$

Для унитарных дробно-линейных преобразований имеем

$$\begin{gathered}
X^* + X \equiv \tilde{X}^* + \tilde{X} = 0, \\
0 = (AX+B)(CX+D)^{-1} + (CX+D)^{*-1}(AX+B)^*, \\
0 = (CX+D)^*(AX+B) + (AX+B)^*(CX+D) = \\
= X^*(A^*C + C^*A)X + X^*(A^*D + C^*B) + (B^*C + D^*A)X + D^*B + B^*D \equiv \\
\equiv X^* + X.
\end{gathered} \tag{16.80}$$

Откуда

$$\begin{gathered}
A^*C + C^*A = 0, \quad B^*D + D^*B = 0, \quad A^*D + C^*B = E, \quad S^*\hat{E}S = \hat{E}, \\
\hat{E} := \begin{pmatrix} 0 & E \\ E & 0 \end{pmatrix}.
\end{gathered} \tag{16.81}$$

Пусть

$$X = YZ^{-1}, \quad \tilde{Y} = AY + BZ, \quad \tilde{Z} = CY + DZ. \tag{16.82}$$

Положим

$$\begin{gathered}
X := \begin{pmatrix} x_1 & x_2 \\ x_3 & x_4 \end{pmatrix}, \quad \hat{X} := \begin{pmatrix} \hat{x}_1 & \hat{x}_2 \\ \hat{x}_3 & \hat{x}_4 \end{pmatrix}, \quad dX := \begin{pmatrix} dx_1 & dx_2 \\ dx_3 & dx_4 \end{pmatrix}, \quad \hat{d}X := \begin{pmatrix} d\hat{x}_1 & d\hat{x}_2 \\ d\hat{x}_3 & d\hat{x}_4 \end{pmatrix}, \\
\tfrac{\partial}{\partial X} := \begin{pmatrix} \dfrac{\partial}{\partial x_1} & \dfrac{\partial}{\partial x_2} \\ \dfrac{\partial}{\partial x_3} & \dfrac{\partial}{\partial x_4} \end{pmatrix}, \quad \tfrac{\partial}{\partial \hat{X}} := \begin{pmatrix} \dfrac{\partial}{\partial \hat{x}_1} & \dfrac{\partial}{\partial \hat{x}_2} \\ \dfrac{\partial}{\partial \hat{x}_3} & \dfrac{\partial}{\partial \hat{x}_4} \end{pmatrix}.
\end{gathered} \tag{16.83}$$

Тогда

$$\begin{gathered}
dX + d\hat{X} = d(X + \hat{X}), \\
d(X\hat{X}) = d\begin{pmatrix} x_1\hat{x}_1 + x_2\hat{x}_3 & x_1\hat{x}_2 + x_2\hat{x}_1 \\ x_3\hat{x}_1 + x_4\hat{x}_3 & x_3\hat{x}_2 + x_4\hat{x}_4 \end{pmatrix} = \\
= \begin{pmatrix} dx_1\hat{x}_1 + dx_2\hat{x}_3 & dx_1\hat{x}_2 + dx_2\hat{x}_1 \\ dx_3\hat{x}_1 + dx_4\hat{x}_3 & dx_3\hat{d}x_2 + x_4\hat{x}_4 \end{pmatrix} + \begin{pmatrix} x_1d\hat{x}_1 + x_2d\hat{x}_3 & x_1d\hat{x}_2 + x_2d\hat{x}_1 \\ x_3d\hat{x}_1 + x_4d\hat{x}_3 & x_3d\hat{x}_2 + x_4d\hat{x}_4 \end{pmatrix} = \\
= (dX)\ \hat{X} + X\ (d\hat{X}).
\end{gathered} \tag{16.84}$$

Поэтому верны тождества

$$\hat{\hat{X}} = AX + B \ \Rightarrow \ d\hat{\hat{X}} = AdX, \quad \hat{\hat{X}} = X^{-1} \ \Rightarrow \ d\hat{\hat{X}} = -X^{-1}\, dX\ X^{-1}. \tag{16.85}$$

Доказательство второго таково

$$\begin{gathered} \hat{\hat{X}} X = 1, \ \ (d\hat{\hat{X}})\ X + \hat{\hat{X}}\ (dX) = 0, \\ (d\hat{\hat{X}})\ X + X^{-1}\ (dX) = 0, \ \ d\hat{\hat{X}} = -X^{-1}\ (dX)\ X^{-1}. \end{gathered} \tag{16.86}$$

Тогда получим цепочку тождеств

$$\begin{gathered} \tilde{X}^* + \tilde{X} = 0, \ \ -\tilde{X}^* = \tilde{X}, \\ -(CX+D)^{*-1}(AX+B)^* = (AX+B)(CX+D)^{-1}, \\ -(AX+B)^* = (CX+D)^*(AX+B)(CX+D)^{-1}. \end{gathered} \tag{16.87}$$

Домножим обе части на CdX

$$\begin{gathered} (-X^*A^* - B^*)CdX = (CX+D)^*(AX+B)(CX+D)^{-1}CdX, \\ (-X^*A^*C - B^*C)dX = (CX+D)^*(AX+B)(CX+D)^{-1}CdX, \\ (X^*C^*A + D^*A - E)dX = (CX+D)^*(AX+B)(CX+D)^{-1}CdX, \\ (CX+D)^*AdX - (CX+D)^*(AX+B)(CX+D)^{-1}CdX = dX, \\ A(dX)(CX+D)^{-1} - (AX+B)(CX+D)^{-1}C(dX)(CX+D)^{-1} = \\ = (CX+D)^{*-1}(dX)(CX+D)^{-1}, \\ (d(AX+B))\ (CX+D)^{-1} + (AX+B)d((CX+D)^{-1}) = \\ = (CY+DZ)^{*-1}Z^*(dX)Z(CY+DZ)^{-1}, \\ \tilde{Z}^*\ d\tilde{X}\tilde{Z} = Z^*\ dXZ. \end{gathered} \tag{16.88}$$

Таким образом получается первый инвариант.

16.8.8 Доказательство формул о втором инварианте

Пусть

$$\hat{X} = AX + B, \tag{16.89}$$

или в покомпонентной записи

$$\begin{gathered} \hat{x}_1 = a_1x_1 + a_2x_3 + b_1, \quad \hat{x}_2 = a_1x_2 + a_2x_4 + b_2, \\ \hat{x}_3 = a_3x_1 + a_4x_3 + b_3, \quad \hat{x}_4 = a_3x_2 + a_4x_4 + b_4. \end{gathered} \tag{16.90}$$

Тогда

$$\begin{gathered} \tfrac{\partial}{\partial x_1} = a_1\tfrac{\partial}{\partial \hat{x}_1} + a_3\tfrac{\partial}{\partial \hat{x}_3}, \qquad \tfrac{\partial}{\partial x_2} = a_1\tfrac{\partial}{\partial \hat{x}_2} + a_3\tfrac{\partial}{\partial \hat{x}_4}, \\ \\ \tfrac{\partial}{\partial x_3} = a_2\tfrac{\partial}{\partial \hat{x}_1} + a_4\tfrac{\partial}{\partial \hat{x}_3}, \qquad \tfrac{\partial}{\partial x_4} = a_2\tfrac{\partial}{\partial \hat{x}_2} + a_4\tfrac{\partial}{\partial \hat{x}_4}, \end{gathered} \tag{16.91}$$

или в сокращенной записи

$$\tfrac{\partial}{\partial X} = A^T \tfrac{\partial}{\partial \hat{X}} \quad \Rightarrow \quad \tfrac{\partial}{\partial \hat{X}^T} = \tfrac{\partial}{\partial X^T} A^{-1} \ (det(A) \neq 0). \tag{16.92}$$

Пусть теперь

$$\hat{X} = X^{-1}, \tag{16.93}$$

или в покомпонентной записи

$$\hat{x}_1 = \tfrac{x_4}{x_1 x_4 - x_2 x_3} \quad \hat{x}_2 = -\tfrac{x_2}{x_1 x_4 - x_2 x_3}, \quad \hat{x}_3 = -\tfrac{x_3}{x_1 x_4 - x_2 x_3} \quad \hat{x}_4 = \tfrac{x_1}{x_1 x_4 - x_2 x_3}. \tag{16.94}$$

Тогда

$$\begin{gathered}
\tfrac{\partial}{\partial x_1} = -\tfrac{x_4 x_4}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_1} + \tfrac{x_2 x_4}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_2} + \tfrac{x_3 x_4}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_3} - \tfrac{x_2 x_3}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_4}, \\
\tfrac{\partial}{\partial x_2} = \tfrac{x_4 x_3}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_1} - \tfrac{x_1 x_4}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_2} - \tfrac{x_3 x_3}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_3} + \tfrac{x_1 x_3}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_4}, \\
\tfrac{\partial}{\partial x_3} = \tfrac{x_4 x_2}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_1} - \tfrac{x_2 x_2}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_2} - \tfrac{x_1 x_4}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_3} + \tfrac{x_1 x_2}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_4}, \\
\tfrac{\partial}{\partial x_4} = -\tfrac{x_2 x_3}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_1} + \tfrac{x_1 x_2}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_2} + \tfrac{x_1 x_3}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_3} - \tfrac{x_1 x_1}{(x_1 x_4 - x_2 x_3)^2} \tfrac{\partial}{\partial \hat{x}_4},
\end{gathered} \tag{16.95}$$

или в сокращенной записи

$$\tfrac{\partial}{\partial X} = -X^{-1T} \tfrac{\partial}{\partial \hat{X}} X^{-1T} \quad \Leftrightarrow \quad \tfrac{\partial}{\partial \hat{X}^T} = -X \tfrac{\partial}{\partial X^T} X. \tag{16.96}$$

Предположим, что $det(C) \neq 0$, то

$$\begin{gathered}
\tilde{X} = (AX + B)(CX + D)^{-1} = AC^{-1} + (B - AC^{-1}D)(CX + D)^{-1}, \\
\tfrac{\partial}{\partial \tilde{X}^T} = \tfrac{\partial}{\partial (CX+D)^{T-1}} (B - AC^{-1}D)^{-1} = \\
= -(CX + D) \tfrac{\partial}{\partial (CX+D)^T} (CX + D)(B - AC^{-1}D)^{-1} = \\
= -(CX + D) \tfrac{\partial}{\partial X^T} C^{-1} (CX + D)(B - AC^{-1}D)^{-1} = \\
= -(CX + D) \tfrac{\partial}{\partial X^T} (XC^* + C^{-1}DC^*)(BC^* - AC^{-1}DC^*)^{-1} = \\
= (CX + D) \tfrac{\partial}{\partial X^T} (CX + D)^* (BC^* + AD^*)^{-1} = (CX + D) \tfrac{\partial}{\partial X^T} (CX + D)^*.
\end{gathered} \tag{16.97}$$

Откуда

$$\tilde{Z}^{-1} \frac{\partial}{\partial \tilde{X}^T} \tilde{Z}^{*-1} = Z^{-1} \frac{\partial}{\partial X^T} Z^{*-1}. \tag{16.98}$$

Если $det(C) = 0$, то

$$\tilde{\tilde{X}} = \tilde{X}^{-1} = (CX + D)(AX + B)^{-1}, \quad \tilde{\tilde{Z}} = AY + BZ, \quad \tilde{\tilde{C}} = A, \quad det(\tilde{\tilde{C}}) \neq 0. \tag{16.99}$$

Если $det(A) \neq 0$, то

$$\begin{gathered}
Z^{-1}\tfrac{\partial}{\partial \tilde{X}^T}Z^{*-1} = \tilde{\tilde{Z}}^{-1}\tfrac{\partial}{\partial \tilde{\tilde{X}}^T}\tilde{\tilde{Z}}^{*-1} = \\
= -(AY+BZ)^{-1}\tilde{X}\tfrac{\partial}{\partial \tilde{X}^T}\tilde{X}(AY+BZ)^{*-1} = |\tilde{X}^* = -\tilde{X}| = \\
= (AY+BZ)^{-1}(AY+BZ)(CY+DZ)^{-1}\tfrac{\partial}{\partial \tilde{X}^T}\times \\
\times(CY+DZ)^{*-1}(AY+BZ)^{*}(AY+BZ)^{*-1} = \tilde{Z}^{-1}\tfrac{\partial}{\partial \tilde{X}^T}\tilde{Z}^{*-1}.
\end{gathered} \tag{16.100}$$

Если все-таки $det(A) = 0$, то можно положить

$$\tilde{\tilde{X}} = \tilde{X} + \tilde{D},\quad \tilde{\tilde{\tilde{X}}} = \tilde{\tilde{X}}^{-1},\quad \tilde{\tilde{\tilde{X}}} = (\tilde{\tilde{\tilde{A}}}X + \tilde{\tilde{\tilde{B}}})(\tilde{\tilde{\tilde{C}}}X + \tilde{\tilde{\tilde{D}}})^{-1},\quad \tilde{\tilde{\tilde{C}}} = A + \tilde{D}C,\quad det(\tilde{\tilde{\tilde{C}}}) \neq 0. \tag{16.101}$$

Очевидно, что $\tilde{D}$ всегда можно выбрать так, что $det(\tilde{\tilde{\tilde{C}}}) \neq 0$. Поэтому

$$Z^{-1}\tfrac{\partial}{\partial X^T}Z^{*-1} = \tilde{\tilde{\tilde{Z}}}^{-1}\tfrac{\partial}{\partial \tilde{\tilde{\tilde{X}}}^T}\tilde{\tilde{\tilde{Z}}}^{*-1} = \tilde{\tilde{Z}}^{-1}\tfrac{\partial}{\partial \tilde{\tilde{X}}^T}\tilde{\tilde{Z}}^{*-1} = \tilde{Z}^{-1}\tfrac{\partial}{\partial \tilde{X}^T}\tilde{Z}^{*-1}, \tag{16.102}$$

что и даст второй инвариант.

16.8.9 Доказательство тождества альтернативности для алгебры октав

Докажем тождество (14.95). В случае $P_{\Psi A} = \eta_{\Psi X A}X^X$

$$\begin{gathered}
\tfrac{1}{2}\eta_{(\Lambda\Psi)}{}^{\Theta}\eta_{\Theta\Phi}{}^{\Upsilon} = \eta_{(\Lambda}{}^{AB}P_{\Psi)A}P^{\Theta}{}_{B}\eta_{\Theta CD}P_{\Phi}{}^{C}P^{\Upsilon D} = \\
= (-\eta_{(\Lambda}{}^{AB}\eta_{\Psi)AX} + 2\eta_{(\Lambda}{}^{AB}\eta_{\Psi)(AX)})X^X P^{\Theta}{}_{B}\eta_{\Theta CD}P_{\Phi}{}^{C}P^{\Upsilon D} = \\
= -\tfrac{1}{2}g_{\Lambda\Psi}X^B X^Y\varepsilon_{YBCD}P_{\Phi}{}^{C}P^{\Upsilon D} + \tfrac{1}{4\sqrt{2}}\eta_{(\Psi}{}^{K}{}_{|K}\eta_{\Theta\Phi}{}^{\Upsilon}P^{\Theta}{}_{B|}P_{\Lambda)}{}^{B} = \\
= -\tfrac{1}{2}g_{\Lambda\Psi}\delta_{\Phi}{}^{\Upsilon} + \tfrac{1}{4\sqrt{2}}\eta_{(\Psi}{}^{K}{}_{|K|}\eta_{\Lambda)\Phi}{}^{\Upsilon},
\end{gathered}$$

$$\begin{gathered}
\tfrac{1}{2}\eta_{(\Lambda|\Theta|}{}^{\Upsilon}\eta_{\Psi)\Phi}{}^{\Theta} = \eta_{(\Lambda}{}^{AB}\eta_{\Psi)}{}^{CD}P_{\Theta A}P^{\Upsilon}{}_{B}P_{\Phi C}P^{\Theta}{}_{D} = \\
= (-\eta_{(\Lambda}{}^{AB}\eta_{\Psi)A}{}^{C} + 2\eta_{(\Lambda}{}^{AB}\eta_{\Psi)K}{}^{K}\cdot\tfrac{1}{8}\delta_{A}{}^{C})P^{\Upsilon}{}_{B}P_{\Phi C} = \\
= -\tfrac{1}{2}g_{\Lambda\Psi}\delta_{\Phi}{}^{\Upsilon} + \tfrac{1}{4}\eta_{(\Lambda}{}^{CB}\eta_{\Psi)K}{}^{K}P^{\Upsilon}{}_{B}P_{\Phi C} = -\tfrac{1}{2}g_{\Lambda\Psi}\delta_{\Phi}{}^{\Upsilon} + \tfrac{1}{4\sqrt{2}}\eta_{(\Psi}{}^{K}{}_{|K|}\eta_{\Lambda)\Phi}{}^{\Upsilon},
\end{gathered}$$

$$\begin{gathered}
\tfrac{1}{2}\eta_{\Lambda\Theta}{}^{\Phi}\eta_{(\Upsilon\Psi)}{}^{\Theta} = \eta_{\Lambda}{}^{AB}P_{\Theta A}P^{\Phi}{}_{B}\eta_{(\Psi}{}^{CD}P_{\Upsilon)C}P^{\Theta}{}_{D} = \eta_{\Lambda D}{}^{B}\eta_{(\Psi}{}^{CD}P_{\Upsilon)C}P^{\Phi}{}_{B} = \\
= \eta_{\Lambda D}{}^{B}(-\eta_{(\Psi}{}^{CD}\eta_{\Upsilon)CX}X^X + \tfrac{1}{4}P_{(\Psi}{}^{D}\eta_{\Upsilon)K}{}^{K})P^{\Phi}{}_{B} = -\tfrac{1}{2}g_{\Upsilon\Psi}\delta_{\Lambda}{}^{\Phi} + \tfrac{1}{4\sqrt{2}}\eta_{\Lambda(\Psi}{}^{\Phi}\eta_{\Upsilon)K}{}^{K},
\end{gathered}$$

$$\begin{gathered}
\tfrac{1}{2}\eta_{\Lambda(\Psi}{}^{\Theta}\eta_{|\Theta|\Upsilon)}{}^{\Phi} = \eta_{\Lambda}{}^{AB}P_{(\Psi|A}P^{\Theta}{}_{B}\eta_{\Theta|}{}^{CD}P_{\Upsilon)C}P^{\Phi}{}_{D} = \eta_{\Lambda}{}^{AB}P_{(\Psi|A|}P_{\Upsilon)C}\varepsilon_{XB}{}^{CD}X^X P^{\Phi}{}_{D} = \\
= (\eta_{\Lambda}{}^{CD}X^A - \eta_{\Lambda X}{}^{B}X^X\varepsilon^{A}{}_{B}{}^{CD})P_{(\Psi|A|}P_{\Upsilon)C}P^{\Phi}{}_{D} = -\tfrac{1}{2}g_{\Upsilon\Psi}\delta_{\Lambda}{}^{\Phi} + \tfrac{1}{4\sqrt{2}}\eta_{\Lambda(\Psi}{}^{\Phi}\eta_{\Upsilon)K}{}^{K}.
\end{gathered} \tag{16.103}$$

16.8.10 Доказательство центрального тождества Муфанг

Докажем тождество (14.96). Отметим, что $r^\Phi \eta_{\Phi\Theta}{}^\Omega$ и $\eta_{\Omega\Upsilon}{}^\Gamma r^\Upsilon$ являются ортогональными преобразованиями.

$$\begin{aligned} r^\Phi \eta_{\Phi\Theta}{}^\Omega \eta_{\Lambda\Psi}{}^\Theta \eta_{\Omega\Upsilon}{}^\Gamma r^\Upsilon &= r^\Phi \eta_{\Phi\Lambda}{}^\Theta \eta_{\Theta\Omega}{}^\Gamma \eta_{\Psi\Upsilon}{}^\Omega r^\Upsilon, \\ r^\Phi \eta_{\Phi\Lambda}{}^\Theta \eta_{\Theta\Delta}{}^\Gamma &= r^\Phi \eta_{\Phi\Theta}{}^\Omega \eta_{\Lambda\Psi}{}^\Theta \eta_{\Omega\Upsilon}{}^\Gamma r^\Upsilon \eta^\Psi{}_{\Xi\Delta} r^\Xi. \end{aligned} \tag{16.104}$$

Поскольку $r^\Lambda = \eta^\Lambda{}_{AB} X^A Y^B = P^\Lambda{}_B Y^B$, то

$$\begin{aligned} R_S{}^Q P^\Theta{}_Q \eta_\Theta{}^{CD} &= \underbrace{R_B{}^K}_{[=\varepsilon_B{}^{KMN} X_M Y_N]} \underbrace{P^\Lambda{}_S \eta_\Lambda{}^{AB}}_{[=X^P \varepsilon_{PS}{}^{AB}]} \underbrace{P^\Omega{}_K \eta_\Omega{}^{VD} Y_V}_{[=\varepsilon_K{}^{LDV} X_L Y_V]} \underbrace{P^\Psi{}_A \eta_\Psi{}^{YC} Y_Y}_{[\varepsilon_A{}^{XCY} X_X Y_Y]}, \\ X_X Y_Y \varepsilon_S{}^{QXY} \varepsilon_{RQ}{}^{CD} X^R &= \varepsilon_B{}^{KMN} X_M Y_N X^P \varepsilon_{PS}{}^{AB} \varepsilon_K{}^{LDV} X_L Y_V \varepsilon_A{}^{XCY} X_X Y_Y. \end{aligned} \tag{16.105}$$

1. Правая часть.

$$\begin{gathered} \varepsilon_B{}^{KMN} X_M Y_N X^P \varepsilon_{PS}{}^{AB} \varepsilon_K{}^{LDV} X_L Y_V \varepsilon_A{}^{XCY} X_X Y_Y = \\ = X^P \varepsilon_{PS}{}^{AB} \varepsilon_A{}^{XCY} X_X Y_Y \varepsilon_B{}^{KMN} X_M Y_N \varepsilon_K{}^{LDV} X_L Y_V = \\ = X^P \varepsilon_{PS}{}^{AB} \varepsilon_A{}^{XCY} X_X Y_Y (-\varepsilon^{KMN}{}_B + \delta_B{}^M \varepsilon^{KN}) X_M Y_N \varepsilon_K{}^{LDV} X_L Y_V = \\ = X^P \varepsilon_{PS}{}^{AB} \varepsilon_A{}^{XCY} X_X Y_Y (-\varepsilon^{DVN}{}_B Y_V Y_N + \varepsilon^{NLDV} X_B Y_N Y_V X_L) = \\ = X^P (\varepsilon_{PS} \varepsilon^{AB} + \delta_P{}^A \delta_S{}^B - \delta_P{}^B \delta_S{}^A - \varepsilon^B{}_S{}^A{}_P) \varepsilon_A{}^{XCY} X_X Y_Y \\ (-Y^D Y_B + \delta_B{}^D + X_L Y^L Y^D X_B - X^D X_B) = \\ = (X^S \varepsilon^{BXCY} X_X Y_Y + \delta_S{}^B Y^C - X^B \varepsilon_S{}^{XCY} X_X Y_Y - \varepsilon^B{}_S{}^{CY} Y_Y) \\ (-Y^D Y_B + \delta_B{}^D + X_L Y^L Y^D X_B - X^D X_B) = \\ = -X_S Y^D X_X Y_B Y_Y \varepsilon^{BXCY} + X_S \varepsilon^{DXCY} Y_Y X_X + X_S Y^L X_L Y^D X_B \varepsilon^{BXCY} X_X Y_Y - \\ -X_S X^D X_B \varepsilon^{BXCY} X_X Y_Y - Y^C Y^D Y_S + Y^C \delta_S{}^D + Y^C Y^D X_S X_L Y^L - Y^C X^D X_S + \\ +X^B \varepsilon_S{}^{XCY} X_X Y_Y Y^D Y_B - X^D \varepsilon_S{}^{XCY} X_X Y_Y - Y^L X_L Y^D X_B X^B \varepsilon_S{}^{XCY} X_X Y_Y + \\ +X^B X_B X^D \varepsilon_S{}^{XCY} X_X Y_Y + Y^D Y_B \varepsilon^B{}_S{}^{CY} Y_Y - \varepsilon^D{}_S{}^{CY} Y_Y - \\ -Y^L X_L Y^D X_B \varepsilon^B{}_S{}^{CY} Y_Y + X^D X_B \varepsilon^B{}_S{}^{CY} Y_Y = \\ = -X_S Y^D X_X (\underbrace{Y^X Y^C}_{1} - \varepsilon^{CX}) + X_S \varepsilon^{DXCY} Y_Y X_X + \underbrace{X_S Y^C Y^D Y^L X_L}_{2} - \\ - \underbrace{X_S X^D Y^C}_{3} - \underbrace{Y^C Y^D Y_S}_{4} + Y_C \delta_S{}^D + \underbrace{Y^C Y^D X_S X_L Y^L}_{1} - Y^C X^D X_S + \\ + \underbrace{X^B Y_B Y^D \varepsilon_S{}^{XCY} X_X Y_Y}_{2} - \underbrace{X^D \varepsilon_S{}^{XCY} X_X Y_Y}_{3} - \underbrace{2 X^L Y_L Y^D \varepsilon_S{}^{XCY} X_X Y_Y}_{2} + \\ +2 \underbrace{X^D \varepsilon_S{}^{XCY} X_X Y_Y}_{3} + Y^D (\underbrace{Y_S Y^C}_{4} - \delta_S{}^C) - \varepsilon^D{}_S{}^{CY} Y_Y - \\ - \underbrace{Y^L X_L Y^D X_B \varepsilon^B{}_S{}^{CY} Y_Y}_{1} + \underbrace{X^D X_B \varepsilon^B{}_S{}^{CY} Y_Y}_{3} = \\ = X_S \varepsilon^{DXCY} Y_Y X_X + X_S Y^D X^C - X_S X^D Y^C - \varepsilon^D{}_S{}^{CY} Y_Y + Y^C \delta_S{}^D - Y^D \delta_S{}^C = \\ = X_S \varepsilon^{CDXY} Y_Y X_X - \varepsilon^{CD}{}_S{}^Y Y_Y. \end{gathered} \tag{16.106}$$

2. Левая часть.

$$\begin{aligned}X_X Y_Y \varepsilon_S{}^{QXY} \varepsilon_{RQ}{}^{CD} X^R = (\delta_S{}^X \varepsilon^{QY} - \varepsilon^{XQ}{}_S{}^Y) \varepsilon_{RQ}{}^{CD} X^R X_X Y_Y = \\ = X_S Y^Q X^R \varepsilon_{RQ}{}^{CD} - \varepsilon_S{}^{YCD} Y_Y.\end{aligned} \tag{16.107}$$

Это прямое доказательство. Более простое доказательство основано на формуле (10.1, случай 2).

16.8.11 Доказательство тождеств для оператора A для n=8

Докажем тождество (14.99).

1. Во-первых.

$$\begin{aligned}\eta_\Lambda{}^{KA} \eta_{\Psi K}{}^B \eta^\Lambda{}_{LC} \eta^{\Psi L}{}_D = \varepsilon_{LC}{}^{KA} \varepsilon_K{}^{BL}{}_D = \\ = (\varepsilon_{LC} \varepsilon^{KA} - \delta_L{}^A \delta_C{}^K + \varepsilon^K{}_{LC}{}^A)(\delta_K{}^B \delta_D{}^L - \varepsilon_{KD} \varepsilon^{BL} + \varepsilon_K{}^L{}_D{}^B) = \\ = \varepsilon_{CD} \varepsilon^{AB} - \delta_D{}^A \delta_C{}^B + \varepsilon^B{}_{DC}{}^A - \delta_D{}^A \delta_C{}^B + \varepsilon_{CD} \varepsilon^{AB} - \varepsilon_D{}^B{}_C{}^A + \\ + \varepsilon^A{}_{CD}{}^B - \varepsilon_C{}^A{}_D{}^B + 4\varepsilon_C{}^A{}_D{}^B = 2(\varepsilon_{CD} \varepsilon^{AB} + \delta_C{}^A \delta_D{}^B - \delta_D{}^A \delta_C{}^B).\end{aligned} \tag{16.108}$$

2. Во-вторых.

$$\begin{aligned}\eta_\Lambda{}^{KA} \eta_{\Psi K}{}^B \eta^\Lambda{}_{CL} \eta^\Psi{}_D{}^L = \varepsilon_{CL}{}^{KA} \varepsilon_K{}^B{}_D{}^L = (\delta_C{}^K \delta_L{}^A - \varepsilon^K{}_{LC}{}^A)(\varepsilon_{KD} \varepsilon^{BL} - \varepsilon_K{}^L{}_D{}^B) = \\ = \varepsilon_{CD} \varepsilon^{AB} - \varepsilon_D{}^B{}_C{}^A - \varepsilon_C{}^A{}_D{}^B + 4\varepsilon_C{}^A{}_D{}^B = \varepsilon_{CD} \varepsilon^{AB} + 2\varepsilon_C{}^A{}_D{}^B.\end{aligned} \tag{16.109}$$

3. В-третьих.

   На основании тождества (8.27)

$$\eta_\Lambda{}^{AB} \eta_{\Psi AD} \eta^{\Omega CD} \eta^\Theta{}_{CB} = 2(g_{\Lambda\Psi} g^{\Omega\Theta} + \delta_\Lambda{}^\Theta \delta_\Psi{}^\Omega - \delta_\Psi{}^\Theta \delta_\Lambda{}^\Omega), \tag{16.110}$$

   тогда получим

$$\begin{aligned}\eta_{[\Lambda}{}^C{}_{|A|} \eta_{\Psi]CB} \eta_{\Upsilon K}{}^A \eta_\Omega{}^{KB} = 4 g_{\Upsilon[\Lambda} g_{\Psi]\Omega}, \\ \eta_{[\Lambda}{}^C{}_{|A|} \eta_{\Psi]CB} \eta_\Upsilon{}^A{}_K \eta_\Omega{}^{BK} = 4 g_{\Upsilon[\Lambda} g_{\Psi]\Omega} + 8\eta_{[\Upsilon} g_{\Omega][\Lambda} \eta_{\Psi]}, \\ \eta_{[\Lambda|A|}{}^C \eta_{\Psi]BC} \eta_\Upsilon{}^A{}_K \eta_\Omega{}^{BK} = 4 g_{\Upsilon[\Lambda} g_{\Psi]\Omega}, \\ \eta_{[\Lambda|A|}{}^C \eta_{\Psi]BC} \eta_{\Upsilon K}{}^A \eta_\Omega{}^{KB} = 4 g_{\Upsilon[\Lambda} g_{\Psi]\Omega} + 8\eta_{[\Upsilon} g_{\Omega][\Lambda} \eta_{\Psi]}.\end{aligned} \tag{16.111}$$

   При этом

$$\begin{aligned}\eta_{[\Lambda}{}^C{}_{|A|} \eta_{\Psi]CB} \eta_{\Upsilon K}{}^A \eta_\Omega{}^{KB} = 8 g_{\Upsilon[\Lambda} g_{\Psi]\Omega} + \eta_\Upsilon{}^C{}_A \eta_{[\Psi|CB|} \eta_{\Lambda]}{}^{KB} \eta_{\Omega K}{}^A = \\ = 8 g_{\Upsilon[\Lambda} g_{\Psi]\Omega} + (\eta_{[\Lambda}{}^{BK} \eta_{\Psi]BC} + 2\eta_{[\Lambda}{}^K{}_{|C|} \eta_{\Psi]})(\eta_{[\Upsilon|A|}{}^C \eta_{\Omega]}{}^A{}_K + 2\eta_{[\Omega|K|}{}^C \eta_{\Upsilon]}) = \\ = 8 g_{\Upsilon[\Lambda} g_{\Psi]\Omega} - \eta_{[\Lambda}{}^C{}_{|A|} \eta_{\Psi]CB} \eta_{\Upsilon K}{}^A \eta_\Omega{}^{KB} - 2\eta_{[\Psi}{}^K{}_{|C|} \eta_{\Lambda]} \eta_{[\Upsilon|A|}{}^C \eta_{\Omega]}{}^A{}_K + \\ + 2\eta_{[\Omega|K|}{}^C \eta_{\Upsilon]} \eta_{[\Lambda}{}^{BK} \eta_{\Psi]BC} - 16\eta_{[\Lambda} g_{\Psi][\Omega} \eta_{\Upsilon]}.\end{aligned} \tag{16.112}$$

Литература

[1] Андреев К. В. Спинорный формализм и геометрия шестимерных римановых пространств. Кандидатская диссертация, Уфа, 1997. [K.V. Andreev. Spinor formalism and the geometry of six-dimensional Riemannian spaces / Ph. D. Thesis, Ufa, 1997], arXiv:1204.0194v1.

[2] Андреев К. В. О структуре тензора кривизны 6-мерных римановых пространств. Вестик Башкирского унверситета, (1996) №2, с. 44-47. [K. V. Andreev. On the structure of the curvature tensor of 6-dimensional Riemannian spaces. Vestnik Bashkirskogo Universiteta, (1996) №2, 44-47].

[3] Андреев К. В. О спинорном формализме при n=6. Известия Высших учебных заведений, 2001, №1 (№464), 11-23.[K. V. Andreev. On spinor formalism for n=6. Izvestiya Vysshikh Uchebnykh Zavedeni., Matematika, 2001, №1 (№464), 11-23. [(The article is freely available online at the Mathnet URL:`http://mi.mathnet.ru/eng/ivm838`, the two figures of the article are missing in the electronic version, however). English translation: K. V. Andreyev: On spinor formalism for n = 6. Russian Mathematics (Iz. VUZ) 45:1(2001)9-20.]

[4] Андреев К. В. О структурных константах алгебры октав. Уравнение Клиффорда , Известия вузов, Математика, 2001,№3, 3–6. [K. V. Andreev, Structure constants of the algebra of octaves. The Clifford equation, Izv. Vyssh. Uchebn. Zaved. Mat., 2001, №3, 1-4, Mathnet URL:`www.ksu.ru/journals/izv_vuz/arch/2001/03/01-3.PDF`]

[5] Андреев К. В. О твисторах 6-мерного пространства. ВИНИТИ -2469-В-98, август 1998. 23 с. [K.V. Andreev. On the twistors of six-dimensional space. VINITI-2469-V-98, Aug 1998. 23pp. Paper deponed on Aug 3, 1998 at VINITI (Moscow), ref. №2469-V 98]

[6] Андреев К. В. Принцип тройственности для двух квадрик. ВИНИТИ -2470-В-98, август 1998. 19 с. [K.V. Andreev. Triality principle for two quadrics. VINITI-2470-V-98, Aug 1998. 19pp. Paper deponed on Aug 3, 1998 at VINITI (Moscow), ref. №2470-V 98.]

[7] Андреев К. В. О внутренних геометриях многообразия плоских образующих 6-мерной квадрики, Изв. вузов. Матем., 1998,№6, 3-8. [K. V. Andreev. On intrinsic geometries of the manifold of plane generators of a 6-dimensional quadric, Izv. Vyssh. Uchebn. Zaved. Mat., 1998, №6, 1-6, Mathnet URL:`http://www.mathnet.ru/links/31e5d8f5c91e320943c14afce86b45aa/ivm436.pdf`]

[8] Баэз Джон С. Октонионы. Гиперкомплексные числа в геометрии и физике. №1(5), Vol 3 (2006), с.120-177. [John C. Baez The Octonions. arXiv:math.RA/0105155v4]

[9] Билялов Р.Ф., Никитин Б.С. Спиноры в произвольных реперах. Ковариантная производная и производная Ли спиноров. Известия Высших учебных заведений, 1998, N 6 (N 433), 9-19. [R. F. Bilyalov, B. S. Nikitin. Spinors in arbitrary frames. The covariant derivative and the Lie derivative of spinors. Izv. Vyssh. Uchebn. Zaved. Mat., 1998, №. 6, 7–15.]

[10] Берже М. Геометрия т.1,2. Москва, Мир, 1984. [Berger, Marcel: Geometry. II. Translated from the French by M. Cole and S. Levy. Universitext. Springer-Verlag, Berlin, 1987. Berger, Marcel: Geometry. I. Translated from the French by M. Cole and S. Levy. Universitext. Springer-Verlag, Berlin, 1987.]

[11] Бухдаль Х.А. H.A. Buchdahl: On the calculus of for-spinors. Proceedings of the Royal Society of London, Series A, Mathematical and Physical Sciences 303(1968)355-378.

[12] Дитмар С. Dietmar Salamon: Spin Geometry and Seiberg-Witten invariants. Warwick, 1996.

[13] Клотц Ф.С. F. S. Klotz. Twistors and the conformal group. Journal of Mathematical Phisics, 15(1974) 2242-2247.

[14] Котельников А.П. Винтовое счисление и некоторые приложения его к геометрии механики. Казань, 1895 г. [Screw calculation and some of its applications to geometry, mechanics. Kazan, 1895.]

[15] Нейфельд Э.Г. Об инволюциях в комплексных пространствах. ТГС, Казань, 1989, Выпуск 19, с. 71-82. [E. G. Neifel'd. On the involutions in complex spaces. Tr. Geom. Semin, Kazan, 1989, № 19, pp. 71-82.]

[16] Нейфельд Э.Г. Нормализация комплексных грассманианов и квадрик. ТГС, Казань, 1990, Выпуск 20, с. 58-69 [E. G. Neifel'd. Normalization of complex Grassmannians and quadrics. Tr. Geom. Semin, Kazan, 1990, Nø 20, pp. 58-69.]

[17] Норден А.П. О комплексном представлении тензоров пространства Лоренца. Известия ВУЗов, Казань, 1959, т. 8, № 1, с. 156-164, Mathnet URL:`http://www.mathnet.ru/links/456d047e23563a194e6ea7d5c013a792/ivm2415.pdf`.

[18] Постников М.М. Лекции по геометрии. V семестр. Группы и алгебры Ли. Москва, Наука, 1982 г.

[19] Розенфельд Б.А. Проективная геометрия как метрическая геометрия. Труды семинара по векторному и тензорному анализу с их приложениями к геометрии, механике и физике.- М., 1950.-Т8.-с. 328-354. [B. A. Rosenfeld. Projective geometry as the metric geometry. Proceedings of the Seminar on Vector and tensor analysis with their applications.- Moscow, 1950.-v8.-p. 328-354.]

[20] Розенфельд Б.А. Неевклидовы геометрии. Москва, ГИТО, 1955. [B. A. Rosenfeld. Non-euclidian geometry. Moscow, Gosudarstv. Izdat. Tehn., 1955]

[21] Энтони Зи Квантовая теория в двух словах. РХД, Москва, Ижевск, 2009. [A. Zee. Quantum field theory in a nutshell. Princeton university press, 2003]

[22] Пенроуз Р. Риндлер В. Спиноры и пространство-время. Т.1, Мир, Москва, 1987. Спиноры и пространство-время. Т.2, Мир, Москва, 1988.

[23] Синцов Д.М. Теория коннексов в пространстве в связи с теорией дифференциальных уравнений в частных производных первого порядка. Казань, 1894 г. [Theory of connex in space in relation to the theory of partial differential equations of first order. Kazan. 1894]

[24] Степановский Ю.П. Спиноры 6-мерного пространства и их применение к описанию поляризованных частиц со спином 1/2. Проблемы Ядерной Физики и Космических Лучей, Харьков, 4(1976)9-21. [Yu. P. Stepanovskii]: Spinors of a sixdimensional space and their application to the description of polarized particles with spin 1/2. Problems of Nuclear Physics and Cosmic Rays, Kharkiv, 4(1976)9-21.

[25] Степановський Ю.П. Алгебра матриць Дірака у шестімірному вигляді. Український Фізичний Журнал. Київ, т. XI, 8(1968) 813-824. [Yu. P. Stepanovskii: Algebra of Dirac matrices in six-dimensional form. Ukrainian Journal of Physics, Kiev, v.XI, 8(1968) 813-824].

[26] Схоутен Я.А. J.A. Schouten, J. Haantjes: Konforme Feldtheorie II; R_6 und Spinuraum [Conformal field theory II and spin space]. Annali della R.Scuola Normale Superiore di Piza, Scienze Fisiche e Matematiche,2. Series, 4(1935)175-189.

[27] Хьюзмоллер Д. Расслоенные пространства. Москва, Мир, 1970 г. [Dale Husemoller. Fibre bundles. McGraw-Hill Book Company,1966]

[28] К. Шарнхорст, Ж.-В. ван Хольтен K. Scharnhorst, J.-W. van Holten: Nonlinear Bogolyubov-Valatin transformations: 2 modes. NIKHEF preprint NIKHEF/2010-005, arXiv:1002.2737v3, 114 pp.

17 Добавление

Рассмотрим вложение $\mathbb{R}^4_{(1,3)} \subset \mathbb{C}^4$ с операторами [5]

$$\bar{H}_i{}^{\Lambda'} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -i \end{pmatrix}, \quad H^i{}_\Lambda = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -i \end{pmatrix}, \quad S_\Lambda{}^{\Lambda'} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}. \tag{17.1}$$

Для значений комплексных операторов из алгоритма 14.1 метрика имеет вид

$$g_{\Lambda\Psi} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad g_{ij} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}. \tag{17.2}$$

Тогда определение (14.182) станет таким

$$(P^*_K)_i{}^A := \tfrac{1+i}{4} \begin{pmatrix} 0 & p_1 - p_2 \\ 0 & 1 - p_1 p_2 \\ p_1 - p_2 & 0 \\ i(1 - p_1 p_2) & 0 \end{pmatrix}, \quad (P_K)^i{}_A := \tfrac{1-i}{4} \begin{pmatrix} 0 & p_1 - p_2 \\ 0 & 1 - p_1 p_2 \\ p_1 - p_2 & 0 \\ -i(1 - p_1 p_2) & 0 \end{pmatrix}. \tag{17.3}$$

Посколькy$x^i := (P_2)^i{}_A (X_2)^A + (P_3)^i{}_A (X_3)^A$ то из (6.33), (14.84) и следствия (8.3) следует, что $(X_K)^A := x^i (P^*_K)_i{}^A$, $(X^*_K)_A := x^i (P_K)_{iA}$, а из (6.33), (14.84), следствия (8.3) и соотношения $(M_K)^{iA}(M_K)_{iA} = 0$, $(M^*_K)^{iA}(M^*_K)_{iA} = 0$ (смотри (14.181)) следует, что $(X^*_2)_C = (P_2)^i{}_C (P_3)_{iB} (X_3)^B$. Поэтому мы можем построить взаимно однозначное соответствие $(a, b, ... = \overline{1,4})$

$$x^i \cdot \underbrace{((P^*_2)_i{}^A, (P^*_3)_i{}^B)}_{:=P_i{}^a} = \underbrace{((X_2)^A, (X_3)^B)}_{:=X^a}, \quad x^i \cdot \underbrace{((P_2)_{iA}, (P_3)_{iB})}_{:=P_{ia}} = \underbrace{((X^*_2)_A, (X^*_3)_B)}_{:=X_a}. \tag{17.4}$$

---

[5] Этот пример не вошел в опубликованную статью.

Тогда на спин-парах $X^a = ((X_2)^A, (X_3)^B)$ индуцируется метрика и инволюция вида

$$\begin{gathered}
\varepsilon_{ab} = \begin{pmatrix} 0 & (P_2)^i{}_A (P_3)_{iB} \\ (P_3)^i{}_C (P_2)_{iD} & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & i \\ -i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{pmatrix}, \\
S_a{}^{b'} = \begin{pmatrix} (P_2)^i{}_A (\bar{P}^*_2)_i{}^{C'} & (P_2)^i{}_A (\bar{P}^*_3)_i{}^{D'} \\ (P_3)^i{}_B (\bar{P}^*_2)_i{}^{C'} & (P_3)^i{}_A (\bar{P}^*_3)_i{}^{D'} \end{pmatrix} = \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & -i \end{pmatrix}.
\end{gathered} \tag{17.5}$$

Поэтому возможно построить на спин-парах лиевские аналоги

$$L_x(Y)^a = x^i \nabla_i Y^a - P^j{}_b Y^b P_i^a \nabla_j x^i, \quad L_X(Y)^a = X^c \nabla_c Y^a - Y^c \nabla_c X^a. \tag{17.6}$$

Заметим, что разложение $x^i := (P_2)^i{}_A (X_2)^A + (P_3)^i{}_A (X_3)^A$ является разложением по двум изотропным векторам. А для таких векторов лиевские аналоги были построены в [22, т. 2, с. 127].

Ограниченное преобразование Лоренца переводит вектор x^i в вектор $\tilde{x}^i$. Это индуцирует спинорные преобразования $(X_K)^A \longmapsto (\tilde{X}_K)^A$. Они воздействуют на оператор $(P^*_K)_i{}^A$, $(P_K)^i{}_A$, или более точно, на управляющий спинор $X^{\tilde{\tilde{A}}}$ из определения (11.59). В свою очередь, это индуцирует преобразования на спин-парах $X^a = ((X_2)^A, (X_3)^B) \longmapsto \tilde{X}^a$.

Contents/Содержание

I English edition

1 Introduction 5

2 Clifford algebra 6

3 Complex and real representations 9

4 Real inclusion. Involution 14

5 Elementary transformations of the orthogonal group 16

6 Clifford equation. Double covering 24

7 Quaternions 35

8 Particular solutions of the Clifford equation 37

9 On the structural constants of the non-division algebra. Sedenions 47

10 Infinitesimal transformations 52

11 Complex and real connections 53

12 On the classification of tensors with the symmetries of the Riemann curvature tensor. Curvature spinors 61

13 Twistor equation 65

14 Spinor formalism for n=6 and n=8 68

- 14.1 General isomorphisms 68
- 14.2 On the classification of tensors with the symmetries of the Riemann curvature tensor for n=6. Curvature spinors 69
- 14.3 Geometric representation of a twistor in $\mathbb{R}^6_{(2,4)}$ 72
- 14.4 Structure constants of the octonion algebra. Spinor formalism for n = 8 77
- 14.5 Geometry of the inductive step from n=6 to n=8 79
  - 14.5.1 Rosenfeld null-pair 79
  - 14.5.2 Construction of the quadrics $\mathbb{C}Q_6$ and $\mathbb{C}\tilde{Q}_6$ 79

14.6 Correspondence $\mathbb{C}Q_6 \longmapsto \mathbb{C}\tilde{Q}_6$ 80
  14.6.1 Geometry of the transition to the connecting operators ${\eta_\Lambda}^{KL}$ for n $= 8$. 83
14.7 Correspondence $\mathbb{C}\tilde{Q}_6 \longmapsto \mathbb{C}Q_6$ 87
  14.7.1 On the Majorana spinors. 89
  14.7.2 Theorem on two quadrics. 91
14.8 Explicit construction of the Lie spin-operators for n=4. 93

15 Acknowledgments **98**

16 Appendix **99**
16.1 Proof of the metric tensor and involution relations 99
16.2 Proof of the connecting operator relations 101
16.3 Proof of the infinitesimal transformation relations 103
16.4 Proof of the Lie bracket identity 104
16.5 Proof of the curvature tensor relations 105
16.6 Proof of the auxiliary identities for the theorem on the spin analogs of the curvature tensor 105
16.7 Proof of the twistor equation relations 107
16.8 Proof of the identities for n $= 6$ and n=8 109
  16.8.1 Proof of the fundamental identities for n $= 6$ 109
  16.8.2 Proof of the 4-vector identities 110
  16.8.3 Proof of the 6-vector identities 110
  16.8.4 Proof of the Bianchi identity 112
  16.8.5 Proof of the identities related to the Weyl tensor 112
  16.8.6 Proof of the relations associated with the metric induced in the cross section of the cone K_6 113
  16.8.7 Proof of the first invariant relations 114
  16.8.8 Proof of the second invariant relations 116
  16.8.9 Proof of the octave alternative identity 117
  16.8.10 Proof of the central Moufang identity 118
  16.8.11 Proof of the A-operator identities for n $= 8$ 119

II Русская редакция

1 Введение **128(5)**

2 Алгебра Клиффорда **129(6)**

3 Комплексная и вещественные реализации **133(10)**

4 Вещественные вложения. Инволюции **138(15)**

5 Элементарные преобразования ортогональной группы **140(17)**

6 Уравнение Клиффорда. Двулистное накрытие **150(27)**

7 Кватернионы **162(39)**

8 Частные решения уравнения Клиффорда **166(43)**

9 О структурных константах алгебр без деления. Седенионы **178(54)**

10 Инфинитизимальные преобразования **183(59)**

11 Комплексная и вещественная связности **184(60)**

12 О классификации тензоров с симметриями тензора кривизны. Спиноры кривизны **193(69)**

13 Твисторное уравнение **197(73)**

14 Спинорный формализм для n=6 и n=8 **200(76)**

14.1 Основные изоморфизмы . . . . 200(76)

14.2 О классификации тензоров с симметриями тензора кривизны для n=6. Спиноры кривизны . . . . 202(78)

14.3 Геометрическое представление твистора в $\mathbb{R}^6_{(2,4)}$ . . . . 205(81)

14.4 О структурных константах алгебры октонионов. Спинорный формализм для n = 8 . . . . 211(87)

14.5 Геометрия индуктивного перехода от n=6 к n=8 . . . . 213(89)

14.5.1 Нуль-пары Розенфельда . . . . 213(89)

14.5.2 Построение квадрик $\mathbb{C}Q_6$ и $\mathbb{C}\tilde{Q}_6$ . . . . 214(90)

14.5.3 Соответствие $\mathbb{C}Q_6 \longmapsto \mathbb{C}\tilde{Q}_6$ . . . . 215(91)

14.5.4 Геометрия перехода к связующим операторам ${\eta_\Lambda}^{KL}$ для n=8 . . . . . 218(94)
14.5.5 Соответствие $\mathbb{C}\tilde{Q}_6 \longmapsto \mathbb{C}Q_6$ . . . . 223(99)
14.5.6 О майорановских спинорах. . . . . 225(101)
14.5.7 Теорема о двух квадриках. . . . . 227(103)
14.6 Явное построение аналогов операторов Ли при n=4. . . . . 229(105)

15 Благодарности **234(110)**

16 Приложение **235(111)**
16.1 Доказательство формул о метрических тензорах и инволюции . . . . 235(111)
16.2 Доказательство формул о связующих операторах . . . . 237(113)
16.3 Доказательство формул об инфинитезимальных преобразованиях . . . . . . 239(115)
16.4 Доказательство тождества о скобке Ли . . . . 240(116)
16.5 Доказательство формул о тензоре кривизны . . . . 241(117)
16.6 Доказательство вспомогательных тождеств для теоремы о спинорных аналогах тензора кривизны . . . . 241(117)
16.7 Доказательство формул о твисторном уравнении. . . . . 244(120)
16.8 Доказательство тождеств для n=6 . . . . 246(122)
16.8.1 Доказательство основных тождеств для n=6 . . . . 246(122)
16.8.2 Доказательство тождеств о 4-векторе . . . . 247(123)
16.8.3 Доказательство формул о 6-векторе . . . . 247(123)
16.8.4 Доказательство тождества Бианки . . . . 249(125)
16.8.5 Доказательство тождеств, касающихся тензора Вейля. . . . . 249(125)
16.8.6 Доказательство формул, связанных с метрикой, индуцированной в сечении конуса K_6 . . . . 250(126)
16.8.7 Доказательство формул о первом инварианте . . . . 251(127)
16.8.8 Доказательство формул о втором инварианте . . . . 253(129)
16.8.9 Доказательство тождества альтернативности для алгебры октав . . . 255(131)
16.8.10 Доказательство центрального тождества Муфанг . . . . 256(132)
16.8.11 Доказательство тождеств для оператора A для n=8 . . . . 257(133)

17 Добавление **262**

Tables/Список таблиц

1 The matrix form of the tensor S for the real inclusions of some 2-dimensional spaces in the complex space for $r^i = (T, Z)$. 45
2 The matrix form of the tensor S for the real inclusions of some 4-dimensional spaces in the complex space for $r^i = (T, X, Y, Z)$. 45
3 The matrix form of the tensor S for the real inclusions of some 6-dimensional spaces in the complex space for $r^i = (T, X, V, W, Y, Z)$. 46
1 Вид матрицы тензора S для соответствующих действительных вложений 2-мерных пространств в комплексное для $r^i = (T, Z)$. . . . . 176(52)
2 Вид матрицы тензора S для соответствующих действительных вложений 4-мерных пространств в комплексное для $r^i = (T, X, Y, Z)$. . . . . 176(52)
3 Вид матрицы тензора S для соответствующих действительных вложений 6-мерных пространств в комплексное для $r^i = (T, X, V, W, Y, Z)$. . . . . 177(53)

Figures/Список рисунков

1 Correspondence $\forall CP_2 \subset CP_3 \leftrightarrow R$ 82
2 Correspondence $CP_3 \supset CP_1 \leftrightarrow CP_1 \subset K_6$ 82
1 Соответствие $\forall CP_2 \subset CP_3 \leftrightarrow R$ . . . . 217(93)
2 Соответствие $CP_3 \supset CP_1 \leftrightarrow CP_1 \subset K_6$ . . . . 217(93)